

QUANTIZATION OF GAUGE FIELDS, GRAPH POLYNOMIALS AND GRAPH HOMOLOGY*

DIRK KREIMER, MATTHIAS SARS, AND WALTER D. VAN SUIJLEKOM

ABSTRACT. We review quantization of gauge fields using algebraic properties of 3-regular graphs. We derive the Feynman integrand at n loops for a non-abelian gauge theory quantized in a covariant gauge from scalar integrands for connected 3-regular graphs, obtained from the two Symanzik polynomials.

The transition to the full gauge theory amplitude is obtained by the use of a third, new, graph polynomial, the corolla polynomial.

This implies effectively a covariant quantization without ghosts, where all the relevant signs of the ghost sector are incorporated in a double complex furnished by the corolla polynomial —we call it cycle homology— and by graph homology.

1. INTRODUCTION

Feynman rules for scalar field theories reveal astonishing connections to algebraic geometry. These are most easily seen upon the use of parametric representations, which furnish renormalized integrands suitable to be analysed in mathematical terms. Crucial are here the two Symanzik polynomials, which underlie the parametric representation [1, 2, 3, 4, 5, 6, 7].

If one wants to generalize this approach into the realm of gauge theories, a pedestrian approach would be to turn every tensor integral appearing from any single graph into the parametric representation, creating a bewildering number of tensor integrals for each and any contributing graph in a non-abelian gauge theory. While effective on-shell methods have been suggested as an alternative and indeed succeeded at sufficiently low loop orders [8, 9], the question how the transition from scalar to gauge field theory can be formulated mathematically remained open.

Here we establish an answer and suggest a more succinct and, we believe, more elegant approach. It allows us to obtain the renormalized integrand of a n -loop scattering amplitude in a gauge theory from the use of the scalar amplitude with cubic interaction and through one further graph polynomial, beyond the use of the two Symanzik polynomials, in one go.

We eliminate thereby the need to introduce ghosts in the context of covariant quantization and replace their use by the use of the corolla polynomial. This polynomial, which incorporates all the necessary signs of closed ghost loops, is still a strictly positive polynomial. This property is maintained in the pure Yang–Mills sector. We also generalize it to enable the inclusion of fermions. Positivity then depends on the representation of the gauge group which we choose for fermionic matter fields.

We believe our approach is useful as it is an approach to gauge theory amplitudes which does not rely on on-shell recursion relations and methods of cut-reconstructibility. This has become the standard approach to gauge theory amplitudes in particular in the context of $N = 4$ supersymmetric theories in recent years. It nevertheless has shortcomings with respect to the consideration of loop amplitudes, and seems rather restricted to the context of a planar graphs, preferably to be considered in a theory with a vanishing β -function.

Here, we offer an approach which is based on a mere study of the two Kirchhoff polynomials, the corolla polynomial of [10], the combinatorics of parametric renormalization [17], and nothing else to obtain the renormalized integrand of a generic gauge theory amplitude from the amplitude of a scalar field theory with 3-valent vertices only.

In this paper, we introduce our approach. Comparison of graph- and cycle homology with BRST homology, an interpretation of Slavnov–Taylor identities through corolla differentials, and computations of gauge theory

This work is supported by an Alexander von Humboldt Professorship of the Alexander von Humboldt Foundation and the BMBF for Dirk Kreimer.

amplitudes will be presented in future work, as will be a discussion of gravity Feynman rules from this approach.

Acknowledgments. M.S. thanks Erik Panzer for many helpful discussions. D.K. is grateful to David Broadhurst, Francis Brown, Spencer Bloch, Erik Panzer, Oliver Schnetz and Karen Yeats for shared collaborations and insights into the topics of this work.

1.1. Results. We prove that the Feynman integrand at any finite loop order n in gauge theory can be obtained from the study of two different complexes: graph homology and cycle homology, acting on 3-regular connected scalar graphs.

Both complexes are reflected in the analytic structure of the Feynman integrand for an amplitude: residues in parametric space along co-dimension k hypersurfaces biject with the integrands of graphs having k distinct 4-valent vertices and thus reflect graph homology, while cycle homology pairs with the structure of the corolla polynomial defined below, see also [10].

In fact, we prove that there exists a corolla differential D_Γ such that the Feynman integrand I_Γ for connected 3-regular graphs Γ ,

$$(1) \quad I_\Gamma = \frac{e^{-\frac{|N_\Gamma|_{\text{pr}}}{\psi_\Gamma}}}{\psi_\Gamma^2},$$

gives rise, when summed over connected graphs Γ , to the total gauge theory amplitude, using $D_\Gamma I_\Gamma$. The differential D_Γ arises from the corolla polynomial C_Γ , which is a graph polynomial based on half-edge variables, upon replacing each half-edge variable by a suitable differential operator assigned to any half-edge.¹

In a gauge theory, we must deal with the simultaneous requirements of unitarity of the scattering matrix S and of covariance of fields which we assume to be representatives of the Poincaré group. These two requirements severely restrict possible Feynman rules [11, 12, 13]. In particular, unphysical degrees of freedom must be eliminated from observable amplitudes.

For the 3- and 4-valent vertices of gauge boson interactions this requires that those vertices are not independent [14]. This leads us to graph homology, as shrinking an edge between two 3-valent vertices gives a 4-valent vertex.

More precisely, let e be an edge connecting two 3-gluon vertices in a graph Γ , χ_e be the operator which shrinks edge e , and we extend χ_e to zero when acting between any other two vertices. Let $\sum_e \chi_e$ sum over all internal edges. Let S , with $S^2 = 0$, be the generalized graph homology operator introduced below. Then, for a gauge theory amplitude r :

Theorem 1.1. *Let $X_{0\chi;j}^{r,n}$ be the sum of all 3-regular connected graphs, with j ghost loops, and with external legs determined by r and loop number n , weighted by colour and symmetry, let $X_{/\chi;j}^{r,n}$ be the same allowing for 3- and 4-valent vertices. We have*

$$\begin{aligned} i) : e^{\chi} X_{0\chi;j}^{r,n} &= X_{/\chi;j}^{r,n}, \\ ii) : S e^{\chi} X_{0\chi;j}^{r,n} &= 0. \end{aligned}$$

This theorem ensures that 3- and 4-valent vertices match to fulfil simultaneously the requirement of relativistic field theory and S -matrix theory.

These two requirements also demand that unphysical degrees of freedom propagating in closed loops do cancel.

Let δ_\pm^C be the operator which marks a cycle C through 3-valent vertices and unmarked edges, extend δ_\pm^C to zero on any other cycle. Let $\sum_C \delta_\pm^C$ sum over all cycles. Let T , with $T^2 = 0$, be the generalized cycle homology operator introduced below. Then:

Theorem 1.2. *Let $X_{j\chi;0}^{r,n}$ be the sum of all connected graphs with j 4-vertices contributing to amplitude r and loop number n and no ghost loops, weighted by colour and symmetry, $X_{j\chi;j}^{r,n}$ be the same allowing for*

¹Equation (1) above needs a correction for quadratic sub-divergences, which will be provided in detail later on

any possible number of ghost loops. We have

$$\begin{aligned} i) : e^{\delta+} X_{j\mathbb{X};0}^{r,n} &= X_{j\mathbb{X};/\circ}^{r,n}, \\ ii) : T e^{\delta+} X_{j\mathbb{X};0}^{r,n} &= 0. \end{aligned}$$

These two operations are compatible:

Theorem 1.3. i) We have $[s, t] = [S, T] = 0$ and

$$T e^{\delta++\chi+} X_{0\mathbb{X};0}^{r,n} = 0, \quad S e^{\delta++\chi+} X_{0\mathbb{X};0}^{r,n} = 0.$$

ii) Together, they generate the whole gauge theory amplitude from 3-regular graphs:

$$e^{\delta++\chi+} X_{0\mathbb{X};0}^{r,n} = X_{/\mathbb{X};/\circ}^{r,n} =: X^{r,n}.$$

Remark 1.4. $X^{r,n}$ is the only non-trivial element in the cycle and graph homology which we here construct. It will be an object of future study.

Finally, we get the Feynman integrands in the unrenormalized and renormalized case for a gauge theory amplitude r from 3-regular connected graphs of scalar fields.

Theorem 1.5. The full Yang–Mills amplitude \bar{U}_Γ for a graph Γ can be obtained by acting with a corolla differential operator (see below) on the scalar integrand $U_\Gamma(\{\xi_e\})$ for Γ , setting the edge momenta $\xi_e = 0$ afterwards.

Moreover, \bar{U}_Γ gives rise to a differential form $J_\Gamma^{\bar{U}_\Gamma}$ and there exists a vector H_Γ such that the unrenormalized Feynman integrand for the sum of all Feynman graphs contributing to the connected k -loop amplitude r is

$$\Phi(X^{r,k}) = \sum_{|\Gamma|=k, \text{res}(\Gamma)=r} \frac{\text{colour}(\gamma)}{\text{sym}(\Gamma)} \int (H_\Gamma \cdot J_\Gamma^{\bar{U}_\Gamma}),$$

The renormalized analogue is given by writing \bar{U}_Γ^R instead of \bar{U}_Γ .

1.2. Organization of the paper. The next section gives a detailed account of our graph-theoretic notions including graph homology. Then, we turn to the structure of Feynman rules for scalar fields in the parametric representation, including renormalization as rigorously detailed for parametric representations in [17]. The peculiar situation in gauge theory is discussed in the fourth section, and a review of gauge theory from the viewpoint of Hochschild cocycles and the corresponding combinatorial Green functions is provided in the fifth. The sixth section combines these results with the corolla polynomial and the corolla differentials, culminating in Theorem 1.5 above. Short conclusions finish the paper.

2. GRAPHS

We first define the necessary graph theoretic notions.

2.1. Vertices, edges, half-edges. We consider connected graphs with labelled edges and vertices. We consider graphs as elements of a free commutative \mathbb{Q} -algebra H , which is graded as a vectorspace by the first Betti number, the number of algebraically independent cycles in a graph.

For a graph Γ , we let $\Gamma^{[1]} = \Gamma_E^{[1]} \cup \Gamma_I^{[1]} \equiv E^\Gamma = E_E^\Gamma \cup E_I^\Gamma$ be the set of its external and internal edges and let $\Gamma^{[0]} \equiv V^\Gamma$ be the set of its vertices.

We do not allow for internal edges which form self-loops (tadpoles): every internal edge can be considered as a pair of two distinct vertices.²

For a vertex $v \in V^\Gamma$, let $n(v)$ be the set of edges adjacent to v , and its cardinality $|n(v)|$ be the valence of v . For an edge $e \in E^\Gamma$, we let $v(e)$ be the set $e^{[1]} \equiv e \cap V^\Gamma$. If $|v(e)| = 2$, the edge e is an internal edge. If $|v(e)| = 1$, e is an external edge, as self-loops are excluded.

Definition 2.1. A pair (v, e) with $e \in n(v)$ is called a half-edge. We let H^Γ be the set of half-edges of Γ .

²The \mathbb{Q} -vectorspace of graphs with self-loops forms an ideal and co-ideal I_{tad} , and we can effectively work in a quotient H/I_{tad} .

An internal edge e defines two half-edges uniquely.³ An ordering of the set $v(e)$ defines an orientation of that edge. Reversing that ordering is called an edge swap.

An oriented internal edge e connects two vertices, which we call source, $s(e) \in V^\Gamma$, and target, $t(e) \in V^\Gamma$, for an edge oriented from source to target.

Half-edges will play an important role for us as our new graph polynomial, which we dub the corolla polynomial below, is actually based on the half-edges of a graph.

Definition 2.2. The set of all half-edges incident to a given vertex,

$$\text{cor}(v) := \bigcup_{e \in n(v)} (v, e),$$

is called the corolla at v .

Definition 2.3. We denote

$$P_e := \text{cor}(s(e))\text{cor}(t(e)),$$

as an ordered pair of corollas.

We call two such pairs P_e, P_f disjoint if the edges e and f are disjoint.

External edges e at v are identified with the half-edge (v, e) and are always regarded as oriented to the vertex.

Definition 2.4. We say that a graph is j -regular if all vertices have valence j , $|n(v)| = j$, $\forall v \in V^\Gamma$.

Let Γ be 3-regular. Let \mathcal{C}^Γ be the set of all its cycles (not circuits!). For $C \in \mathcal{C}^\Gamma$ and $v \in C$ a vertex, let $v_C \in H^\Gamma$ be the unique half-edge at v not in C_i .

Definition 2.5. A graph is n -connected if it is connected after removal of any n of its internal edges.

Remark 2.6. A 2-connected graph is commonly called one-particle irreducible (1PI) in physics.

2.2. Orientation and Cycles. We need oriented graphs for two reasons: to define graph homology, and to have cyclic ordering at each corolla so that each half-edge incident to a vertex has a precursor and a successor at that vertex.

Let M_k be an oriented Riemann surface of genus k . We call a graph k -compatible, if it can be drawn on M_k without self-intersections.

Definition 2.7. We say a graph is of genus k if it is k -compatible but not j -compatible for any $j < k$. A planar graph is of genus 0.

Definition 2.8. An orientation (Γ, or) of a graph Γ is an ordering of V^Γ together with an ordering of each $v(e)$ for all internal edges e . Here, an ordering of $v(e)$ is the choice of one of the two possible bijections between $v(e)$ and the set $\{s(e), t(e)\}$.

Two orderings distinguished by an even number of vertex permutations and edge swaps are equivalent. We write (Γ, or) for an oriented graph. We set $(\Gamma, or) = -(\Gamma, or')$ for or, or' inequivalent orientations of the same graph.

Lemma 2.9. (*Conant, Vogtmann [15]*) *An ordering of a 3-regular graph is equivalent to a cyclic ordering of all its corollas.*

Definition 2.10. For an oriented 3-regular graph of genus k and $e \in n(v)$, let e_+ and e_- be the edges before (e_-) and after (e_+) edge e in the cyclic ordering of the corolla at v , induced by the orientation of M_k .

Remark 2.11. Note that such an orientation of a graph is compatible with a strict ordering of edges: the ordering of vertices orders the pair $s(e), t(e)$ lexicographically (with $s(e) < t(e)$ say), while multiple edges having the same source and target are ordered by the orientation of the underlying Riemann surface.

³Note that we discard self-loops indeed. As a consequence, a chosen vertex and an edge incident to that vertex label a half-edge uniquely.

2.3. Graph homology. We will use various homologies on graphs, with corresponding boundary operators, and suitable variants to study the filtrations of graphs by the number of ghost cycles and the number of internal 4-valent vertices. We start with standard graph homology for scalar graphs with 3- and 4-valent vertices.

2.3.1. *Graph homology \tilde{s} (following Conant, Vogtmann [15]).* For an edge e in a graph Γ , let Γ_e be the graph where e shrinks to zero length. Its orientation is obtained as follows: we permute vertex labels collecting signs until the edge e connects vertex 1, $s(e) = 1$, to vertex 2, $t(e) = 2$. Let σ be the sign of the necessary permutations. Then we shrink edge e and the so-obtained vertex is labelled 1. We inherit all remaining edge orientations and the ordering of vertices remains unchanged, with vertices $3, 4, \dots, |V^\Gamma|$ relabelled to $2, 3, \dots, |V^\Gamma| - 1$. This defines an orientation of Γ_e . If σ is negative, we change the orientation by an edge swap.

For an oriented graph Γ , let

$$\tilde{s}\Gamma = \sum_{e \in E_I} \Gamma_e,$$

be a sum of graphs obtained by shrinking edge e and assigning the orientation as above. Graph homology comes from the classical result

Theorem 2.12. (*graph homology*) $\tilde{s} \circ \tilde{s} = 0$.

2.3.2. *Graphs with marked vertices.* We can restrict graph homology to graphs with vertex valence bounded by $|n(v)| \leq 4$ by setting all terms which have vertices of valence higher than four to zero in the image of s .

For such graphs, let V_3 be the set of 3-valent vertices, V_4 be the set of 4-valent vertices, so $V^\Gamma = V_3 \cup V_4$.

Let $W_4 \subset V_4$ be a chosen subset. If a vertex $w \in W_4$, call it a marked vertex. A pair (Γ, W_4) is a graph with marked vertices.

Let u_w be the obvious map which removes the marking at vertex w , and $W_4(\Gamma)$ be the map which marks the vertices of Γ which are in W_4 . So $u_w(\Gamma, W_4) = (\Gamma, W_4 - w)$. Note that an orientation of Γ induces an ordering of the set W_4 . Set $\sigma(w) = j$ if and only if $w \in W_4$ is in the j -th place in that ordering.

We extend graph homology with boundary \tilde{s} to graphs with marked 4-valent vertices and boundary \tilde{S} by setting

$$(2) \quad \tilde{S}(\Gamma, W_4) = (s(\Gamma), W_4) + (-1)^{|V_4|} \sum_w (-1)^{\sigma(w)} (\Gamma, W_4 - w).$$

Proposition 2.13. \tilde{S} is a differential:

$$\tilde{S}^2 = 0.$$

Proof. We compute:

$$\begin{aligned} \tilde{S}^2(\Gamma, W_4) &= \tilde{S} \left((\tilde{s}(\Gamma), W_4) + (-1)^{|V_4|} \sum_w (-1)^{\sigma(w)} (\Gamma, W_4 - w) \right) \\ &= (-1)^{|V_4|+1} \sum_w (-1)^{\sigma(w)} (\tilde{s}(\Gamma), W_4 - w) + (-1)^{|V_4|} \sum_w (-1)^{\sigma(w)} (\tilde{s}(\Gamma), W_4 - w) \\ &\quad + (-1)^{|V_4|} \sum_{w \in W_4} (-1)^{\sigma(w)} \sum_{u \in W_4 - w} (-1)^{\sigma(u)} (\tilde{s}(\Gamma), W_4 - w - u) \\ &= 0, \end{aligned}$$

where the last line vanishes due to the ordering of vertices in W_4 , which makes sure that each pair (w, u) appears twice with a relative sign. \square

We will soon see that the sum over all connected graphs is in the kernel of \tilde{S} .

Remark 2.14. While the above operators \tilde{s}, \tilde{S} were defined on scalar graphs, we also have variants for gauge theory graphs where we have internal gauge boson or ghost propagators or fermion propagators. There are obvious restrictions then to shrink only edges which connect two 3-gluon vertices. The detailed homology operations available in such circumstances are exhibited in Section 4.

We now make graphs into a Hopf algebra.

2.4. Algebra of graphs. As we said before, we consider graphs as generators of a free commutative \mathbb{Q} -algebra of graphs H . We write \mathbb{I} for the unit represented by the empty set, with disjoint union of graphs furnishing the product.

Definition 2.15. The number of external edges $n_E := |E_E^\Gamma|$ assigns the weight

$$\omega^\Gamma := 4 - n_E = |V_3^\Gamma| - 2|E_I^\Gamma| + 4|\Gamma|,$$

to a graph.

A graph has positive valuation if $\omega^\Gamma \geq 0$. A graph has ι -valuation if $\omega^\Gamma \geq \iota$. Note that the valuation of a graph is invariant under shrinking edges.

The most obvious Hopf algebra structure is given by a co-product based on subgraphs of non-negative weights:

$$\Delta\Gamma = \Gamma \otimes \mathbb{I} + \mathbb{I} \otimes \Gamma + \sum_{\emptyset \neq \gamma = \coprod \gamma_i, \omega^{\gamma_i} \geq 0} \gamma \otimes \Gamma/\gamma,$$

is a coproduct, for a connected commutative Hopf algebra with unit \mathbb{I} and the span of all non-trivial graphs as augmentation ideal, as usual [16].

Definition 2.16. Let $f := \{\gamma_i\}$ be a subset of proper positive valued 1PI subgraphs $\gamma_i \subset \Gamma$ such that any two elements γ_i, γ_j of f fulfill: $\gamma_i \cap \gamma_j = \emptyset$, or $\gamma_i \subset \gamma_j$ or $\gamma_j \subset \gamma_i$. Then, f is called a forest.

Definition 2.17. It is maximal, if and only if Γ/f contains no positive valued proper subgraph. It is complete, if it contains all positive valued proper subgraphs of all its elements.

We denote by $|f|$ is the number of elements of f and by \mathcal{F}^Γ the set of all forests of Γ . For a union of graphs $\gamma = \bigcup_i \gamma_i$ we say it has ι valuation if all its components have. There are Hopf algebras for any ι -valuation:

$$\Delta_\iota(\Gamma) = \Gamma \otimes \mathbb{I} + \mathbb{I} \otimes \Gamma + \sum_{\gamma, \omega(\gamma) \geq \iota} \gamma \otimes \Gamma/\gamma.$$

In particular the antipode for positive valued graphs can be written as

$$S(\Gamma) = - \sum_{f \in \mathcal{F}^\Gamma} (-1)^{|f|} f \times \Gamma/f,$$

where the sum includes the empty set.

If the number of external edges of a subgraph γ is greater than two, $|\gamma_E^{[1]}| > 2$, then γ shrinks to a vertex in Γ/γ . If it equals two, the two external edges are identified to a single edge in Γ/γ .⁴

These Hopf algebras on scalar graphs straightforwardly generalize to gauge theory graphs and in particular act on the sum of all graphs contributing to a given amplitude—the combinatorial Green function—filtered by the number of 4-valent vertices and the number of ghost loops.

From now on, we demand that orientations of a graph Γ are such that for any proper subgraph $\gamma \subset \Gamma$ with two external edges e, f , $|\gamma_E^{[1]}| = 2$, the edges e, f form a consistently oriented edge in Γ/γ . Then, the orientation of Γ determines the orientations of f and Γ/f , for all forests f .

Definition 2.18. We call e_+^γ the edge pointing towards γ , e_-^γ the edge pointing away from it.

Lemma 2.19. Let $\gamma_1, \gamma_2 \subset \Gamma$ be two proper propagator subgraphs, $|\gamma_{i,E}^{[1]}| = 2$. Then, either: $\gamma_1 \subset \gamma_2$ or $\gamma_2 \subset \gamma_1$ or $\gamma_1 \cap \gamma_2 = \emptyset$.

Proof. This follows from [17, Lemma 51]. □

⁴If we were to have massive particles, we had to blow up notation slightly.

3. FEYNMAN RULES FOR SCALAR INTEGRANDS

First, some general remarks. For n external edges and l loops, an overall factor $c = (-i)^{n+3(l-1)}g^{n+2(l-1)}$ is not explicitly given below. All momentum integrals are understood in Euclidean space, all parametric integrals over the real simplex $\sigma : \{A_e > 0\}$, with boundary $\prod_{e \in \Gamma^{[1]}} A_e = 0$. A pairing between a parametric integrand and a simplex as in

$$\Phi(\Gamma) = \int_{\sigma} d\underline{A}_{\Gamma^{[1]}} I_{\Gamma},$$

means just that: the pair of the simplex and a form, and is to be regarded as an honest integral only when the integrand is replaced by its suitably renormalized form as defined below, typically indicated by a superscript R , so that the integral actually exists. We often simply write \int for \int_{σ} .

Let Γ be a 3-regular graph as defined in Definition 2.4. In order to define the momenta, choose an orientation on Γ , which we represent by ε (the incidence matrix):

$$\varepsilon_{ve} = \begin{cases} 1 & \text{if the vertex } v \text{ is the endpoint of the edge } e, \\ -1 & \text{if the vertex } v \text{ is the starting point of the edge } e, \\ 0 & \text{if } e \text{ is not incident on } v. \end{cases}$$

Furthermore, choose a basis of loops $L \subset \mathcal{C}^{\Gamma}$ of $l =: |\Gamma|$ independent loops of Γ and for each $\ell \in L$ an orientation given by ε^{ℓ} .⁵

The Feynman amplitude of Γ is⁶

$$\Phi(\Gamma) = \mathbf{Q}_{\Gamma}^E \int \frac{d\underline{k}_L}{(2\pi)^{dl}} \left(\prod_{e \in \Gamma^{[1]}} \frac{1}{\xi_e^{r_2}} \right),$$

where

$$\int \frac{d\underline{k}_L}{(2\pi)^{dl}} := \prod_{\ell \in L} \int_{\mathbb{R}^4} \frac{d^d k_{\ell}}{(2\pi)^d},$$

and

$$\varepsilon_{ve} \xi'_e := \varepsilon_{ve} \xi_e + \sum_{\substack{\ell \in L \\ \ell^{[1]} \ni e}} \varepsilon_{ve}^{\ell} k_{\ell}.$$

Note that we also include the external momenta, which is useful for our purposes. This gives just an overall factor above,

$$\mathbf{Q}_{\Gamma}^E = \prod_{e \in \Gamma_{\text{ext}}^{[1]}} \xi_e^2.$$

We define

$$- \sum_{e \in \Gamma^{[1]}, e \text{ adj. to } v} \varepsilon_{ve} \xi_e =: \xi_v,$$

to be the external momentum for every vertex v of our graph. Note that if v is a vertex to which an external edge e is adjacent, ξ_v does not equal ξ_e .

In Schwinger parametric form, the Feynman amplitude for generic ξ_e is

$$\Phi(\Gamma) = \int d\underline{A}_{\Gamma^{[1]}} I_{\Gamma},$$

where the integration is a short-hand notation for

$$\int d\underline{A}_{\Gamma^{[1]}} = \prod_{e \in \Gamma^{[1]}} \int_0^{\infty} dA_e = \int_{\sigma} dA_e,$$

⁵ ε^{ℓ} is such that $\varepsilon_{ve_1}^{\ell} = -\varepsilon_{ve_2}^{\ell}$, where e_1 and e_2 are the two edges adjacent to v , which are inside ℓ .

⁶ d is the dimension of spacetime, and can be safely set to four as later on we will renormalize the parametric integrand before integrating. If the reader wishes to integrate first, d can be used as a regulator, which is mathematically questionable though [18] when combined with minimal subtraction despite its popularity in physics.

and the integrand is obtained from integrating the universal quadric ([1])

$$Q_\Gamma := \sum_{e \in \Gamma^{[1]}} A_e \xi_e'^2,$$

so that analytically, we study the expression

$$I_\Gamma := \mathbf{Q}_\Gamma^E \int \frac{dk_L}{(2\pi)^{dL}} \prod_e e^{-Q_\Gamma} \prod_v \delta^4 \left(\sum \varepsilon_{ev} k(e) \right),$$

which corresponds to a graph where at each vertex v , an external momentum ξ_v is entering, and which can be written in the form of Eq.(1), by use of the graph polynomials, to which we now turn.

3.1. The first Kirchhoff polynomial ψ_Γ . For the first Kirchhoff polynomial consider the short exact sequence

$$(3) \quad 0 \rightarrow H^1 \rightarrow \mathbb{Q}^E \xrightarrow{\partial} \mathbb{Q}^{V,0} \rightarrow 0.$$

Here, H^1 is provided by a chosen basis for the algebraically independent loops of a graph Γ . $E = |E^\Gamma|$ is the number of edges and $V = |V^\Gamma|$ the number of vertices, so \mathbb{Q}^E is an E -dimensional \mathbb{Q} -vectorspace generated by the edges, similar $\mathbb{Q}^{V,0}$ for the vertices with a side constraint setting the sum of all vertices to zero.

Consider the matrix (see [1, 19])

$$N_0 \equiv (N_0)_{ij} = \sum_{e \in l_i \cap l_j} A_e,$$

for $l_i, l_j \in H^1$.

Define the *first Kirchhoff polynomial* as the determinant

$$\psi_\Gamma := |N_0|.$$

Proposition 3.1. ([1, Prop.2.2]) *The first Kirchhoff polynomial can be written as*

$$\psi_\Gamma = \sum_T \prod_{e \notin T} A_e$$

where the sum on the right is over spanning trees T of Γ .

3.2. The second Kirchhoff polynomial ϕ_Γ and $|N|_{\text{Pf}}$. Let σ^i , $i \in 1, 2, 3$ be the three Pauli matrices, and $\sigma^0 = \mathbb{I}_{2 \times 2}$ the unit matrix.

For the second Kirchhoff polynomial, augment the matrix N_0 to a new matrix N in the following way:

- i. Assign to each edge e a quaternion

$$\mathbf{q}_e := q_0 \sigma^0 - i \sum_{j=1}^3 q_j \sigma^j,$$

so that $\xi_e^2 \mathbb{I}_{2 \times 2} = \mathbf{q}_e \bar{\mathbf{q}}_e$, and to the loop l_i , the quaternion

$$u_i = \sum_{e \in l_i} A_e q_e.$$

- ii. Consider the column vector $u = (u_i)$ and the conjugated transposed row vector \bar{u} . Augment u as the rightmost column vector to M , and \bar{u} as the bottom row vector.
- iii. Add a new diagonal entry at the bottom right $\sum_e q_e \bar{q}_e A_e$.

Note that by momentum conservation, to each vertex, we assign a momentum ξ_v . and a corresponding quaternion \mathbf{q}_v .

The matrix N has a well-defined Pfaffian determinant (see [19]) with a remarkable form obtained for generic ξ_e and hence generic ξ_v :

Lemma 3.2. ([19, Eq.3.12])

$$|N|_{\text{Pf}} = - \sum_{T_1 \cup T_2} \left(\sum_{e \notin T_1 \cup T_2} \tau(e) \xi_e \right)^2 \prod_{e \notin T_1 \cup T_2} A_e,$$

where $\tau(e)$ is $+1$ if e is oriented from T_1 to T_2 and -1 else.

Proof. See [19]. □

Note that $|N|_{\text{Pf}} = |N|_{\text{Pf}}(\{\xi_v\})$ is a function of all ξ_v , $v \in \Gamma^{[0]}$. It gives the second Symanzik polynomial upon setting the ξ_e in accordance with the external momenta:

$$Q : \xi_e \rightarrow \xi_e + q_e,$$

and setting $\xi_e = 0$ for all edges e afterwards.

Proposition 3.3.

$$\phi_{\Gamma} \mathbb{I}_{2 \times 2} = Q(|N|_{\text{Pf}})|_{\xi_e=0} = - \sum_{T_1 \cup T_2} Q(T_1) \bar{Q}(T_2) \prod_{e \notin T_1 \cup T_2} A_e,$$

where the sum is over spanning two-forests $T_1 \cup T_2$, so T_1, T_2 are two disjoint non-empty trees which together cover all vertices of Γ . Here, $Q(T_i) = \sum_{v \in T_i^0} \mathbf{q}_v$, v a vertex of T_i , the momentum q_v incoming at that vertex expressed in the quaternionic basis.

Remark 3.4. For a Feynman graph Γ contributing to a scattering amplitude r with k external edges adjacent to $m \leq k$ vertices of Γ , ϕ_{Γ} is the quantity of interest. The quantity $|N|_{\text{Pf}}$, which assigns a momentum ξ_v to every vertex (not only to those m which have external edges attached) is more natural from a graph-theoretic viewpoint. For us, it has the added advantage that it assigns a four-momentum ξ_e to every vertex. Derivatives with respect to such four-momenta will generate gauge theory Feynman rules for us below.


Corollary 3.5. $\frac{\partial}{\partial \xi_{e\mu}} |N|_{\text{Pf}}$ is linear in $(A_e \xi_e^\mu)$.

For edges $e \neq f$, $\frac{\partial^2}{\partial \xi_{e\mu} \partial \xi_{(f)\nu}} |N|_{\text{Pf}}$ is linear in $(A_e A_f g^{\mu\nu})$ and constant in all ξ_e .

For $e = f$, $\frac{\partial^2}{\partial \xi_{e\mu} \partial \xi_{e\nu}} |N|_{\text{Pf}}$ is linear in $g^{\mu\nu}$, constant in all ξ_e and linear in A_e .

Proof. This follows readily from Lemma 3.2. □

Example 3.6. We let



with $\{1, 2, 3\}, \{1, 2, 4\}$ a basis for the cycles of Γ . Then,

$$N := \begin{pmatrix} N_0 := \left(\sum_{e \in h_i \cap h_j} A_e \right)_{ij} & \sum_{e \in h_j} \mu_e A_e \\ \sum_{e \in h_j} \bar{\mu}_e A_e & \sum_{e \in \Gamma^{[1]}} \bar{\mu}_e \mu_e A_e \end{pmatrix}$$

so

$$N_{\Gamma} = \begin{pmatrix} A_1 + A_2 + A_3 & A_1 + A_2 & A_1 \mu_1 + A_2 \mu_2 + A_3 \mu_3 \\ A_1 + A_2 & A_1 + A_2 + A_4 & A_1 \mu_1 + A_2 \mu_2 + A_4 \mu_4 \\ A_1 \bar{\mu}_1 + A_2 \bar{\mu}_2 + A_3 \bar{\mu}_3 & A_1 \bar{\mu}_1 + A_2 \bar{\mu}_2 + A_4 \bar{\mu}_4 & \sum_{i=1}^4 A_i \bar{\mu}_i \mu_i \end{pmatrix}$$

$$\psi_{\Gamma} = (A_1 + A_2)(A_3 + A_4) + A_3 A_4 = \sum_{\text{sp. Tr. } T} \prod_{e \notin T} A_e$$

and

$$\begin{aligned} \phi_{\Gamma} &= -(A_3 + A_4) A_1 A_2 p_a^2 + A_2 A_3 A_4 p_b^2 + A_1 A_3 A_4 p_c^2 \\ &= \sum_{\text{sp. 2-Tr. } T_1 \cup T_2} Q(T_1) \cdot Q(T_2) \prod_{e \notin T_1 \cup T_2} A_e. \end{aligned}$$

Remark 3.7. We use the two Symanzik polynomials to integrate out loop momenta in scalar integrands upon using

$$\frac{1}{\xi'_e \cdot \xi'_e} = \int_0^\infty e^{-A \xi'_e \cdot \xi'_e} dA, \quad \xi'_e \cdot \xi'_e = \xi_e'^2,$$

so that we get back Eq.(1).

3.3. Correction for quadratic subdivergences. The renormalized massless quadratically divergent two-point self-energy

$$\Sigma_R(q^2, \mu^2) = q^2 \sigma(q^2/\mu^2),$$

vanishes at $q^2 = 0$, and $\sigma = \Sigma_R/q^2$ vanishes at $q^2 = \mu^2$:

$$(4) \quad \Sigma_R(q^2, \mu^2)|_{q^2=0} = 0, \quad \left(\frac{1}{q^2} \Sigma_R(q^2, \mu^2) \right)|_{q^2=\mu^2} = 0.$$

This fixes the two renormalization conditions for any graph contributing to a massless quadratically divergent two-point functions which we employ. Transversality of the gluon propagator self-energy

$$\Pi_{\mu\nu} = q^2 (g_{\mu\nu} - \hat{q}_\mu \hat{q}_\nu) \Pi(q^2/\mu^2),$$

renders this as a single condition $\Pi(1) = 0$ on the sum of all contributing graphs at each loop order.

All graphs contributing to other Green functions are renormalized by simple subtractions at chosen kinematics (see [17] for a complete discussion).

Let us now discuss the correction factor for each quadratically divergent (sub)graph. The scalar integrand in parametric variables is

$$I_\Gamma = \frac{e^{-\frac{|N_\Gamma| \text{Pf}}{\psi_\Gamma}}}{\psi_\Gamma^2} dA_1 \cdots dA_{|\Gamma^{[1]}|}.$$

In fact, for gauge theory we need to study integrands which are slightly generalized:

$$\tilde{I}_{\Gamma, F} = F \frac{e^{-\frac{|N_\Gamma| \text{Pf}}{\psi_\Gamma}}}{\psi_\Gamma^2} dA_1 \cdots dA_{|\Gamma^{[1]}|}.$$

Here,

$$F = \frac{F_N(\{A_e\})}{F_D(\{A_e\})}$$

is a rational function of the variables A_e , which is a quotient of homogeneous polynomials F_N, F_D . In fact, our integrand in Yang–Mills or gauge theory will give us a finite sum of such terms with different F .

We want to analyse the degree of the form

$$A_\Gamma^F = \frac{F}{\psi_\Gamma^2} dA_1 \cdots dA_{|\Gamma^{[1]}|}.$$

Consider a set of variables $\gamma_I^{[1]} \subseteq \Gamma_I^{[1]}$, for $\gamma \subset \Gamma$ a subgraph (the case $\gamma = \Gamma$ is allowed). For a polynomial function $f = f(\{A_e\})$ we let $|f|_{\gamma_I^{[1]}}$ be twice⁷ the degree of that polynomial in the variables indicated. For a rational function which is a quotient f_1/f_2 of such functions, we have $|f_1/f_2|_{\gamma_I^{[1]}} = |f_1|_{\gamma_I^{[1]}} - |f_2|_{\gamma_I^{[1]}}$. Also, $|dA_1 \cdots dA_{|\Gamma^{[1]}|}|_{\gamma_I^{[1]}} = |\gamma_I^{[1]}|$.

Proposition 3.8. *For any A_Γ^F appearing in the integrand of a graph Γ , we have $|A_\Gamma^F|_{\gamma_I^{[1]}} \leq \omega^\gamma$.*

Proof. All short-distance singularities of a Feynman integrand correspond to forests and their degree is bounded by the powercounting ω^γ , for all γ appearing as components of forests. \square

Now assume $|A_\Gamma^F|_{\gamma_I^{[1]}} = 2$ for some γ . Such quadratically divergent short-distance singularities can only originate from self-energy subgraphs. Such graphs have two distinguished vertices at which external edges are adjacent. Moreover, it follows from Lemma 2.19 that quadratically divergent subgraphs are either disjoint or nested. We have immediately from the definition of the second Kirchhoff polynomial $\phi_\gamma =: q^2 \psi_{\gamma_\bullet}$, which defines γ_\bullet to be the graph where those two vertices are identified [17].

⁷The mass dimension of an edge variable A_e is -2 .

For each $|A_\Gamma^F|_{\gamma_I^{[1]}}$ above, let 2_Γ^F be the set of subgraphs $\gamma \subseteq \Gamma$ such that $|A_\Gamma^F|_{\gamma_I^{[1]}} = 2$, for all $\gamma \in 2_\Gamma^F$. Define, for all A_Γ^F ,

$$(5) \quad \overline{A}_\Gamma^F := A_\Gamma^F \prod_{\gamma \in 2_\Gamma^F} \frac{\psi_{\gamma \bullet}}{\psi_\gamma A_{e_+^\gamma}}.$$

Lemma 3.9. *The form \overline{A}_Γ^F has only logarithmic poles along divergent subgraphs including self-energy subgraphs. Renormalizing these remaining logarithmic poles of self-energy subgraphs at a fixed μ^2 imposes renormalization conditions Eq.(4).⁸*

Proof. A partial integration with respect to the quadratic subgraph variables renders its overall divergence logarithmic. The boundary term is eliminated by our renormalization conditions, which adopt the BPHZ conditions of massive propagators to the renormalization conditions adopted here for massless gluons. See [17, Section 3.5]. \square

Let us exhibit this in an example. Integrating the universal quadric, Eq.(3), only the propagators for the quadratically divergent subgraph γ leaves us with a contribution

$$\frac{1}{\xi_{e_-^\gamma}'} F \frac{e^{-\frac{\xi_e'^2 \psi_{\gamma \bullet}}{\psi_\gamma}} (dA)_\gamma}{\psi_\gamma^2} \frac{1}{\xi_{e_+^\gamma}'},$$

with e_\pm^γ defined in Def.(2.18).

Setting $A_i = t_\gamma a_i$ isolates the quadratic divergence:

$$\frac{1}{\xi_{e_-^\gamma}'} F e^{-t_\gamma \frac{\xi_e'^2 \psi_{\gamma \bullet}}{\psi_\gamma}} \Omega_\gamma \wedge t_\gamma \frac{1}{\xi_{e_+^\gamma}'}$$

Consider the integral $\lim_{c_\gamma \rightarrow 0} \int_{c_\gamma}^\infty$ against the above, and partially integrate with respect to t_γ . This gives, modulo terms which vanish when $c_\gamma \rightarrow 0$, a boundary term

$$\frac{1}{\xi_{e_-^\gamma}'} \frac{F}{\psi_\gamma^2} \left(1 - \xi_e'^2 \frac{\psi_{\gamma \bullet}}{\psi_\gamma} \right) \frac{1}{\xi_{e_+^\gamma}'}$$

which is polynomial in ξ_e' and hence vanishes in our renormalization conditions.

What remains though is the logarithmically divergent

$$\frac{1}{\xi_{e_-^\gamma}'} \frac{F \psi_{\gamma \bullet} \xi_e'^2 e^{-t_\gamma \frac{\xi_e'^2 \psi_{\gamma \bullet}}{\psi_\gamma}} \Omega_\gamma \wedge t_\gamma}{t_\gamma \psi_\gamma^A} \frac{1}{\xi_{e_+^\gamma}'}$$

which justifies the result above, upon using momentum conservation $\xi_e' = \xi_{e_+^\gamma}'$, and

$$\frac{\xi_e'}{\xi_{e_+^\gamma}'} = 1 = \frac{1}{2\pi i} \oint_{\gamma_e} \frac{e^{-\xi_e'^2 A_e}}{A_e},$$

with γ_e a curve which picks the residue at $A_e = 0$. Collecting such residues will be automatic in our approach below.

Remark 3.10. For a self-energy graph γ and an F such that $|A_\gamma^F|_\gamma = 2$, and $\tilde{I}_\Gamma^F = \tilde{I}_\Gamma^F(q^2)$, the factor Eq.(5) is in accordance with subtractions $(\tilde{I}_\Gamma^F(q^2) - \tilde{I}_\Gamma^F(0))/q^2 - (\tilde{I}_\Gamma^F(\mu^2) - \tilde{I}_\Gamma^F(0))/\mu^2$, in accordance with the conditions (4).

⁸We consider the case of massless propagators. For the incorporation of masses, see [19].

3.4. Renormalization. By summing over all F , we have finally constructed an integrand which we will call \bar{U}_Γ with log-poles only along any forest. We can hence render it finite by the usual forest formula. We define

$$U_\Gamma^R := \sum_{f \in \mathcal{F}_\Gamma} (-1)^{|f|} Q(\bar{U}_{\Gamma/f}) Q_0(\bar{U}_f),$$

where for $f = \bigcup_i \gamma_i$, $U_f = \prod_i U_{\gamma_i}$ and Q, Q_0 are the maps

$$Q : \xi_e \rightarrow \xi_e + q_e, \quad Q_0 : \xi_e \rightarrow \xi_e + q_{e,0},$$

where q_e are the momenta as prescribed by the amplitude under consideration, $q_{e,0}$ those prescribed by the renormalization scheme for this amplitude.

Remark 3.11. Note that a single $\tilde{I}_{\Gamma,F}$ renormalizes similarly,

$$\tilde{I}_{\Gamma,F}^R = \bar{F} \sum_{f \in \mathcal{F}_\Gamma} (-1)^{|f|} Q \left(\frac{e^{-\frac{|N_{\Gamma/f}| Pf}}{\psi_{\Gamma/f}}}}{\psi_{\Gamma/f}^2} \right) Q_0 \left(\frac{e^{-\frac{|N_f| Pf}}{\psi_f}}}{\psi_f^2} \right),$$

with notation as in [17]. Note in particular how derivatives with respect to four-vectors ξ_e still act on $\tilde{I}_{\Gamma,F}^R$: for any monomial in derivatives $X_E := \prod_{e \in E} \partial_{\xi_{e\mu(e)}}$, we have

$$(6) \quad X_E \tilde{I}_{\Gamma,F}^R = \bar{F} \sum_{f \in \mathcal{F}_\Gamma} (-1)^{|f|} \left(\prod_{e \in E \cap \Gamma/f^{[1]}} \partial_{\xi_{e\mu(e)}} \right) Q \left(\frac{e^{-\frac{|N_{\Gamma/f}| Pf}}{\psi_{\Gamma/f}}}}{\psi_{\Gamma/f}^2} \right) \left(\prod_{e \in E \cap f^{[1]}} \partial_{\xi_{e\mu(e)}} \right) Q_0 \left(\frac{e^{-\frac{|N_f| Pf}}{\psi_f}}}{\psi_f^2} \right).$$

This is particularly useful for future work when combined with the projective renormalized integrand, see [17].

Also, even when Γ is a graph without external ghost lines, some of the forests appearing above can correspond to graphs with open ghost lines. Then, Eq.(6) reconfirms the formula Eq.(22) below for amplitudes with open ghost (or fermion, in an obvious modification) lines.

Remark 3.12. The regularized integrand in dimensional regularization is obtained by multiplying \bar{U}_Γ by $1/\psi_\Gamma^{(4-d)/2}$, and treating the Clifford algebra accordingly. Evaluating the parametric integrals on the regularized integrand first and renormalizing in accordance with our renormalization prescription produces the same renormalized results as above. Using a minimal subtraction scheme is different though. See [18] for a discussion of this point.

4. GAUGE THEORY GRAPHS

We now turn to graphs in gauge theory, as contrasted to 3-regular graphs in scalar field theory. While the latter were graphs which can be regarded as corollas with three half-edges, connected by gluing two half-edges from different corollas to an internal edge e which hence determine a pair of corollas P_e , the former are graphs with 3- and 4-valent vertices.

Again, we can consider them based on corollas, this time corollas which have either three or four half-edges of gauge boson type (indicated by wavy lines), or one gauge-boson half-edge with two half-edges of ghost type (indicated by consistently oriented straight dashed lines), or one gauge-boson half-edge with two half-edges of fermion type (indicated by consistently oriented straight full lines).

Also, we will mark in such graphs edges and vertices in various ways and let

$$\mathcal{G}_n^{\times, m, \circ}{}^{r, l}$$

be the set of all graphs with external half-edges specifying the amplitude r , with l loops and n 4-gluon vertices of type, m ghostloops. Similarly, we will indicate the number of marked edges and other qualifiers as needed.

If we want to leave a qualifier l, n, m, \dots unspecified (so that we consider the union of all sets with any number of such items), we replace it by $/$. A similar notation will be used below for series of graphs

$$X_n^{\times, m, \circ}{}^{r, l}$$

which are sums (for fixed l) or series (for $l = /$) of graphs weighted by symmetry, colour and other such factors as defined below. We dub such series combinatorial Green functions.

We now start adopting graph homology to our purposes in gauge theory.

4.1. Marking edges. Recall that the Feynman rule for the 4-valent vertex is

$$\begin{aligned} \Phi\left(\begin{array}{c} 1 \\ \diagdown \quad \diagup \\ \text{---} \quad \text{---} \\ \diagup \quad \diagdown \\ 2 \quad \quad 3 \quad \quad 4 \end{array}\right) &= + f^{a_1 a_2 b} f^{a_3 a_4 b} (g^{\mu_1 \mu_3} g^{\mu_2 \mu_4} - g^{\mu_1 \mu_4} g^{\mu_2 \mu_3}) \\ &+ f^{a_1 a_3 b} f^{a_2 a_4 b} (g^{\mu_1 \mu_2} g^{\mu_3 \mu_4} - g^{\mu_1 \mu_4} g^{\mu_3 \mu_2}) \\ &+ f^{a_1 a_4 b} f^{a_2 a_3 b} (g^{\mu_1 \mu_2} g^{\mu_4 \mu_3} - g^{\mu_1 \mu_3} g^{\mu_4 \mu_2}). \end{aligned}$$

We introduce a new edge type \rightsquigarrow which has the following Feynman rule:

$$(7) \quad \begin{aligned} \Phi\left(\begin{array}{c} 1 \\ \diagdown \quad \diagup \\ \rightsquigarrow \quad \rightsquigarrow \\ \diagup \quad \diagdown \\ 2 \quad \quad 3 \quad \quad 4 \end{array}\right) &= f^{a_1 a_2 b} f^{a_3 a_4 b} (g^{\mu_1 \mu_3} g^{\mu_2 \mu_4} - g^{\mu_1 \mu_4} g^{\mu_2 \mu_3}) \\ &=: \text{colour}\left(\begin{array}{c} \diagdown \quad \diagup \\ \rightsquigarrow \quad \rightsquigarrow \\ \diagup \quad \diagdown \end{array}\right) W_e, \end{aligned}$$

so that we can write the 4-point vertex as

$$(8) \quad \begin{array}{c} \diagdown \quad \diagup \\ \rightsquigarrow \quad \rightsquigarrow \\ \diagup \quad \diagdown \end{array} \sim \begin{array}{c} \diagdown \quad \diagup \\ \rightsquigarrow \quad \rightsquigarrow \\ \diagup \quad \diagdown \end{array} + \begin{array}{c} \diagdown \quad \diagup \\ \rightsquigarrow \quad \rightsquigarrow \\ \diagup \quad \diagdown \end{array} + \begin{array}{c} \diagdown \quad \diagup \\ \rightsquigarrow \quad \rightsquigarrow \\ \diagup \quad \diagdown \end{array}.$$

(The relation \sim denotes that the left- and right-hand side have the same Feynman amplitude.) Note that because of this relation, the internal marked edge does not correspond to a propagator. It is just a graphical way of writing the three terms of the 4-valent vertex.

For any graph Γ with marked edges, let $\bar{\Gamma}$ be the graph where the marked edges shrink to zero length.⁹ The Feynman–Schwinger integrand of a graph with marked edges is given for us by

$$(9) \quad I_\Gamma = \left(\prod_{e \in \Gamma_{\text{marked}}^{[1]}} W_e \right) \left(\prod_{v \in V^\Gamma, v \cap \Gamma_{\text{marked}}^{[1]} = \emptyset} V_v \right) I_{\bar{\Gamma}}.$$

Here, V_v is the colour-stripped part of the Feynman rule of a 3-gluon vertex,

$$(10) \quad V_v = \sum_{\text{cycl}(1,2,3)} (\xi_1 - \xi_2)_{\mu_3} g_{\mu_1 \mu_2}.$$

Note that the scalar integrand $I_{\bar{\Gamma}}$ does obviously not contain the edge variables of the \rightsquigarrow -marked edges.

For future use, we define

$$(11) \quad \mathcal{I}_\Gamma = \left(\prod_{e \in \Gamma_{\text{marked}}^{[1]}} W_e \right) \left(\prod_{v \in V^\Gamma, v \cap \Gamma_{\text{marked}}^{[1]} = \emptyset} V_v \right) e^{-\sum_{e \in \Gamma_{\text{unmarked}}} A_e \xi_e'^2},$$

which is such that

$$\int \frac{d^k_L}{(2\pi)^{dl}} \mathcal{I}_\Gamma = I_\Gamma.$$

Definition 4.1. Define a derivation $\chi_+ : H \rightarrow H$ on generators by

$$\chi_+ \Gamma = \sum_{e \in \Gamma_{\text{int}}^{[1]}} \chi_+^e \Gamma,$$

where

$$\chi_+^e \Gamma = \begin{cases} 0 & \text{if } e \text{ shares a vertex with a marked, fermion or ghost edge,} \\ \Gamma_{e \rightsquigarrow} & \text{otherwise.} \end{cases}$$


The next lemma shows how symmetry factors relate upon exchanging 4-valent vertices for a pair of corollas with a marked edge in-between. We consider graphs with l loops, k 4-gluon vertices and k' marked edges, for an amplitude r . Also, \mathcal{F} denotes unlabelled graphs, in contrast to labelled graphs in \mathcal{G} .

⁹If we have k marked edges, here are 3^k different graphs Γ which have the same $\bar{\Gamma}$.

Lemma 4.2. For any graph $\Gamma \in \mathcal{G}_{k, \chi, k' \sim \sim}^{r, l}$ we have

$$(12) \quad \frac{1}{\text{Sym}(\Gamma)} \Gamma \sim \frac{1}{k} \sum_{\substack{\Gamma' \in \mathcal{G}_{k-1, \chi, k'+1 \sim \sim}^{r, l} \\ \exists e \in \Gamma_{\text{marked}}^{[1]}: \Gamma'/e = \Gamma}} \frac{1}{\text{Sym}(\Gamma')} \Gamma'.$$

Proof. Let $v \in \Gamma_4^{[0]}$ with adjacent edges 1, 2, 3, 4:



(We do not show Γ 's external edges in the diagram.) Apply (8):

$$\frac{1}{\text{Sym}(\Gamma)} \Gamma = \frac{1}{\text{Sym}(\Gamma)} \left(\Gamma_1 + \Gamma_2 + \Gamma_3 \right).$$

The following three cases can occur:

1. The four edges adjacent to v are each un-interchangeable. In this case

$$\text{Sym}(\Gamma_1) = \text{Sym}(\Gamma_2) = \text{Sym}(\Gamma_3) = \text{Sym}(\Gamma_4),$$

so that

$$\frac{1}{\text{Sym}(\Gamma)} \Gamma \sim \frac{1}{\text{Sym}(\Gamma_1)} \Gamma_1 + \frac{1}{\text{Sym}(\Gamma_2)} \Gamma_2 + \frac{1}{\text{Sym}(\Gamma_3)} \Gamma_3 + \frac{1}{\text{Sym}(\Gamma_4)} \Gamma_4.$$

Note that the three graphs at the right-hand side are all non-isomorphic.

2. Two of v 's adjacent edges are interchangeable, say 1 and 2. Then

$$\Gamma_1 = \Gamma_2.$$

The symmetry factors of the new graphs are

$$\text{Sym}(\Gamma_1) = \text{Sym}(\Gamma_2),$$

and

$$\text{Sym}(\Gamma_3) = \frac{1}{2} \text{Sym}(\Gamma_1),$$

so that

$$\begin{aligned} \frac{1}{\text{Sym}(\Gamma)} \Gamma &\sim \frac{1}{\text{Sym}(\Gamma_1)} \left(\Gamma_1 + 2 \Gamma_3 \right) \\ &= \frac{1}{\text{Sym}(\Gamma_1)} \Gamma_1 + \frac{1}{\text{Sym}(\Gamma_3)} \Gamma_3. \end{aligned}$$

Note that the two graphs at the right-hand side are unequal.

3. Three of v 's adjacent edges are interchangeable, say 1, 2 and 3. Then

$$\Gamma_1 = \Gamma_2 = \Gamma_3,$$

and

$$\text{Sym}(\Gamma_1) = \frac{1}{3} \text{Sym}(\Gamma_4).$$

So:

$$\frac{1}{\text{Sym}(\Gamma)} \Gamma \sim \frac{3}{\text{Sym}(\Gamma_1)} \Gamma_1 = \frac{1}{\text{Sym}(\Gamma_4)} \Gamma_4.$$

Thus we can conclude that

$$\frac{1}{\text{Sym}(\Gamma)}\Gamma = \sum_{\substack{\Gamma' \in \overline{\mathcal{G}}^{n,l} \\ \Gamma' \sim_{k-1} \chi_{k',k'+1} \\ \exists e \in \Gamma'_{\text{marked}}^{[1]} : \Gamma'/e = \Gamma \\ \text{where the new vertex is } v}} \frac{1}{\text{Sym}(\Gamma')}\Gamma'.$$

The result follows up summing this over all 4-valent vertices in Γ , giving rise to the factor $\#\Gamma_4^{[0]} = k$:

$$\Gamma \sim \frac{1}{k} \sum_{\substack{\Gamma' \in \overline{\mathcal{G}}^{n,l} \\ \Gamma' \sim_{k-1} \chi_{k',k'+1} \\ \exists e \in \Gamma'_{\text{marked}}^{[1]} : \Gamma'/e = \Gamma}} \frac{1}{\text{Sym}(\Gamma')}\Phi(\Gamma').$$

□

Example 4.3. Take $\Gamma = \text{diagram}$ and apply equation (8) to the 4-valent vertex:

$$(13) \quad \frac{1}{2} \text{diagram} \sim \frac{1}{2} \left(\text{diagram} + \text{diagram} + \text{diagram} \right) = \text{diagram} + \frac{1}{2} \text{diagram}.$$

Example 4.4. Take $\Gamma = \text{diagram}$ and apply (8) to one of the two vertices:

$$(14) \quad \frac{1}{6} \text{diagram} \sim \frac{1}{6} \left(\text{diagram} + \text{diagram} + \text{diagram} \right) = \frac{1}{2} \text{diagram}.$$

Analogously, we get

$$\frac{1}{6} \text{diagram} \sim \frac{1}{2} \text{diagram},$$

so that we can write

$$(15) \quad \frac{1}{6} \text{diagram} \sim \frac{1}{2} \left(\frac{1}{2} \text{diagram} + \frac{1}{2} \text{diagram} \right).$$

Example 4.5.

$$\frac{1}{2} \text{diagram} \sim \frac{1}{2} \left(\text{diagram} + \text{diagram} + \text{diagram} \right) = \frac{1}{2} \text{diagram} + \text{diagram}.$$

The next lemma is crucial, as it shows that the fundamental relation between a 4-gluon vertex and a pair of 3-gluon vertices, in all three channels, gives a relation between combinatorial Green functions. We would have no chance at getting a well-defined gauge theory without such a relation.

Lemma 4.6. i. For any k and k' , $k' < k$:

$$\frac{1}{\binom{k}{k'}} X_{k-k'}^{n,l} \chi_{k',k'+1} \sim \frac{1}{\binom{k}{k'+1}} X_{k-k'-1}^{n,l} \chi_{k',k'+1}.$$

ii. For any k :

$$X_{k \chi}^{n,l} \sim X_{k \sim}^{n,l}.$$

Proof. For i. we have from lemma 4.2 for the Green's function $X_{k-k', \chi', k' \rightsquigarrow}^{n,l}$:

$$\begin{aligned}
X_{k-k', \chi', k' \rightsquigarrow}^{n,l} &= \sum_{\Gamma \in \overline{\mathcal{G}}_{k-k', \chi', k' \rightsquigarrow}^{n,l}} \frac{1}{\text{Sym}(\Gamma)} \Gamma \\
&\sim \frac{1}{k-k'} \sum_{\Gamma \in \overline{\mathcal{G}}_{k-k', \chi', k' \rightsquigarrow}^{n,l}} \sum_{\substack{\Gamma' \in \overline{\mathcal{G}}_{k-k'-1, \chi', k'+1 \rightsquigarrow}^{n,l} \\ \exists e \in \Gamma'^{[1]}_{\text{marked}}: \Gamma'/e = \Gamma}} \frac{1}{\text{Sym}(\Gamma')} \Gamma' \\
&= \frac{k'+1}{k-k'} \sum_{\Gamma' \in \overline{\mathcal{G}}_{k-k'-1, \chi', k'+1 \rightsquigarrow}^{n,l}} \frac{1}{\text{Sym}(\Gamma')} \Gamma' \\
&= \frac{k'+1}{k-k'} X_{k-k'-1, \chi', k'+1 \rightsquigarrow}^{n,l}.
\end{aligned}$$

The factor $k'+1$ appears because every graph in $\overline{\mathcal{G}}_{k-k', \chi', k' \rightsquigarrow}^{n,l}$ can be obtained from $k'+1$ graphs in $\overline{\mathcal{G}}_{k-k'-1, \chi', k'+1 \rightsquigarrow}^{n,l}$ by applying (8). Using the identity

$$\binom{k}{k'+1} = \frac{k-k'}{k'+1} \binom{k}{k'},$$

it follows that

$$\frac{1}{\binom{k}{k'}} X_{k-k', \chi', k' \rightsquigarrow}^{n,l} \sim \frac{1}{\binom{k}{k'+1}} X_{k-k'-1, \chi', k'+1 \rightsquigarrow}^{n,l}.$$

For ii. we have

$$X_{k, \chi}^{n,l} \sim \frac{1}{\binom{k}{k'}} X_{k-k', \chi', k' \rightsquigarrow}^{n,l}.$$

This is true by induction: it is an equality for $k'=0$ and the inductive step is true by i.. Taking $k'=k$ gives:

$$X_{k, \chi}^{n,l} \sim X_{k \rightsquigarrow}^{n,l}. \quad \square$$

In the following, if r is an n -gluon amplitude, we often replace the subscript r by n , as in this example:

Example 4.7. Take $n=3$, $l=1$ and $k=1$.

$$\begin{aligned}
X_{1, \chi}^{3,1} &= \frac{1}{2} \text{diagram}_1 + \frac{1}{2} \text{diagram}_2 + \frac{1}{2} \text{diagram}_3 \\
&\sim \text{diagram}_4 + \text{diagram}_5 + \text{diagram}_6 + \frac{1}{2} \text{diagram}_7 + \frac{1}{2} \text{diagram}_8 + \frac{1}{2} \text{diagram}_9 = X_{1 \rightsquigarrow}^{3,1}
\end{aligned}$$

where we have used Eq.(13).

Example 4.8. Take $n=2$, $l=2$ and $k=2$. Note that $\overline{\mathcal{G}}_{2, \chi}^{2,2}$ contains just one graph and use (14) and (15):

$$X_{2, \chi}^{2,2} = \frac{1}{6} \text{diagram}_1 \sim \frac{1}{2} \text{diagram}_2 \sim \frac{1}{2} \text{diagram}_3 + \text{diagram}_4 = X_{2 \rightsquigarrow}^{2,2}.$$

Lemma 4.9. i. For any graph $\Gamma \in \overline{\mathcal{G}}^{n,l}$:

$$\frac{1}{\text{Sym}(\Gamma)} \frac{(\chi_+)^k}{k!} \Gamma = \sum_{\substack{\Gamma' \in \overline{\mathcal{G}}^{n,l} \\ \text{skeleton}(\Gamma') = \Gamma}} \frac{1}{\text{Sym}(\Gamma')} \Gamma'.$$

where the skeleton of a marked graph is the graph with its markings removed.

- ii. For any $k \geq 0$, $\frac{(\chi_+)^k}{k!} X_{k \rightsquigarrow}^{n,l} = X_{k \rightsquigarrow}^{n,l}$.
- iii. We have $e^{\chi_+} X_{\rightsquigarrow}^{n,l} = X_{\rightsquigarrow}^{n,l} = X_{\chi}^{n,l}$.


Proof. For i. we have

$$\begin{aligned}
\frac{1}{\text{Sym}(\Gamma)} \frac{(\chi_+)^k}{k!} [\Gamma] &= \frac{1}{\text{Sym}(\Gamma)} \sum_{\{e_1, \dots, e_k\} \subset \Gamma_{\text{int}}^{[1]}} [\chi_+^{e_1} \dots \chi_+^{e_k} \Gamma] \\
&= \frac{1}{\text{Sym}(\Gamma)} \sum_{\substack{\Gamma' \in \overline{\mathcal{G}}_{k \rightsquigarrow}^{n,l} \\ \text{skeleton}(\Gamma') = [\Gamma]}} \#\{\{e_1, \dots, e_k\} \subset \Gamma_{\text{int}}^{[1]} \mid [\chi_+^{e_1} \dots \chi_+^{e_k} \Gamma] = \Gamma'\} \Gamma' \\
&= \sum_{\substack{\Gamma' \in \overline{\mathcal{G}}_{k \rightsquigarrow}^{n,l} \\ \text{skeleton}(\Gamma') = [\Gamma]}} \frac{1}{\text{Sym}(\Gamma')} \Gamma'.
\end{aligned}$$

For ii. we apply i. to the combinatorial Green's function $X^{n,l}$, instead of a single graph. Summing over all graphs in $\overline{\mathcal{G}}^{n,l}$ yields

$$\begin{aligned}
\frac{(\chi_+)^k}{k!} X^{n,l} &= \sum_{\Gamma \in \overline{\mathcal{G}}^{n,l}} \frac{1}{\text{Sym}(\Gamma)} \frac{(\chi_+)^k}{k!} \Gamma \\
&= \sum_{\Gamma' \in \overline{\mathcal{G}}_{k \rightsquigarrow}^{n,l}} \frac{1}{\text{Sym}(\Gamma')} \Gamma' = X_{k \rightsquigarrow}^{n,l}.
\end{aligned}$$

Finally, iii. follows directly from ii. by taking the sum over k . □

Example 4.10. Take  and $k = 2$; then lemma 4.9.i reads

$$\frac{1}{2} \frac{(\chi_+)^2}{2} \text{loop} = \frac{1}{2} (\text{loop} + \text{loop}) = \text{loop}.$$

Remark 4.11. Note that χ_+ has a non-trivial kernel as it can create self-loops graphs, for example:

$$\frac{1}{2} \chi_+ \text{loop} = \text{loop} \sim \frac{1}{2} \text{loop}$$

Here we have used that

$$\text{loop} \sim \text{loop} + \text{loop} + \text{loop} \sim 2 \text{loop}.$$

This does not influence our results, since self-loops have amplitude zero.

It is now time to study graph homology, again by studying marked edges, but now the labelling plays a crucial role.

4.2. The graph differential s for gauge theory graphs.

Definition 4.12. The derivation $s : H \rightarrow H$ is given on generators by

$$s\Gamma = \sum_{e \in \Gamma_{\text{int}}^{[1]}} (-) \#\{e' \in \Gamma_{\text{marked}}^{[1]} \mid e' < e\} s_e \Gamma$$

where

$$s_e \Gamma = \begin{cases} 0 & \text{if } e \text{ shares a vertex with a marked or ghost edge} \\ \Gamma_{e \rightsquigarrow} & \text{otherwise} \end{cases}$$

In the above, $<$ is a (strict) total ordering on $\Gamma^{[1]}$.

Next, we want to distinguish the markings created by χ_+ and s . Therefore we draw the latter with two lines instead of one. So two lines indicate the action of s , and we denote: $\Gamma_{\rightsquigarrow}^{[1]} \cup \Gamma_{\rightsquigarrow}^{[1]} = \Gamma_{\text{marked}}^{[1]} \subset \Gamma_{\text{int}}^{[1]}$.


Proposition 4.13. s is a differential: $s^2\Gamma = 0$.

Proof. We compute

$$\begin{aligned}
s^2\Gamma &= \sum_{e_1, e_2 \in \Gamma_{\text{int}}^{[1]}} (-) \#\{e'_1 \in \Gamma_{\text{marked}}^{[1]} \mid e'_1 < e_1\} + \#\{e'_2 \in \Gamma_{\text{marked}}^{[1]} \cup \{e_1\} \mid e'_2 < e_2\} s_{e_1} s_{e_2} \Gamma \\
&= \sum_{\substack{e_1, e_2 \in \Gamma_{\text{int}}^{[1]} \\ e_1 < e_2}} (-) \#\{e \in \Gamma_{\text{marked}}^{[1]} \mid e_1 < e < e_2\} + 1 s_{e_1} s_{e_2} \Gamma \\
&\quad + \sum_{\substack{e_1, e_2 \in \Gamma_{\text{int}}^{[1]} \\ e_1 > e_2}} (-) \#\{e \in \Gamma_{\text{marked}}^{[1]} \mid e_2 < e < e_1\} s_{e_1} s_{e_2} \Gamma \\
&= 0.
\end{aligned}$$

□

Example 4.14. We work with labelled graphs, eg.



for which

$$\begin{aligned}
s^2 \text{ (graph)} &= s \text{ (graph)} + s \text{ (graph)} + s \text{ (graph)} + s \text{ (graph)} + s \text{ (graph)} \\
&= - \text{ (graph)} - \text{ (graph)} + \text{ (graph)} + \text{ (graph)} = 0.
\end{aligned}$$

4.3. The differential S for gauge theories. Marking edges, which corresponds upon summation of connected diagrams to shrinking pairs of two 3-gluon vertices to 4-gluon vertices, should match with the graphs with 4-gluon vertices present in the theory. This can be phrased homologically.

Definition 4.15. A derivation $S : H \rightarrow H$ is given by $S = s + \sigma$ where

$$\sigma\Gamma = (-) \#\Gamma_{\text{marked}}^{[1]} \sum_{e \in \Gamma^{[1]} \rightsquigarrow} (-) \#\{e' \in \Gamma^{[1]} \rightsquigarrow \mid e' > e\} \sigma_e \Gamma,$$

and

$$\sigma_e \Gamma = \Gamma_{e \rightsquigarrow \rightsquigarrow}.$$

Proposition 4.16. S is a differential: $S^2\Gamma = 0$.

Proof. A calculation shows that

$$\begin{aligned}
s\sigma\Gamma &= (-) \Gamma_{\text{marked}}^{[1]} \sum_{e_1 \in \Gamma_{\rightsquigarrow}^{[1]}} \sum_{e_2 \in \Gamma_{\text{int}}^{[1]}} (-) \#\{e'_1 \in \Gamma^{[1]} \rightsquigarrow \mid e'_1 > e_1\} + \#\{e'_2 \in \Gamma_{\text{marked}}^{[1]} \mid e'_2 < e_2\} s_{e_2} \sigma_{e_1} \Gamma \\
\sigma s\Gamma &= (-) \Gamma_{\text{marked}}^{[1]} + 1 \sum_{e_2 \in \Gamma_{\text{int}}^{[1]}} \sum_{e_1 \in \Gamma^{[1]} \rightsquigarrow} (-) \#\{e'_2 \in \Gamma_{\text{marked}}^{[1]} \mid e'_2 < e_2\} + \#\{e'_1 \in \Gamma^{[1]} \rightsquigarrow \mid e'_1 > e_1\} s_{e_2} \sigma_{e_1} \Gamma \\
&= -s\sigma\Gamma,
\end{aligned}$$

and also

$$\begin{aligned}
\sigma^2\Gamma &= \sum_{e_1 \in \Gamma^{[1]} \rightsquigarrow} \sum_{e_1 \in \Gamma^{[1]} \rightsquigarrow \setminus \{e_1\}} (-) \#\{e'_1 \in \Gamma^{[1]} \rightsquigarrow \mid e'_1 > e_1\} + \#\{e'_2 \in \Gamma^{[1]} \rightsquigarrow \setminus \{e_1\} \mid e'_2 > e_2\} \sigma_{e_1} \sigma_{e_2} \Gamma \\
&= \sum_{\substack{e_1, e_2 \in \Gamma^{[1]} \rightsquigarrow \\ e_1 < e_2}} (-) \#\{e \in \Gamma^{[1]} \rightsquigarrow \mid e_1 < e < e_2\} + 1 \sigma_{e_1} \sigma_{e_2} \Gamma \\
&\quad + \sum_{\substack{e_1, e_2 \in \Gamma^{[1]} \rightsquigarrow \\ e_1 > e_2}} (-) \#\{e \in \Gamma^{[1]} \rightsquigarrow \mid e_1 < e < e_2\} \sigma_{e_1} \sigma_{e_2} \Gamma \\
&= 0.
\end{aligned}$$

so that

$$S^2\Gamma = s^2\Gamma + s\sigma\Gamma + \sigma s\Gamma + \sigma^2\Gamma = 0.$$

□

Remark 4.17. Note that upon summing the markings in a 3-valent corolla, and identifying such a sum with a 4-valent vertex, the operators \tilde{s} and \tilde{S} here reduce to the operators s, S we had before in Eq.(2).

Example 4.18.

$$\begin{aligned} s\sigma \text{---} &= -s \text{---} = \text{---} \\ \sigma s \text{---} &= -\sigma \text{---} = -\text{---} \\ \sigma^2 \text{---} &= -\sigma \text{---} = 0 \\ S^2 \text{---} &= 0 \end{aligned}$$

Example 4.19.

$$\begin{aligned} s\sigma \text{---} &= -s \text{---} + s \text{---} = 0 \\ \sigma s \text{---} &= 0 \\ \sigma^2 \text{---} &= -\sigma \text{---} + \sigma \text{---} = -\text{---} + \text{---} = 0 \\ S^2 \text{---} &= 0 \end{aligned}$$

The cancellations between 3-gluon and 4-gluon vertices necessary to obtain a unitary and covariant gauge theory demand that shrinking internal edges in graphs with k_3 3-gluon vertices and k_4 4-gluon vertices matches with the graphs having $(k_3 - 2)$ 3-gluon vertices, and $(k_4 + 1)$ 4-gluon vertices. Rephrased in terms of our marked edges and using our sign conventions, that precisely is captured by

Proposition 4.20. *Let Γ be a graph without marked edges. Then:*

$$(16) \quad Se^{\chi+\Gamma} = 0.$$

Proof. By definition:

$$e^{\chi+\Gamma} = \sum_{k \geq 0} \sum_{\substack{e_1, \dots, e_k \in \Gamma_{\text{int}}^{[1]} \\ e_1 < \dots < e_k}} \chi_+^{e_1} \cdots \chi_+^{e_k} \Gamma$$

on which

$$\begin{aligned} se^{\chi+\Gamma} &= \sum_{k \geq 0} \sum_{\substack{e_1, \dots, e_{k+1} \in \Gamma_{\text{int}}^{[1]} \\ e_1 < \dots < e_{k+1}}} \sum_{l=1}^{k+1} (-)^{l-1} \chi_+^{e_1} \cdots s_{e_l} \cdots \chi_+^{e_{k+1}} \Gamma \\ &= \sum_{k \geq 1} \sum_{\substack{e_1, \dots, e_k \in \Gamma_{\text{int}}^{[1]} \\ e_1 < \dots < e_k}} \sum_{l=1}^k (-)^{l-1} \chi_+^{e_1} \cdots s_{e_l} \cdots \chi_+^{e_k} \Gamma \\ \sigma e^{\chi+\Gamma} &= \sum_{k \geq 1} \sum_{\substack{e_1, \dots, e_k \in \Gamma_{\text{int}}^{[1]} \\ e_1 < \dots < e_k}} (-)^k \sum_{l=1}^k (-)^{k-l} \chi_+^{e_1} \cdots s_{e_l} \cdots \chi_+^{e_k} \Gamma \\ &= -se^{\chi+\Gamma}. \end{aligned}$$

We conclude that

$$Se^{\chi+\Gamma} = (s + \sigma)e^{\chi+\Gamma} = 0.$$

□

Example 4.21.

$$\begin{aligned}
e^{X^+} \text{---} \text{---} &= \text{---} + \text{---} + \text{---} + \text{---} + \text{---} + \text{---} + \text{---} + \text{---} \\
se^{X^+} \text{---} \text{---} &= \text{---} + \text{---} + \text{---} + \text{---} + \text{---} \\
&\quad - \text{---} - \text{---} + \text{---} + \text{---} \\
\sigma e^{X^+} \text{---} \text{---} &= -\text{---} - \text{---} - \text{---} - \text{---} - \text{---} \\
&\quad - \text{---} + \text{---} - \text{---} + \text{---} \\
Se^{X^+} \text{---} \text{---} &= 0
\end{aligned}$$

This finishes our considerations of graph homology; we have proved Theorem 1.1.

4.4. The ghost cycle generator δ_+^C . We now turn to an investigation of the ghost sector through cycle homology.

Definition 4.22. Let \mathcal{C}_Γ be the set of cycles in Γ . We write $\mathcal{C}_\Gamma = \{C_1, C_2, \dots\}$.

$$\delta_+^C \Gamma = \sum_{C \in \mathcal{C}_\Gamma} \delta_+^C \Gamma,$$

where

$$\delta_+^C \Gamma = \begin{cases} 0 & \text{if } C \text{ has a vertex which has an adjacent marked or ghost edge} \\ \Gamma_{C \rightsquigarrow \dots} & \text{otherwise.} \end{cases}$$

Note that an (unoriented) ghost cycle is the short-hand notation for the sum of the two orientations:

$$\text{---} = \text{---} + \text{---}.$$

Lemma 4.23. i. For any graph $\Gamma \in \overline{\mathcal{G}}^{n,l}$:

$$\frac{1}{\text{Sym}(\Gamma)} \frac{(\delta_+)^k}{k!} \Gamma = \sum_{\substack{\Gamma' \in \overline{\mathcal{G}}^{n,l} \\ \text{skeleton}(\Gamma') = \Gamma}} \frac{1}{\text{Sym}(\Gamma')} \Gamma'.$$

ii. For any $k \geq 0$, $\frac{(\delta_+)^k}{k!} X^{n,l} = X_{k \text{---}}^{n,l}$.

iii. We have $e^{\delta_+} X^{n,l} = X_{/\text{---}/}^{n,l}$.

Proof. Analogous to Lemma 4.6. □

Example 4.24. An example for Lemma 4.23.i is

$$\frac{1}{2} \delta_+ \text{---} = \frac{1}{2} \left(\text{---} + \text{---} + \text{---} \right) = \text{---} + \frac{1}{2} \text{---}$$

Remark 4.25. The operators χ_+^e and δ_+^C commute, hence so do χ_+ and δ_+ .

Corollary 4.26. i. The combinatorial Green's functions $X_{k \text{---}, \bar{l} \text{---}}^{n,l}$ can be written as

$$X_{k \text{---}, \bar{l} \text{---}}^{n,l} = \frac{\chi_+^k \delta_+^{\bar{l}}}{k! \bar{l}!} X_{0 \text{---}, 0 \text{---}}^{n,l}$$

ii. The full combinatorial Green's function can be written as

$$X_{/\text{---}/, / \text{---} /}^{n,l} = e^{X^+} e^{\delta_+} X_{0 \text{---}, 0 \text{---}}^{n,l}$$

4.5. The cycle differential t .

Definition 4.27. For a graph Γ choose a labelling of the cycles $C_1, C_2, \dots \in \mathcal{C}_\Gamma$. We define a derivation $t : H \rightarrow H$ acting on graphs as:

$$t\Gamma = \sum_{C_i \in \mathcal{C}_\Gamma} (-)^{\#\{C_{i'} \in \mathcal{C}_{\Gamma_{\text{gh}}} \mid i' < i\}} t_{C_i} \Gamma$$

where


$$t_{C_i} \Gamma = \begin{cases} 0 & \text{if } C \text{ has a vertex which has an adjacent marked or ghost edge} \\ \Gamma_{C_i \rightsquigarrow \dots\dots} & \text{otherwise.} \end{cases}$$

Next, we want to distinguish the markings created by δ_+ and t . Therefore we draw the former with little circles instead of dots. We denote: $\mathcal{C}_{\Gamma, \dots\dots} \cup \mathcal{C}_{\Gamma, \dots\dots} = \mathcal{C}_{\Gamma_{\text{gh}}} \subset \mathcal{C}_\Gamma$.


Proposition 4.28. t is a differential: $t^2\Gamma = 0$.

Proof. Analogous to Proposition 4.13. □

Example 4.29. Consider the graph



and label the two cycles:



Then:

$$\begin{aligned} t^2 \text{---} \text{---} &= t \text{---} \text{---} + t \text{---} \text{---} \\ &= \text{---} \text{---} + \text{---} \text{---} = 0. \end{aligned}$$

4.6. The differential T . The T -homology checks that the longitudinal degrees of freedom in a loop through 3-gluon vertices are appropriately matched by ghost loops, so that physical amplitudes are in the kernel of T . Hence, we define

Definition 4.30. A derivation $T : H \rightarrow H$ is given by $T = t + \tau$ where

$$\tau\Gamma = (-)^{\#\mathcal{C}_{\Gamma_{\text{gh}}}} \sum_{C_i \in \mathcal{C}_{\Gamma, \dots\dots}} (-)^{\#\{C_{i'} \in \mathcal{C}_{\Gamma, \dots\dots} \mid i' > i\}} \tau_{C_i} \Gamma$$

and

$$\tau_{C_i} \Gamma = \Gamma_{C_i \rightsquigarrow \dots\dots}$$

Proposition 4.31. T is a differential: $T^2\Gamma = 0$.

Proof. As in Proposition 4.16 this follows from $t\tau\Gamma = -\tau t\Gamma$ and $\tau^2\Gamma = 0$. □

Example 4.32.

$$\begin{aligned} t\tau \text{---} \text{---} &= -t \text{---} \text{---} = \text{---} \text{---} \\ \tau t \text{---} \text{---} &= -\tau \text{---} \text{---} = \text{---} \text{---} \\ \tau^2 \text{---} \text{---} &= -\tau \text{---} \text{---} = 0 \\ T^2 \text{---} \text{---} &= 0 \end{aligned}$$

Example 4.33.

$$\begin{aligned} t\tau \text{---} \text{---} &= -t \text{---} \text{---} + t \text{---} \text{---} = 0 \\ \tau t \text{---} \text{---} &= 0 \\ \tau^2 \text{---} \text{---} &= -\tau \text{---} \text{---} + \tau \text{---} \text{---} = \text{---} \text{---} + \text{---} \text{---} = 0 \\ T^2 \text{---} \text{---} &= 0 \end{aligned}$$

Proposition 4.34. *Let Γ be a graph without ghost edges. Then $Te^{\delta+}\Gamma = 0$.*

Proof. Analogous to Proposition 4.20. □

Symmetry factors are no issue in the following example as we sum over both orientations for the two ghost lines.

Example 4.35.

$$\begin{aligned}
 e^{\delta+} &= \text{diagram 1} + \text{diagram 2} + \text{diagram 3} + \text{diagram 4} \\
 te^{\delta+} &= \text{diagram 5} + \text{diagram 6} - \text{diagram 7} + \text{diagram 8} \\
 \tau e^{\delta+} &= -\text{diagram 9} - \text{diagram 10} - \text{diagram 11} + \text{diagram 12} \\
 Te^{\delta+} &= 0
 \end{aligned}$$

This homology ensures that longitudinal degrees of freedom propagating in loops cancel. We summarize:

Theorem 4.36. *Let Γ be a graph without marked and ghost edges. Then*

$$Se^{\delta+}e^{\chi+}\Gamma = 0, \quad \text{and} \quad Te^{\delta+}e^{\chi+}\Gamma = 0.$$

4.7. **The bicomplex.** As $[s, t] = [S, T] = 0$, we get a double complex:

$$\begin{array}{ccccccc}
 & & \vdots & & \vdots & & \\
 & & \downarrow s & & \downarrow s & & \\
 \cdots & \xrightarrow{t} & H_{k,\tilde{l}} & \xrightarrow{t} & H_{k,\tilde{l}+1} & \xrightarrow{t} & \cdots \\
 & & \downarrow s & & \downarrow s & & \\
 \cdots & \xrightarrow{t} & H_{k+1,\tilde{l}} & \xrightarrow{t} & H_{k+1,\tilde{l}+1} & \xrightarrow{t} & \cdots \\
 & & \downarrow s & & \downarrow s & & \\
 & & \vdots & & \vdots & &
 \end{array}$$

Here, H_{\dots} are to be regarded as reflecting the relevant vector space structure only of these spaces. The corresponding Hopf algebras and combinatorial Green functions are discussed now. This bicomplex above and its relation to gauge symmetry and BRST cohomology will be the study of future work.

5. COMBINATORIAL GREEN FUNCTIONS

The Hopf algebras on scalar graphs straightforwardly generalize to gauge theory graphs. In particular, the coproduct acts on the sum of all graphs contributing to a given amplitude —the combinatorial Green function— filtered by the number of 4-valent vertices and the number of ghost loops. Let us make this more precise.

5.1. Gradings on the Hopf algebra. Recall that the Hopf algebra H is graded by the loop number, since the number of loops in a subgraph $\gamma \subset \Gamma$ and in the graph Γ/γ add up to $|\Gamma| \equiv n(\Gamma)$. Another (multi)grading is given by the number of vertices. In order for this to be compatible with the coproduct —creating an extra vertex in the quotient Γ/γ — we say a graph Γ with $E_E(\Gamma)$ external edges, is of multi-vertex-degree (j_3, j_4, \dots) if the number of m -valent vertices is equal to $j_m + \delta_{m, E_E(\Gamma)}$. One can check that this grading is compatible with the coproduct. Moreover, the two degrees are related via $\sum_m (m-2)j_m(\Gamma) = 2|\Gamma|$. This grading can be extended to involve other types of vertices —such as \tilde{j} ghost-gluon vertices— cf. [20] for full details.




FIGURE 1. A 3-regular gluon cycle (left) and an oriented ghost cycle (right)

5.2. **Series of graphs.** As said, from a physical point of view, it is not so interesting to study individual graphs; rather, one considers whole sums of graphs with the same number of external lines. In this section, we will study series of 1PI graphs in the Hopf algebra H :

$$G_0^{k,n} = \sum_{|\Gamma|=n, |E_E(\Gamma)|=k} \Gamma \frac{\text{colour}(\Gamma)}{\text{sym}(\Gamma)},$$

This is the sum of all 1PI 3-regular (0 4-valent vertices) graphs with first Betti number n and k external gluon edges (which fixes the amplitude r under consideration), normalized by their symmetry factors $\text{sym}(\Gamma)$, the rank of their automorphism groups, in the denominator, and also weighted in the numerator by the corresponding colour factor $\text{colour}(\Gamma)$:

$$\text{colour}(\Gamma) := \prod_{v \in V^\Gamma} R_v \prod_{e \in E_I^\Gamma} \delta_{s(e), t(e)}.$$

Here, R_v is determined by a choice of a representation of the gauge group at v , and $s(e), t(e)$ are the vertex labels for source and target of the internal edge e . Typical, R_v is the adjoint representation for gluon self-interactions or the fundamental representation for a gluon interacting with fermionic matter fields.

Similarly, we write $G_j^{k,n}$ for series of graphs which have j 4-valent vertices, with all other vertices 3-valent. Also, we consider external ghost edges and loops. We let $G_{j;\tilde{n}}^{k,\tilde{k},n}$ denote the sum of graphs which have k external gluon edges, \tilde{k} external ghost edges, j 4-valent vertices, with all other vertices 3-valent, and \tilde{n} ghost cycles (Figure 1). We let $\widehat{G}_{j;\tilde{n}}^{k,n}$ be the same sum where we consider all j 4-valent vertices, and all \tilde{n} ghost cycles as marked.

Summarizing, the superscript on G always indicates the external structure of the graphs in the series, whereas the subscripts indicate the 4-vertex degree, the loop degree, or the ghost cycle degree.

We have shown in [21] that we can impose the Slavnov–Taylor identities on the Hopf algebra H , compatibly with the coproduct, equating all of the following formal elements:

$$(17) \quad Q^{k,\tilde{k}} := \left(\frac{G_{/}^{k,\tilde{k},/}}{(G_{/}^{2,0,/})^{k/2} (G_{/}^{0,\tilde{2},/})^{\tilde{k}/2}} \right)^{1/(k+\tilde{k}-2)},$$

independent of the numbers k and \tilde{k} of external gluon and ghost edges, respectively. The thus-defined single formal series $Q \equiv Q^{k,\tilde{k}}$ will play the role of a ‘charge’ element in the Hopf algebra.

Proposition 5.1. *The coproduct on the Green’s functions reads*

$$\Delta(G_{j_3 j_4; \tilde{n}}^{k,n}) = \sum_{\substack{j_m = j'_m + j''_m \\ n = n' + n'' \\ \tilde{n} = \tilde{n}' + \tilde{n}''}} (G^{k,n'} Q^{2n''})_{j'_3 j'_4; \tilde{n}'} \otimes G_{j''_3 j''_4; \tilde{n}''}^{k,n''}$$

with $G_{j_3 j_4; \tilde{n}}^{k,n}$ the above series of graphs of vertex multidegree (j_3, j_4) , first Betti number n and \tilde{n} ghost cycles.

After taking the Slavnov–Taylor identities (17) into account, the coproduct reads on the above series of graphs

$$\Delta(G^{k,n}) = \sum_{n=n'+n''} (G^k Q^{2n''})_{n'} \otimes G^{k,n''}.$$

Remark 5.2. The inclusion of fermions is parallel to the study of ghost edges and loops, and a mere notational exercise.

Another way to describe the Green's function G^k is in terms of so-called grafting operators, defined in terms of 1PI primitive graphs. We start by considering maps $B_+^\gamma : H \rightarrow \text{Aug}$, with Aug the augmentation ideal, which will soon lead us to non-trivial one co-cycles in the Hochschild cohomology of H . They are defined as follows.

$$B_+^\gamma(h) = \sum_{\Gamma \in \langle \Gamma \rangle} \frac{\mathbf{bij}(\gamma, h, \Gamma)}{|h|_\vee} \frac{1}{\max f(\Gamma)} \frac{1}{(\gamma|h)} \Gamma,$$

where $\max f(\Gamma)$ is the number of maximal forests of Γ , $|h|_\vee$ is the number of distinct graphs obtainable by permuting edges of h , $\mathbf{bij}(\gamma, h, \Gamma)$ is the number of bijections of external edges of h with an insertion place in γ such that the result is Γ , and finally $(\gamma|h)$ is the number of insertion places for h in γ [22]. $\sum_{\Gamma \in \langle \Gamma \rangle}$ indicates a sum over the linear span $\langle \Gamma \rangle$ of generators of H .

The sum of the B_+^γ over all primitive 1PI Feynman graphs at a given loop order and with given residue will be denoted by $B_+^{k;n}$, as in [22]. More precisely,

$$B_+^{k;n} = \sum_{\substack{\gamma \text{ prim} \\ |\gamma|=n \\ E_E(\gamma)=k}} \frac{1}{\text{Sym}(\gamma)} B_+^\gamma.$$

With this and the above Proposition, we can show [22, Theorem 5]:

$$(18) \quad G^k = \sum_{l=0}^{\infty} B_+^{k;n} (G^k Q^{2n});$$

$$(19) \quad \Delta(B_+^{k;n} (G^k Q^{2n})) = B_+^{k;n} (G^k Q^{2n}) \otimes \mathbb{I} + (\text{id} \otimes B_+^{k;n}) \Delta(G^k Q^{2n}).$$

Equation (18) is known as the combinatorial Dyson–Schwinger equation, while (19) shows that $B_+^{k;n}$ is a Hochschild cocycle for the Hopf algebra H .

5.3. The generator of ghost loops. We again consider the map $\delta_+ : H \rightarrow H$ that replaces gluon loops in a Feynman graph by ghost loops.

Remark 5.3. In accordance with our previous definition of δ_+ , it becomes an algebra derivation $\delta_+ : H \rightarrow H$ by the assignment

$$\delta_+(\Gamma) = (\tilde{l} + 1) \sum_{g \subset \Gamma} \Gamma_{g \rightarrow \tilde{g}}.$$

for a 1PI Feynman graph Γ at ghost loop order \tilde{l} . The sum is over all oriented 3-regular gluon cycles g , and $\Gamma_{g \rightarrow \tilde{g}}$ denotes the graph Γ with the 3-regular gluon cycle g replaced by a ghost cycle \tilde{g} (cf. Figure 1), of the same orientation. The coefficient $a_{\Gamma, g}$ is given by the number of (equivalence classes of) pairs (Γ', g') , $g' \subset \Gamma'$ that give rise to the same Feynman graph, i.e. such that

$$\Gamma'_{g' \rightarrow \tilde{g}'} \sim \Gamma_{g \rightarrow \tilde{g}}.$$

The notation δ_+ suggests that there is also a δ_- . In fact, such an operator can be defined and would replace a ghost loop by a gluon loop. We will not further study such an operator, since our interest lies in generating physical amplitudes from zero-ghost-loop amplitudes.

Example 5.4. Consider the following one-loop gluon self-energy graph:

$$\Gamma = \text{---} \bigcirc \text{---}.$$

Its symmetry factor is $\text{Sym}(\Gamma) = 2$ so that

$$\delta_+ \left(\text{---} \bigcirc \text{---} \right) = 2 \text{---} \bigcirc \text{---}.$$

A two loop example is given by the graph

$$\Gamma' = \text{---} \langle \text{---} \bigcirc \text{---} \bigcirc \text{---} \rangle \text{---}$$

for which $\text{Sym}(\Gamma') = 2$. Now,

$$\delta_+ \left(\text{---} \langle \text{---} \bigcirc \text{---} \bigcirc \text{---} \rangle \text{---} \right) = 2 \text{---} \langle \text{---} \bigcirc \text{---} \rangle \text{---} + 2 \text{---} \langle \text{---} \bigcirc \text{---} \rangle \text{---} + 2 \text{---} \langle \text{---} \bigcirc \text{---} \rangle \text{---} .$$

Our previous results on δ_+ Lemma 4.9 allow us to conclude:

Proposition 5.5. *When acting on series of graphs with no ghost cycles ($\tilde{n} = 0$):*

$$e^{\delta_+} \circ B_+^{k;n} = \sum_{\substack{\gamma \text{ prim} \\ |\gamma|=n, \tilde{n}(\gamma)=0 \\ E_E(\gamma)=k}} \frac{1}{\text{Sym}(\gamma)} B_+^{e^{\delta_+}(\gamma)} \circ e^{\delta_+}$$

where the sum is over graphs γ with no ghost cycles.

Remark 5.6. There is a similar result for connected graphs on the exponentiation of χ_+ . We give it here without proof. It follows directly though from extending the definition of graph Hopf algebras and their Hochschild cohomology from 1PI to connected graphs. When acting on series of graphs with no marked edges:

$$e^{\chi_+} \circ B_+^{k;n} = \sum_{\substack{\gamma \text{ prim} \\ |\gamma|=n, j(\gamma)=0 \\ E_E(\gamma)=k}} \frac{1}{\text{Sym}(\gamma)} B_+^{e^{\chi_+}(\gamma)} \circ e^{\chi_+}$$

where the sum is over graphs γ with no marked edges and $j(\gamma)$ is the number of 4-valent vertices.

Together, the two results on the interplay of Hochschild cohomology and exponentiation show that gauge invariant combinatorial Green functions are obtained from gauge invariant skeleton graphs into which gauge invariant subgraphs are inserted.

Example 5.7. Let us consider the example of the gluon self-energy at two loops:

$$(20) \quad G_{n=2}^2 = +\frac{1}{6} \text{---} \langle \text{---} \bigcirc \text{---} \bigcirc \text{---} \rangle \text{---} + \frac{1}{2} \text{---} \langle \text{---} \bigcirc \text{---} \rangle \text{---} + \frac{1}{2} \text{---} \langle \text{---} \bigcirc \text{---} \rangle \text{---} \\ + \frac{1}{2} \text{---} \langle \text{---} \bigcirc \text{---} \bigcirc \text{---} \rangle \text{---} + \text{---} \langle \text{---} \bigcirc \text{---} \rangle \text{---} + \text{---} \langle \text{---} \bigcirc \text{---} \rangle \text{---} + \text{---} \langle \text{---} \bigcirc \text{---} \rangle \text{---} \\ + \frac{1}{2} \text{---} \langle \text{---} \bigcirc \text{---} \bigcirc \text{---} \rangle \text{---} + \text{---} \langle \text{---} \bigcirc \text{---} \rangle \text{---} + \text{---} \langle \text{---} \bigcirc \text{---} \rangle \text{---} + \text{---} \langle \text{---} \bigcirc \text{---} \rangle \text{---}$$

whose zero-ghost-loop part is

$$(21) \quad G_{n=2, \tilde{n}=0}^2 = \frac{1}{6} \text{---} \langle \text{---} \bigcirc \text{---} \bigcirc \text{---} \rangle \text{---} + \frac{1}{2} \text{---} \langle \text{---} \bigcirc \text{---} \rangle \text{---} + \frac{1}{2} \text{---} \langle \text{---} \bigcirc \text{---} \rangle \text{---} \\ + \frac{1}{2} \text{---} \langle \text{---} \bigcirc \text{---} \bigcirc \text{---} \rangle \text{---} + \frac{1}{2} \text{---} \langle \text{---} \bigcirc \text{---} \rangle \text{---}$$

One readily checks that $(1 + \delta_+)(G_{\tilde{n}=0}^{n=2})^2 = (G^{n=2})^2$.

Theorem 5.8. *Let \tilde{H} be the Hopf subalgebra of H generated by $G^{k,n}$ for all $n \geq 0$ and $k = 2, 3, 4$. Then $\exp \delta_+$ is an automorphisms of the graded Hopf algebra \tilde{H} :*

$$\exp \delta_+(x_1 x_2) = \exp \delta_+(x_1) \exp \delta_+(x_2); \quad \Delta(\exp \delta_+(x)) = (\exp \delta_+ \otimes \exp \delta_+) \Delta(x).$$

for $x_1, x_2, x \in \tilde{H}$.

Proof. By definition, δ_+ is an algebra derivation so that $\exp \delta_+$ is an algebra automorphism. Note that at a given loop order l , the exponential terminates at that power n and is thus well-defined on the graded algebra underlying \tilde{H} .

Let us then consider the compatibility of δ_+ with the coproduct structure. Recall from [20] the formula

$$\Delta(G^k) = \sum_{j_3, j_4, \tilde{j} \geq 0} G^k(Q^3)^{j_3}(Q^4)^{2j_4}(Q^{1, \tilde{2}})^{\tilde{n}} \otimes G^k_{j_3 j_4 \tilde{j}}.$$

which holds even without the Slavnov–Taylor identities. It continues to hold when restricting to graphs with zero ghost loops:

$$\Delta(G^k_{\tilde{n}=0}) = \sum_{j_3, j_4} G^k_{\tilde{n}=0}(Q^3_{\tilde{n}=0})^{j_3}(Q^4_{\tilde{n}=0})^{2j_4} \otimes G^r_{j_3, j_4; \tilde{n}=0}.$$

We now apply $\exp \delta_+ \otimes \exp \delta_+$ to this equation to obtain after imposing the Slavnov–Taylor identities $Q^3 = Q^4$:

$$\begin{aligned} (\exp \delta_+ \otimes \exp \delta_+) \Delta(G^k_{\tilde{n}=0}) &= \sum_{j_3, j_4} G^k(Q^3)^{j_3}(Q^4)^{2j_4} \otimes \exp \delta_+ (G^k_{j_3 j_4; \tilde{n}=0}) \\ &= \sum_{n \geq 0} G^k Q^{2n} \otimes \exp \delta_+ (G^k_{n, \tilde{n}=0}) \end{aligned}$$

since in the absence of ghost vertices $j_3 + 2j_4 = 2n$ in terms of the first Betti number n . Lemma(4.23) then yields $\exp \delta_+(G^k_{n, \tilde{n}=0}) = G^k_n$, which completes the proof. \square

This can be extended to the *connected* Green’s functions $X_{k,n}$, where also a similar result can be shown for $\exp \chi_+$.

Example 5.9. First, recall the Slavnov–Taylor identities $G^3 G^{\tilde{2}} = G^{1, \tilde{2}} G^2$ which at one-loop order become:

We compute $\Delta'(G^{2, n=2}_{\tilde{n}=0})$ with $G^{2, n=2}_{\tilde{n}=0}$ given in Eq.(21). For the first graph on the last line, we have

$$\Delta' \left(\frac{1}{2} \text{[Diagram]} \right) = \frac{1}{2} \text{[Diagram]} \otimes \text{[Diagram]}$$

If we apply $\exp \delta_+ \otimes \exp \delta_+$ to this expression, we obtain

$$\frac{1}{2} \left(\text{[Diagram]} + 2 \text{[Diagram]} \right) \otimes \left(\text{[Diagram]} + 2 \text{[Diagram]} \right)$$

For the coproduct on the last graph in Eq.(21) we have

$$\Delta' \left(\frac{1}{2} \text{[Diagram]} \right) = \text{[Diagram]} \otimes \text{[Diagram]}$$

and applying $\exp \delta_+ \otimes \exp \delta_+$ to this expression yields

$$\left(\text{[Diagram]} + \text{[Diagram]} + \text{[Diagram]} \right) \otimes \left(\text{[Diagram]} + 2 \text{[Diagram]} \right)$$

On the other hand, $\Delta'(\exp \delta_+ G_{\tilde{n}=0}^{2,n=2}) = \Delta'(G^{2,n=2})$ is computed from Eq.(20):

$$\begin{aligned} \Delta'(G^{2,n=2}) = & \left(\frac{1}{2} \text{---} \bigcirc \text{---} + \text{---} \bigcirc \text{---} \right) \otimes \text{---} \bigcirc \text{---} \\ & + 2 \text{---} \text{---} \otimes \text{---} \bigcirc \text{---} \\ & + \left(\text{---} \bigcirc \text{---} + \text{---} \triangle \text{---} + \text{---} \triangle \text{---} \right) \otimes \text{---} \bigcirc \text{---} \\ & + \left(2 \text{---} \triangle \text{---} + 2 \text{---} \triangle \text{---} \right) \otimes \text{---} \bigcirc \text{---} \end{aligned}$$

We conclude that

$$\begin{aligned} (\exp \delta_+ \otimes \exp \delta_+) \Delta'(G_{\tilde{n}=0}^{2,n=2}) - \Delta'(\exp \delta_+(G_{\tilde{n}=0}^{2,n=2})) = \\ 2 \left(\frac{1}{2} \text{---} \bigcirc \text{---} + \text{---} \bigcirc \text{---} + \text{---} \triangle \text{---} + \text{---} \triangle \text{---} \right. \\ \left. - \text{---} \text{---} + \text{---} \bigcirc \text{---} - \text{---} \triangle \text{---} - \text{---} \triangle \text{---} \right) \otimes \text{---} \bigcirc \text{---} \end{aligned}$$

which vanishes by the Slavnov–Taylor identities upon adding the contribution of 4-valent vertices.

6. THE COROLLA POLYNOMIAL AND DIFFERENTIALS

6.1. The Corolla Polynomial. Finally, we introduce the Corolla Polynomial ([10]). We need the following definitions:

- For a vertex $v \in V$ let $n(v)$ be the set of edges incident to v (internal or external).
- For a vertex $v \in V$ let $D_v = \sum_{j \in n(v)} a_{v,j}$.
- Let \mathcal{C} be the set of all cycles of Γ (cycles, not circuits). This is a finite set.
- For C a cycle and v a vertex in V , since Γ is 3-regular, there is a unique edge of Γ incident to v and not in C , let v_C be this edge.
- For $i \geq 0$ let

$$C^i = \sum_{\substack{C_1, C_2, \dots, C_i \in \mathcal{C} \\ C_j \text{ pairwise disjoint}}} \left(\left(\prod_{j=1}^i \prod_{v \in C_j} a_{v, v_C} \right) \prod_{v \notin C_1 \cup C_2 \cup \dots \cup C_i} D_v \right)$$

- Let

$$C = \sum_{j \geq 0} (-1)^j C^j$$

For any finite graph Γ , this is a polynomial $C = C(\Gamma)$ —the corolla polynomial— because $C^i = 0$ for $i > |\mathcal{C}|$.

Theorem 6.1. ([10]) *Let \mathcal{T}^Γ be the set of sets T of half edges of Γ with the property that*

- *every vertex of Γ is incident to exactly one half edge of T*
- *$\Gamma \setminus T$ has no cycles*

Then

$$C(\Gamma) = \sum_{T \in \mathcal{T}^\Gamma} \prod_{h \in T} a_h$$

Remark 6.2. This shows that the corolla polynomial is strictly positive. As it applies in this form as a corolla differential to pure Yang–Mills theory, this results in a positivity statement on Yang–Mills theory which does not hold for gauge fields coupled to matter fields. Accordingly, the sign of the β -function in gauge theory becomes dependent on the number of fermion families, and their representations.

Remark 6.3. For a graph Γ , let E be a set of pairwise disjoint internal edges of Γ . For $i \geq 0$ let

$$C_E^i(\Gamma) = \sum_{\substack{C_1, C_2, \dots, C_i \in \mathcal{C} \\ C_j \text{ pairwise disjoint} \\ C_j \cap E = \emptyset}} \left(\left(\prod_{j=1}^i \prod_{v \in C_j} a_{v, v_C} \right) \prod_{v \notin C_1 \cup C_2 \cup \dots \cup C_i \cup E} D_v \right)$$

where the sum forbids cycles from sharing either vertices or edges with E .

Let

$$C_E(\Gamma) = \sum_{j \geq 0} (-1)^j C_E^j(\Gamma).$$

Then,

$$C_E(\Gamma) = C(\Gamma - E)$$

where $\Gamma - E$ is the graph with the edges and vertices involved in E removed. Removing a vertex removes all its incident half-edges so that $2|E|$ new external edges are generated. Note that $C_\emptyset(\Gamma) = C(\Gamma)$.

Define

$$C^{\text{fr}}(\Gamma) := \sum_E \left(C_E(\Gamma) \prod_{e \in E} W_e \right),$$

where W_e is defined in (7).

Corollary 6.4.

$$C^{\text{fr}}(\Gamma) = \sum_E \left(\sum_{T \in \mathcal{T}^E} \left(\prod_{h \in T} a_h \right) \prod_{e \in E} W_e \right).$$

Proof. Immediate. □

Remark 6.5. Consider a 3-regular graph Γ which has j , $j \geq 2$, 3-valent vertices, and let \mathcal{P} be a set of m paths, $2m \leq j$, on internal edges and 3-valent vertices in Γ which each connect two external 3-valent vertices (a 3-valent vertex v is external if $n(v)$ contains an external edge) with $p_i \cap p_j = \emptyset$, $\forall p_i, p_j \in \mathcal{P}$.

Consider for chosen set E and \mathcal{P} as above, with $E \cap \mathcal{P} = \emptyset$, for $i \geq 0$,

$$C_{E, \mathcal{P}}^i(\Gamma) = \sum_{\substack{C_1, C_2, \dots, C_i \in \mathcal{C} \\ C_j \text{ pairwise disjoint} \\ C_j \cap E = \emptyset \\ C_j \cap \mathcal{P} = \emptyset}} \left(\left(\prod_{j=1}^i \prod_{v \in C_j} a_{v, v_C} \right) \prod_{v \notin C_1 \cup C_2 \cup \dots \cup C_i \cup E \cup \mathcal{P}} D_v \right) \times \\ \times \left(\prod_{p \in \mathcal{P}} \prod_{v \in p} a_{v, v_p} \right),$$

where the sum forbids cycles from sharing either vertices or edges with E , and v_p is the unique half-edge at v not in p .

Let

$$C_{E, \mathcal{P}}(\Gamma) = \sum_{j \geq 0} (-1)^j C_{E, \mathcal{P}}^j(\Gamma).$$

Finally, we set

$$(22) \quad C_{\mathcal{P}}^{\text{fr}}(\Gamma) := \sum_E \left(C_{E, \mathcal{P}}(\Gamma) \prod_{e \in E} W_e \right).$$

6.2. Corolla differentials. Our main use of the corolla polynomial is to construct differential operators with it. These operators differentiate with respect to momenta ξ_e assigned to edges e of a graph, and act on the second Kirchhoff polynomial written for generic edge momenta ξ_e , that is on $|N|_{\text{Pf}}$.

Only at the end of the computation will we employ the map

$$Q : \xi_e \rightarrow \xi_e + q(e).$$

We then set $\xi_e = 0$ after we have applied the corolla differentials so that we obtain the standard second Symanzik polynomial for specific external momenta as prescribed by gauge theory amplitudes.

For a half edge $h \equiv (w, f) \in H^\Gamma$, we let $e(h) = f$ and $v(h) = w$. We remind the reader that h_+ and h_- are the successor and the precursor of h in the oriented corolla at $v(h)$, and that we assign to a graph Γ :

- i. to each (possibly external) edge e , a variable A_e and a 4-vector ξ_e ;
- ii. to each half edge h , a Lorentz index $\mu(h)$;
- iii. a factor $\text{colour}(\Gamma)$.

6.3. The differential D^0 . The corolla polynomial is an alternating sum over terms C^i , where i counts the number of loops. Similarly, the corolla differentials are a sum of terms D^i . We start with D^0 .

Let $\Gamma \in \mathcal{G}^{n,l}$:

$$U^0(\Gamma) = \int_E \frac{d\mathbf{k}_L}{(2\pi)^{dL}} C_\Gamma^0(\underline{D}) e^{-\sum_{e \in \Gamma^{[1]}} A_e \xi_e'^2},$$

where

$$C_\Gamma^0(\underline{D}) = \prod_{v \in \Gamma^{[0]}} D_v, \quad D_v = D_{v1} + D_{v2} + D_{v3}$$

(the edges incident on v are labelled 1, 2, 3),

$$D_{v1} = -\frac{1}{2} g^{\mu_2 \mu_3} \left(\varepsilon_{v2} \frac{1}{A_2} \frac{\partial}{\partial \xi_{2\mu_1}} - \varepsilon_{v3} \frac{1}{A_3} \frac{\partial}{\partial \xi_{3\mu_1}} \right).$$

Using that all corollas are oriented, we can write this as


$$D_g(h) := -\frac{1}{2} g^{\mu_{h_+} \mu_{h_-}} \left(\varepsilon_{h_+} \frac{1}{A_{e(h_+)}} \frac{\partial}{\partial \xi_{(h_+)\mu_h}} - \varepsilon_{h_-} \frac{1}{A_{e(h_-)}} \frac{\partial}{\partial \xi_{(h_-)\mu_h}} \right),$$

for any half-edge h . The operator D_v is such that if it acts on $e^{-\sum_{e \in \Gamma^{[1]}} A_e \xi_e'^2}$, it gives the 3-vertex Feynman rule of Eq.(10):

$$D_v e^{-\sum_{e \in \Gamma^{[1]}} A_e \xi_e'^2} = V_v^{(3)} e^{-\sum_{e \in \Gamma^{[1]}} A_e \xi_e'^2}.$$

In order to calculate $C_\Gamma^0(\underline{D}) \equiv C_\Gamma^0(h \rightarrow D_g(h))$, we also need to know the Leibniz terms $D_v V_w^{(3)}$, where $v, w \in \Gamma^{[0]}$.

- If v and w do not share an edge, $D_v V_w^{(3)} = 0$.
- Suppose they share exactly one edge; we give it label 5. Let 1 and 2 be the other edges at v and 3 and 4 the other ones at w :




Then:

$$D_v V_w^{(3)} = \frac{1}{A_5} (g^{\mu_4 \mu_2} g^{\mu_3 \mu_1} - g^{\mu_2 \mu_3} g^{\mu_4 \mu_1}) \equiv \frac{W_e}{A_e},$$

where W_e is the Feynman rule for a marked edge (equation (7)). Note that thus

$$D_v V_w^{(3)} = D_w V_v^{(3)}.$$

- Suppose that v and w share two edges, 3 and 4. Let 1 be the other edge at v and 2 the other one at w :



Then:

$$\begin{aligned} D_v V_w^{(3)} &= \left(\frac{1}{A_3} (g^{\mu_4 \mu_4} g^{\mu_2 \mu_1} - g^{\mu_2 \mu_4} g^{\mu_4 \mu_1}) + \frac{1}{A_4} (g^{\mu_3 \mu_3} g^{\mu_2 \mu_1} - g^{\mu_3 \mu_2} g^{\mu_3 \mu_1}) \right) \\ &= \frac{W_3}{A_3} + \frac{W_4}{A_4}. \end{aligned}$$

where we have used equation (7). Note that also in this case

$$D_v V_w^{(3)} = D_w V_v^{(3)}.$$

Contracting the indices further gives self-loops which can be omitted:

$$D_v V_w^{(3)} = 3C_2 \delta^{a_1 a_2} g^{\mu_1 \mu_2} \left(\frac{1}{A_3} + \frac{1}{A_4} \right).$$

6.4. Regular terms and residues. We can now compute immediately the application of D^0 to the scalar integrand I_Γ , that is, $U^0(\Gamma) := D^0 I_\Gamma$.

$$\begin{aligned} U^0(\Gamma) &= \int \frac{d\underline{k}_L}{(2\pi)^{dL}} \left(\prod_{v \in \Gamma^{[0]}} D_v \right) e^{-\sum_{e \in \Gamma^{[1]}} A_e \xi_e'^2} \\ &= \int \frac{d\underline{k}_L}{(2\pi)^{dL}} \left[\left(\prod_{v \in \Gamma^{[0]}} V_v^{(3)} \right) \right. \\ &\quad + \sum_{\substack{w, w' \in \Gamma^{[0]} \\ w \text{ and } w' \text{ share an edge} \\ w < w'}} (D_w V_{w'}^{(3)}) \left(\prod_{\substack{v \in \Gamma^{[0]} \\ v \neq w, w'}} V_v^{(3)} \right) \\ &\quad + \sum_{\substack{w, w', x, x' \in \Gamma^{[0]} \\ w \text{ and } w' \text{ and } x \text{ and } x' \text{ share an edge} \\ w < w' < x < x'}} (D_w V_{w'}^{(3)}) (D_x V_{x'}^{(3)}) \left(\prod_{\substack{v \in \Gamma^{[0]} \\ v \neq w, w', x, x'}} V_v^{(3)} \right) \\ &\quad \left. + \dots \right] e^{-\sum_{e \in \Gamma^{[1]}} A_e \xi_e'^2}. \end{aligned}$$

With the result of the previous subsection we get (recall that we exclude graphs with self-loops):

$$\begin{aligned} U^0(\Gamma) &= \int \frac{d\underline{k}_L}{(2\pi)^{dL}} \left[\left(\prod_{v \in \Gamma^{[0]}} V_v^{(3)} \right) + \sum_{e \in \Gamma_{\text{int}}^{[1]}} \frac{W_e}{A_e} \left(\prod_{\substack{v \in \Gamma^{[0]} \\ (v, e) \notin \Gamma^{[3]}}} V_v^{(3)} \right) \right. \\ &\quad + \sum_{\substack{\{e_1, e_2\} \subset \Gamma_{\text{int}}^{[1]} \\ e_1 \text{ and } e_2 \text{ do not share a vertex}}} \frac{W_{e_1} W_{e_2}}{A_{e_1} A_{e_2}} \left(\prod_{\substack{v \in \Gamma^{[0]} \\ (v, e_1), (v, e_2) \notin \Gamma^{[3]}}} V_v^{(3)} \right) \\ &\quad \left. + \dots \right] e^{-\sum_{e' \in \Gamma^{[1]}} A_{e'} \xi_{e'}'^2}. \\ &= \sum_{k \geq 0} \sum_{\substack{\{e_1, \dots, e_k\} \subset \Gamma_{\text{int}}^{[1]} \\ e_1, \dots, e_k \text{ do not share a vertex}}} \frac{W_{e_1} \dots W_{e_k}}{A_{e_1} \dots A_{e_k}} \int \frac{d\underline{k}_L}{(2\pi)^{dL}} \\ &\quad \times \left(\prod_{\substack{v \in \Gamma^{[0]} \\ (v, e_1), \dots, (v, e_k) \notin \Gamma^{[3]}}} V_v^{(3)} \right) e^{-\sum_{e \in \Gamma^{[1]}} A_e \xi_e'^2}. \end{aligned}$$

The first term we recognise as the Feynman-Schwinger integrand of Γ . The other terms we can write as the integrands of marked versions of Γ (equation (9)). More precisely,

$$U^0(\Gamma) = \sum_{k \geq 0} \sum_{\{e_1, \dots, e_k\} \subset \Gamma_{\text{int}}^{[1]}} \frac{1}{A_{e_1} \cdots A_{e_k}} \int \frac{d\underline{k}_L}{(2\pi)^{dL}} \mathcal{I}(\chi_+^{e_1} \cdots \chi_+^{e_k} \Gamma) e^{-A_{e_1} \xi_{e_1}'^2 - \cdots - A_{e_k} \xi_{e_k}'^2},$$

where $\mathcal{I}(\Gamma)$ is given in equation 11. Recall that in the exponent in the integrand only the unmarked edges are included. That is why the factor $e^{-A_{e_1} \xi_{e_1}'^2 - \cdots - A_{e_k} \xi_{e_k}'^2}$ appears. This factor does not change the residue along $\prod_{e \in \Gamma_{\text{int}}^{[1]}} A_e = 0$.

Each subset of edges here is accompanied by a corresponding set of poles. By construction, the residues along these poles correspond to integrands where the edges shrink to form 4-valent vertices with the correct Feynman rules.

Using the χ_+^e -operator, we can write the integral $\tilde{U}^0(\Gamma)$ as:

$$\tilde{U}^0(\Gamma) = \sum_{k \geq 0} \sum_{\{e_1, \dots, e_k\} \subset \Gamma_{\text{int}}^{[1]}} \int d\underline{A}_{\Gamma^{[1]} \setminus \{e_1, \dots, e_k\}} I(\chi_+^{e_1} \cdots \chi_+^{e_k} \Gamma).$$

In terms of Feynman amplitudes, this is

$$(23) \quad \tilde{U}^0(\Gamma) = \sum_{k \geq 0} \sum_{\{e_1, \dots, e_k\} \subset \Gamma_{\text{int}}^{[1]}} \Phi(\chi_+^{e_1} \cdots \chi_+^{e_k} \Gamma) = \Phi(e^{\chi^+} \Gamma).$$

Instead of applying \tilde{U}^0 to a single graph, we can do this to the combinatorial Green's function $X^{n,l}$. This gives us the Green's function for all graphs in Yang-Mills theory without the ghosts, but including the 4-valent vertices:

Proposition 6.6. *Collecting residues as above produces the evaluation by the Feynman rules of all 3- and 4-valent graphs in gauge theory without internal ghost or fermion edges:*

$$\tilde{U}^0(X^{n,l}) = \Phi(e^{\chi^+} X^{n,l}) = \Phi(X^{n,l}/\chi_+^{\chi}).$$

Proof. The above equation (23) is used, together with Lemma 4.9.iii. □

6.5. Exponentiating residues. Let us discuss the pairing between the integrand with poles along the boundaries of the simplex, with boundaries given by $\sigma_\Gamma : \prod_{i=1}^{|\Gamma_I^{[1]}|} A_i = 0$, and the Feynman integrand $U^0(\Gamma)$ in more detail.

The amplitude $\tilde{U}^0(\Gamma)$ can be obtained from $U^0(\Gamma)$ by taking residues along hypersurfaces $\prod_{e \in E} A_e = 0$ and regular parts and integrating:

$$\tilde{U}^0(\Gamma) = \sum_{k \geq 0} \sum_{\{e_1, \dots, e_k\} \subset \Gamma_{\text{int}}^{[1]}} \int d\underline{A}_{\Gamma^{[1]} \setminus \{e_1, \dots, e_k\}} \underset{A_1, \dots, \widehat{A_{e_1}}, \dots, \widehat{A_{e_k}}, \dots = 0}{\text{Reg}} \underset{A_{e_1}, \dots, A_{e_k} = 0}{\text{Res}} U^0(\Gamma).$$

For a function $f = f(\{A_e\})$ of graph polynomial variables $A_e, e \in \gamma_I^{[1]}$ with at most simple poles at the origin localized in disjoint sets of edges E , we can write

$$f = \sum_E f^E,$$

where the sum is over all such sets and f^E is the part of f which is regular upon setting variables $A_e, e \in (\Gamma_I^{[1]} - E)$ to zero.

For any set E of mutually disjoint internal edges of Γ , consider $\prod_{e \in E} \oint_{\gamma_e} f$, and let f^E be its regular part. For any finite graph Γ , let \mathcal{E}^γ be the set of all sets of mutually disjoint edges (\emptyset included).

Consider the differential form

$$J_\Gamma^f := \left(f^E \bigwedge_{e \in (\Gamma_I^{[1]} - E)} dA_e \right)_{E \in \mathcal{E}^\Gamma}.$$

Let M_Γ^E be the hypercube

$$M_\Gamma^E := \mathbb{R}_+^{|\Gamma_I^{[1]}| - |E|},$$

and the corresponding vector

$$H_\Gamma(M_\Gamma^E)_{E \in \mathcal{E}^\Gamma}.$$

Then, there is a natural pairing

$$\int H_\Gamma \cdot J_\Gamma^f := \sum_{E \in \mathcal{E}^\Gamma} \int_{M_\Gamma^E} \left(f^E \bigwedge_{e \in (\Gamma_I^{[1]} - E)} dA_e \right).$$

6.6. Graph homology and the residue map. Note that in parametric integration we integrate against the simplex $\sigma \equiv \sigma_\Gamma$ with boundary $\prod_{e \in \Gamma^{[1]}} A_e = 0$. We have co-dimension k -hypersurfaces given by

$$A_{i_1} = \dots = A_{i_k} = 0.$$

The Feynman integrand we have constructed above comes from regular parts, and residues along these hypersurfaces. It can be described by the following commutative diagram.

$$\begin{array}{ccc} \mathcal{G}_j \mathfrak{X}_{\mathcal{I}, \mathcal{I}'} \ni \Gamma & \xrightarrow{\chi_+} & \chi_+(\Gamma) \in \mathcal{G}_{(j+1)} \mathfrak{X}_{\mathcal{I}, \mathcal{I}'} \\ \downarrow \Phi & & \downarrow \Phi \\ \Phi(\Gamma) & \xrightarrow{\sum_e \text{Res}_e} & \Phi(\chi_+(\Gamma)) \end{array}$$

The underlying geometry will be interpreted elsewhere.

6.7. Covariant gauges. For an edge e , let

$$G_{\mu\nu}^\rho(e) := \frac{g_{\mu\nu}}{\xi_e'^2} - 2\rho \frac{\xi_{e\mu}' \xi_{e\nu}'}{\xi_e'^4}$$

the corresponding gluon propagator in a covariant gauge ($\rho = 1/2$ being the transversal gauge, $\rho = 0$ the Feynman gauge). One computes

$$G_{\mu\nu}^\rho(e) = \int_0^\infty \frac{-1}{2A\rho} \frac{\partial}{\partial \xi_{e\mu}'} \frac{\partial}{\partial \xi_{e\nu}'} e^{-\rho A \xi_e'^2} dA =: \int_0^\infty g_{\mu\nu}^\rho(e) dA.$$

We set

$$G_\Gamma^\rho := \prod_{e \in \Gamma_I^{[1]}} G_{\mu(s(e), e) \mu(t(e), e)}^\rho(e),$$

for half-edges $(s(e), e)$ and $(t(e), e)$, and g_Γ^ρ accordingly.

We let $I_\Gamma(\rho)$ be the corresponding scalar integrand obtained by substituting $A_e \rightarrow \rho A_e$ for each internal edge e .

G_Γ^ρ acts as a differential operator so that

$$G_\Gamma^\rho I_\Gamma(\rho) = F_G(\rho) I_\Gamma(\rho),$$

with $F_G(\rho)$ a polynomial in ρ , edge variables A_e and 4-momenta ξ_e .

Similarly, the corolla differential D_Γ acts as a differential operator so that

$$D_\Gamma I_\Gamma(1) = F_D I_\Gamma(1),$$

with F_D a polynomial in the 4-momenta ξ_e and a rational function in the edge variables A_e .

To compute in an arbitrary covariant gauge, we then work with

$$F_G(\rho) F_D I_\Gamma(1).$$

6.8. **Yang–Mills theory.** Consider a cycle C through 3-valent vertices in a graph Γ , and consider

$$\prod_{v \in C} D_g(v_C).$$

This is a differential operator with coefficients which are monomials in variables $1/A_e$, where $e \in C$.

Let D_C be the part in this differential operator which is linear in all variables $1/A_e$, for $e \in C$. Let ϕ_C be the Feynman rule for a ghost loop on C , summed over both orientations.

Lemma 6.7.

$$\phi_C := D_C e^{-\sum_{e \in C} A_e \xi_e'^2}.$$

Proof. This follows directly from the Feynman rules for ghost propagators and ghost-gluon vertices. Linearization eliminates all poles with residues corresponding to 4-valent 2-ghost-2-gluon vertices. \square

Now consider the corolla polynomial $C(\Gamma)$ and replace each half-edge variable h by the differential $D_g(h)$. This defines a differential operator

$$d(\Gamma)^{\text{YM}} := C(\Gamma)(h \rightarrow D_g(h)),$$

We consider $d(\Gamma)^{\text{YM}}(I_\Gamma)$ where

$$I_\Gamma := \frac{e^{-\frac{|N_\Gamma|_{\text{Pf}}}{\psi_\Gamma}}}{\psi_\Gamma^2},$$

is the scalar integrand for a graph Γ .

Proposition 6.8. *All poles in $d(\Gamma)^{\text{YM}} I_\Gamma$ are located along co-dimension $|E|$ hypersurfaces $A_e = 0$, $e \in E$ for subsets E of mutually disjoint edges are simple poles.*

Proof. Corollary 3.5 ensures that poles are at most of first order and appear only when two derivatives act on the same edge. By the definition of the corolla polynomial this can only appear in mutually disjoint ordered pairs of corollas. All poles coming from divergent subgraphs are located along subsets of connected edges, as divergent subgraphs have more than a single edge. \square

By our previous results on the Leibniz terms we can summarize now for the parametric integrand:

Corollary 6.9. *The residues of these poles correspond to graphs where each corresponding pair of corollas P_e is replaced by a 4-valent vertex.*

Proof. Setting an edge variable to zero shrinks that edge in the two Symanzik polynomials by the standard contraction-deletion identities [2, 3, 23]. \square

The Leibniz terms serve the useful purpose to shrink an edge between two 3-gluon vertices, and have a residue which is the integrand with the corresponding edge becoming a marked edge, and is hence part of the integrand for a graph with a corresponding 4-valent vertex. As we have checked before, summing over all connected 3-regular graphs, we correctly reproduce the Feynman integrand for all gluon self-interactions.

We stress that in so doing we want to shrink edges only between pairs of corollas which both are corollas for 3-gluon vertices, and will not mark edges between other type of vertices. This leads us to

Definition 6.10. We let $D(\Gamma)^{\text{YM}} I_\Gamma$ be the part $d(\Gamma)^{\text{YM}} I_\Gamma$ which is linear in all variables $1/A_e$.

This eliminates all poles in $D(\Gamma)^{\text{YM}} I_\Gamma$ of the form $1/A_e$. We can regain then the contribution of 4-valent 4-gluon vertices by using Theorem 6.1 together with Remark 6.3:

Lemma 6.11. *Let*

$$U_\Gamma = g_\Gamma^{\text{Pf}} C^{\text{tr}}(\Gamma)(a_h \rightarrow D_g(h)) I_\Gamma.$$

Then \bar{U}_Γ (cf. Eq.(5)) generates the integrand for the complete contribution of Γ to the full Yang–Mills theory amplitude. \bar{U}_Γ^R generates the corresponding integrand for the renormalized contribution.

Proof. Immediate application of Lemma 6.7 and Theorem 1.3. \square

This also proves Theorem 1.5 in the context of Yang–Mills theory.

Remark 6.12. If we were to work with non-linear gauges, we could avoid this linearization and use the Leibniz terms for the graphs with 2-gluon 2-ghost and 4-ghost vertices. Also, note that $\overline{U}_\Gamma^R = \overline{U}_\Gamma^R(\rho)$ depends on the gauge parameter.

6.9. Amplitudes with open ghost or fermion lines. For k open ghost lines we have a straightforward generalization of these differentials by using $C_{\mathcal{F}}^{\text{fr}}(G)$, see Eq.(22), where each half edge h is again replaced by $D_g(h)$ and linearization is understood as before. For fermion lines, see below.

6.10. Gauge Theory. If we include matter fields, we need to add a second differential in particular for fermion fields:

$$D_f(h) := \left(\frac{1}{A(e(h_+))} \frac{\partial}{\partial \xi(e(h_+))_{\mu(h_+)}} \gamma_{\mu(h_+)} \gamma_{\mu(h)} - \frac{1}{A(e(h_-))} \frac{\partial}{\partial \xi(e(h_-))_{\mu(h_-)}} \gamma_{\mu(h)} \gamma_{\mu(h_-)} \right).$$

Now we must carefully distinguish between fermion and ghost cycles.

For a collection of cycles C_1, \dots, C_j contributing to C^j , consider partitions of this set into two subsets I_f, I_g containing $|I_f| + |I_g| = j$ cycles. Replace $a_{v,v_C} \rightarrow b_{v,v_C}$ for each $C \in I_f$. This defines $C^{I_g, I_f}(\Gamma)(a_h, b_h)$. Upon summing over all possible partitions I_g, I_f of the cycles for each j , this gives a further corolla polynomial for which we write in slight abuse of notation $C(\Gamma)(a_h, b_h)$. Assign a differential operator as follows:

$$U_\Gamma = g_\Gamma^p \sum_{j \geq 0} \sum_{|I_g| + |I_f| = j} C^{I_g, I_f}(\Gamma)(D_g(h), D_f(h)) \text{colour}^{I_g, I_f}(\Gamma),$$


where in C^{I_g, I_f} , for $I_g \cup I_f \neq \emptyset$, we keep only terms which are linear in variables $1/A_e$ for edges $e \in C_1 \cup \dots \cup C_j$. We can now proceed with \overline{U}_γ as before.

Note that the restriction to $I_l = \emptyset$ gives back the corresponding operator for Yang–Mills theory. From here on, Theorem 1.5 follows for gauge theory as before for Yang–Mills theory.

Remark 6.13. Note that all this can be turned into a projective integrand, illuminating the slots in the period matrix which are filled in a gauge theory as compared to a scalar field theory. In particular, one hopes that the geometry of Eq.(6.6) is helpful to explain appearances and disappearances of periods in gauge theory.

Remark 6.14. Putting fermions into the same colour rep as gauge bosons allows for immediate cancellations between D_g and D_f . This can be illuminating in studying the simplifications for supersymmetric gauge theories.

6.11. Examples: QED and Yang–Mills theory. In the following two examples, we compute the one-loop vacuum polarization in quantum electrodynamics, and then the one-loop gluon vacuum polarization in Yang–Mills theory. Both examples can be obtained from corolla differentials acting on the simplest possible 3-regular graph:




We label its two internal edges 1, 2, and the external edges 3, 4. We also label the two vertices a, b . Edge 3 is oriented from vertex a to vertex b , and edge 4 vice versa, say.

We have six half-edges: $h_1 := (a, 3), h_2 := (a, 2), h_3 := (a, 1)$ and $h_4 := (b, 1), h_5 := (b, 2), h_6 := (b, 4)$, with corresponding half-edge variables a_{a3}, a_{a2} etc.

We have four 4-vectors $\xi_1, \xi_2, \xi_3, \xi_4$, with $\xi_e \in \mathbb{M}^4$, Minkowski space, with scalar product $\xi_e^2 \equiv \xi_e \cdot \xi_e = \xi_{e0}^2 - \xi_{e1}^2 - \xi_{e2}^2 - \xi_{e3}^2$.

Example 6.15. In order to compute the one-loop vacuum polarisation in massless QED,



we proceed as follows. We have for the corolla polynomial

$$C_1(\Gamma) = a_{a_3} a_{b_4}.$$

The scalar integrand is

$$I(\Gamma) = \frac{1}{2} \xi_3^2 \xi_4^2 \frac{e^{-\frac{(\xi_1 - \xi_2)^2 A_1 A_2 + (A_3 \xi_3^2 + A_4 \xi_4^2)(A_1 + A_2)}{A_1 + A_2}}}{(A_1 + A_2)^2} dA_1 dA_2 dA_3 dA_4.$$

We can directly integrate A_3, A_4 eliminating any appearance of ξ_3^2, ξ_4^2 as in this example no derivatives with respect to external edges appear in the corolla differential.

Indeed, replacing the two half-edge variables in $C_1(\Gamma)$ by the fermion differential and using the linearized corolla differential (we symmetrize below in $\mu(3), \mu(4)$ when allowed)

$$\frac{1}{4A_1 A_2} \left(\frac{\partial}{\partial \xi_{3\mu(3)}} \frac{\partial}{\partial \xi_{4\mu(4)}} + \frac{\partial}{\partial \xi_{4\mu(4)}} \frac{\partial}{\partial \xi_{3\mu(3)}} \right) = \frac{1}{2A_1 A_2} \frac{\partial}{\partial \xi_{3\mu(3)}} \frac{\partial}{\partial \xi_{4\mu(4)}}$$

delivers π_1 , the integrand for Π_1 :

$$\begin{aligned} \pi_1 &:= -\frac{1}{4} \text{Tr}(\gamma_{\mu(3)} \gamma_{\mu(2)} \gamma_{\mu(4)} \gamma_{\mu(1)}) \frac{\partial}{A_1 \partial \xi_{1\mu(1)}} \frac{\partial}{A_2 \partial \xi_{2\mu(2)}} I(\Gamma) \\ &= -\text{Tr}(\gamma_{\mu(3)} \gamma_{\mu(2)} \gamma_{\mu(4)} \gamma_{\mu(1)}) (\xi_1 - \xi_2)_{\mu(1)} (\xi_2 - \xi_1)_{\mu(2)} A_1 A_2 \times \\ &\quad \times \frac{e^{-\frac{(\xi_1 - \xi_2)^2 A_1 A_2}{A_1 + A_2}}}{(A_1 + A_2)^4} dA_1 dA_2 \left(A_\gamma^{F_1} = \frac{A_1 A_2}{(A_1 + A_2)^4}, |A_\gamma^{F_1}|_\gamma = 0 \right) \\ &\quad + \text{Tr}(\gamma_{\mu(3)} \gamma_{\mu(2)} \gamma_{\mu(4)} \gamma_{\mu(1)}) \frac{1}{2} g_{\mu(1)\mu(2)} \times \\ &\quad \times \frac{e^{-\frac{(\xi_1 - \xi_2)^2 A_1 A_2}{A_1 + A_2}}}{(A_1 + A_2)^3} dA_1 dA_2 \left(A_\gamma^{F_2} = \frac{1}{(A_1 + A_2)^3}, |A_\gamma^{F_2}|_\gamma = 2 \right). \end{aligned}$$

Partially integrating the metric tensor term (equivalently, multiplying $A_\gamma^{F_2}$ by $\frac{A_1 A_2}{(A_1 + A_2) A_4}$ before integrating A_4 , see Eq.(5)) gives

$$\begin{aligned} \pi_1 &= \text{Tr}(\gamma_{\mu(3)} \gamma_{\mu(2)} \gamma_{\mu(4)} \gamma_{\mu(1)}) (\xi_1 - \xi_2)_{\mu(1)} (\xi_2 - \xi_1)_{\mu(2)} A_1 A_2 \times \\ &\quad \times \frac{e^{-\frac{(\xi_1 - \xi_2)^2 A_1 A_2}{A_1 + A_2}}}{(A_1 + A_2)^4} dA_1 dA_2 \\ &\quad + \text{Tr}(\gamma_{\mu(3)} \gamma_{\mu(2)} \gamma_{\mu(4)} \gamma_{\mu(1)}) \frac{1}{2} g_{\mu(1)\mu(2)} (\xi_1 - \xi_2)^2 A_1 A_2 \times \\ &\quad \times \frac{e^{-\frac{(\xi_1 - \xi_2)^2 A_1 A_2}{A_1 + A_2}}}{(A_1 + A_2)^4} dA_1 dA_2. \end{aligned}$$

Evaluating the trace, contracting indices and integrating delivers (Q replaces $\xi_1 - \xi_2$ by q , subtraction at $q^2 = \mu^2$ understood, *i.e.* Q_0 replaces $\xi_1 - \xi_2$ by μ)

$$\Pi_1 = 8(q^2 g_{\mu(3)\mu(4)} - q_{\mu(3)} q_{\mu(4)}) \int \frac{A_1 A_2 e^{-\frac{q^2 A_1 A_2}{A_1 + A_2}}}{(A_1 + A_2)^4} \cdots|_{q^2 = \mu^2} dA_1 dA_2$$

which can be written projectively

$$\Pi_1 = 8(q^2 g_{\mu(3)\mu(4)} - q_{\mu(3)} q_{\mu(4)}) \ln \frac{q^2}{\mu^2} \int_{\mathbb{P}^1(\mathbb{R}_+)} \frac{A_1 A_2}{(A_1 + A_2)^4} (A_1 dA_2 - A_2 dA_1)$$

and which correctly evaluates to the expected transversal result

$$\Pi_1 = \frac{4}{3} (q^2 g_{\mu(3)\mu(4)} - q_{\mu(3)} q_{\mu(4)}) \ln \frac{q^2}{\mu^2}.$$

Next, we turn to Yang–Mills theory.

Example 6.16. We have

$$\frac{|N_{\text{O}}|_{\text{Pf}}}{\psi_{\text{O}}} = -\xi_3^2 A_3 - \xi_4^2 A_4 - \frac{(\xi_1 - \xi_2)^2 A_1 A_2}{A_1 + A_2}$$

while the corolla polynomials read

$$\begin{aligned} C_{\text{O}}^0(\underline{a}) &= (a_{a3} + a_{a1} + a_{a2})(a_{b4} + a_{b1} + a_{b2}) \\ C_{\text{O}}^1(\underline{a}) &= a_{a3} a_{b4} \end{aligned}$$

The corresponding differentials then become

$$\begin{aligned} C_{\text{O}}^0(\underline{D}) &= (D_{a3} + D_{a1} + D_{a2})(D_{b4} + D_{b2} + D_{b1}) \\ &= \left(-\frac{1}{2}\right)^2 \left(g^{\mu_1 \mu_2} \left(-\frac{1}{A_1} \frac{\partial}{\partial \xi_{1\mu_1}} - \frac{1}{A_2} \frac{\partial}{\partial \xi_{2\mu_1}} \right) \right. \\ &\quad + g^{\mu_2 \mu_1} \left(\frac{1}{A_2} \frac{\partial}{\partial \xi_{2\mu_1}} - \frac{1}{A_3} \frac{\partial}{\partial \xi_{3\mu_1}} \right) \\ &\quad \left. + g^{\mu_1 \mu_1} \left(\frac{1}{A_3} \frac{\partial}{\partial \xi_{3\mu_2}} + \frac{1}{A_1} \frac{\partial}{\partial \xi_{1\mu_2}} \right) \right) \\ &\quad \times \left(g^{\mu_2 \mu_1} \left(-\frac{1}{A_2} \frac{\partial}{\partial \xi_{2\mu_2}} - \frac{1}{A_1} \frac{\partial}{\partial \xi_{1\mu_2}} \right) \right. \\ &\quad + g^{\mu_1 \mu_2} \left(\frac{1}{A_1} \frac{\partial}{\partial \xi_{1\mu_2}} - \frac{1}{A_4} \frac{\partial}{\partial \xi_{4\mu_2}} \right) \\ &\quad \left. + g^{\mu_2 \mu_2} \left(\frac{1}{A_4} \frac{\partial}{\partial \xi_{4\mu_1}} + \frac{1}{A_2} \frac{\partial}{\partial \xi_{2\mu_1}} \right) \right) \\ C_{\text{O}}^1(\underline{D}) &= D_{a3} D_{b4} \\ &= 4 \left(-\frac{1}{2}\right)^2 \left(-\frac{1}{A_1} \frac{\partial}{\partial \xi_{1\mu_1}} - \frac{1}{A_2} \frac{\partial}{\partial \xi_{2\mu_1}} \right) \\ &\quad \times \left(-\frac{1}{A_2} \frac{\partial}{\partial \xi_{2\mu_2}} - \frac{1}{A_1} \frac{\partial}{\partial \xi_{1\mu_2}} \right) \end{aligned}$$

for which the linear part, without the factor 4 (the space-time dimension), is

$$\tilde{C}_{\text{O}}^1(\underline{D}) = \left(-\frac{1}{2}\right)^2 \frac{1}{A_1 A_2} \left(\frac{\partial^2}{\partial \xi_{1\mu_1} \partial \xi_{2\mu_2}} + \frac{\partial^2}{\partial \xi_{2\mu_1} \partial \xi_{1\mu_2}} \right)$$

We compute

$$\begin{aligned} U_{\text{O}}^0 &= C_{\text{O}}^0(\underline{D}) \frac{e^{\bar{\phi}_{\text{O}}/\psi_{\text{O}}}}{\psi_{\text{O}}^2} \\ &= \frac{1}{(A_1 + A_2)^4} \left((A_1 - A_2) g^{\mu_1 \mu_2} q^{\mu_1} - (2A_1 + A_2) g^{\mu_2 \mu_1} q^{\mu_1} \right. \\ &\quad \left. + (A_1 + 2A_2) g^{\mu_1 \mu_1} q^{\mu_2} \right) \left((A_1 - A_2) g^{\mu_2 \mu_1} q^{\mu_2} \right. \\ &\quad \left. + (A_1 + 2A_2) g^{\mu_1 \mu_2} q^{\mu_2} - (2A_1 + A_2) g^{\mu_2 \mu_2} q^{\mu_1} \right) \\ &\quad \times e^{-q^2 \left(\frac{A_1 A_2}{A_1 + A_2} + A_3 + A_4 \right)} \\ &\quad \rightarrow 0, \text{ as residues are scale-independent self-loops} \\ &\quad + \frac{3}{(A_1 + A_2)^3} \left(1 - \overbrace{\frac{A_1}{A_2} - \frac{A_2}{A_1}} \right) g^{\mu_1 \mu_2} e^{-q^2 \left(\frac{A_1 A_2}{A_1 + A_2} + A_3 + A_4 \right)} \end{aligned}$$

and so

$$\begin{aligned} \tilde{U}_{-\circlearrowleft}^0 &= \left(\frac{1}{(A_1 + A_2)^4} \left(- (2A_1^2 + 2A_2^2 + 14A_1A_2)q^{\mu_1}q^{\mu_2} \right. \right. \\ &\quad \left. \left. + (5A_1^2 + 5A_2^2 + 8A_1A_2)q^2g^{\mu_1\mu_2} \right) \right)_{\text{hence } |A_\gamma^{F_1}|_\gamma=0} e^{-q^2 \frac{A_1A_2}{A_1+A_2}} dA_1 dA_2 \\ &\quad + \left(\frac{3}{(A_1 + A_2)^3} g^{\mu_1\mu_2} \right)_{\text{hence } |A_\gamma^{F_2}|_\gamma=2} e^{-q^2 \frac{A_1A_2}{A_1+A_2}} dA_1 dA_2 \end{aligned}$$

Similarly,

$$\tilde{U}_{-\circlearrowleft}^1 = \left(\frac{2A_1A_2}{(A_1 + A_2)^4} q^{\mu_1}q^{\mu_2} - \frac{1}{(A_1 + A_2)^3} g^{\mu_1\mu_2} \right) e^{-q^2 \frac{A_1A_2}{A_1+A_2}} dA_1 dA_2$$

We thus obtain for the corresponding integrals:

$$\int \tilde{U}_{-\circlearrowleft}^{0R} = \left(\frac{11}{3} q^{\mu_1}q^{\mu_2} - \frac{25}{6} q^2 g^{\mu_1\mu_2} \right) \ln \left(\frac{q^2}{\mu^2} \right),$$

which corresponds to the gauge boson loop:



and

$$(24) \quad \int \tilde{U}_{-\circlearrowleft}^{1R} = - \left(\frac{1}{3} q^{\mu_1}q^{\mu_2} + \frac{1}{6} q^2 g^{\mu_1\mu_2} \right) \ln \left(\frac{q^2}{\mu^2} \right),$$

which corresponds to the ghost loop:



They combine to a transversal result:

$$\begin{aligned} \int \tilde{U}_{-\circlearrowleft}^R &= \int \tilde{U}_{-\circlearrowleft}^{0R} - \int \tilde{U}_{-\circlearrowleft}^{1R} \\ &= 4(q^{\mu_1}q^{\mu_2} - q^2g^{\mu_1\mu_2}) \ln \left(\frac{q^2}{\mu^2} \right). \end{aligned}$$

Multiplying with $\frac{\text{colour}(\gamma)}{\text{sym}(\gamma)} = \frac{1}{2} f^{h_1h_2h_3} f^{h_2h_3h_6}$, this is the result for the 1-loop gluon self-energy in Yang–Mills theory. The gauge theory result is immediate from including the previous example with a suitable colour factor for the fermion loop.

7. CONCLUSION

7.1. Covariant quantization without ghosts. Consider $\overline{U}_\Gamma^R =: \sum_{i=0}^{\infty} (-1)^i \overline{U}_\Gamma^{i,R}$ in a notation which reflects the alternating structure of the corolla polynomial. Set $\overline{U}_{\text{gh}}^R := \sum_{i=1}^{\infty} (-1)^i \overline{U}_\Gamma^{i,R}$.

Covariant quantization delivers naively the integrand $\overline{U}_\Gamma^{0,R}$. Let P_L be a projector onto longitudinal degrees of freedoms so that a physical amplitude is in the kernel of P_L , P_T the corresponding projector such that $P_L + P_T = \text{id}$.

Summing over connected graphs contributing to a physical amplitude $X^{r,n}$ at n loops, we know that

$$P_L \left(\overline{U}_{X^{r,n}}^{0,R} \right) = -P_L \left(\overline{U}_{X^{r,n}}^{\text{gh},R} \right).$$

The undesired longitudinal part of the ghost free sector determines the longitudinal part of the ghost contribution by definition.

But also, to compute the ratio

$$\frac{P_L \left(\overline{U}_{X^{r,n}}^{\text{gh},R} \right)}{P_T \left(\overline{U}_{X^{r,n}}^R \right)}$$

is a combinatorial exercise in determining the interplay of these projectors with the Leibniz terms originating from the corolla differentials in the various topologies. These longitudinal and transversal differentials are determined by the same scalar integrand, and hence not independent. Eq.(24) with the ratio two between the qq and g form-factor is a typical example.

So the transversal part of the ghost sector is determined by the combinatorics of scalar graphs and the longitudinal part. It hence is implicitly determined by the ghost free sector.

7.2. Slavnov–Taylor Identities. Slavnov–Taylor identities are treated here as originating from co-ideals in the corresponding Hopf algebras. We reproduce the Feynman rules in four dimensions as renormalized integrands, and can similarly reproduce them in dimensional regularization, and checked that our renormalized Feynman integrand vanishes on the corresponding co-ideals, as it should.

In future work, we will directly demonstrate the validity of Slavnov–Taylor identities from the structure of the corolla polynomial.

REFERENCES

- [1] S. Bloch, H. Esnault and D. Kreimer, *On Motives associated to graph polynomials*, Commun. Math. Phys. **267** (2006) 181 [math/0510011 [math-ag]].
- [2] F. Brown, *The Massless higher-loop two-point function*, Commun. Math. Phys. **287** (2009) 925 [arXiv:0804.1660 [math.AG]].
- [3] F. C. S. Brown, *On the periods of some Feynman integrals*, arXiv:0910.0114 [math.AG].
- [4] F. Brown and O. Schnetz, *A K3 in ϕ^4* , Duke Math. J. Volume **161**, Number 10 (2012), 1817-1862; arXiv:1006.4064 [math.AG].
- [5] F. Brown and O. Schnetz, *Proof of the zig-zag conjecture*, arXiv:1208.1890 [math.NT].
- [6] F. Brown, O. Schnetz and K. Yeats, *Properties of c_2 invariants of Feynman graphs*, arXiv:1203.0188 [math.AG].
- [7] D. Broadhurst, see <http://www.mathematik.hu-berlin.de/~maphy/QPPIIBroadhurst.pdf> and *Feynman’s sunshine numbers*, arXiv:1004.4238 [physics.pop-ph].
- [8] Z. Bern, J. J. M. Carrasco, L. J. Dixon, H. Johansson and R. Roiban, *Simplifying Multiloop Integrands and Ultraviolet Divergences of Gauge Theory and Gravity Amplitudes*, Phys. Rev. D **85** (2012) 105014 [arXiv:1201.5366 [hep-th]].
- [9] D. A. Kosower, R. Roiban and C. Vergu, *The Six-Point NMHV amplitude in Maximally Supersymmetric Yang-Mills Theory*, Phys. Rev. D **83** (2011) 065018 [arXiv:1009.1376 [hep-th]].
- [10] D. Kreimer and K. Yeats, *Properties of the corolla polynomial of a 3-regular graph*, arXiv:1207.5460 [math.CO].
- [11] S. Weinberg, *Feynman Rules for Any Spin*, Phys. Rev. **133** (1964) B1318.
- [12] S. Weinberg, *Feynman Rules for Any Spin. 2. Massless Particles*, Phys. Rev. **134** (1964) B882.
- [13] S. Weinberg, *Feynman rules for any spin. iii*, Phys. Rev. **181** (1969) 1893.
- [14] P. Cvitanovic, *Field Theory*, ChaosBook.org/FieldTheory, Niels Bohr Institute (Copenhagen 2004), RX-1012 (NORDITA).
- [15] J. Conant, K. Vogtmann, *On a theorem of Kontsevich*, Algebr. Geom. Topol. **3** (2003) 1167-1224, arXiv:math/0208169v2.
- [16] D. Kreimer, *Algebraic Structures in local QFT*, Nucl. Phys. Proc. Suppl. **205-206** (2010) 122 [arXiv:1007.0341 [hep-th]], and references here.
- [17] F. Brown and D. Kreimer, *Angles, Scales and Parametric Renormalization*, arXiv:1112.1180 [hep-th].
- [18] D. Kreimer and E. Panzer, *Renormalization and Mellin transforms*, arXiv:1207.6321 [hep-th].
- [19] S. Bloch and D. Kreimer, *Feynman amplitudes and Landau singularities for 1-loop graphs*, Commun. Num. Theor. Phys. **4** (2010) 709 [arXiv:1007.0338 [hep-th]].
- [20] W. D. van Suijlekom, *The structure of renormalization Hopf algebras for gauge theories I: Representing Feynman graphs on BV-algebras*, Commun. Math. Phys. **290** (2009) 291-319. arXiv:0807.0999 [math-ph]
- [21] W. D. van Suijlekom, *Renormalization of gauge fields: A Hopf algebra approach*, Commun. Math. Phys. **276** (2007) 773 [hep-th/0610137].
- [22] D. Kreimer, *Anatomy of a gauge theory*, Annals Phys. **321** (2006) 2757 [hep-th/0509135].
- [23] P. Aluffi, M. Marcolli, *Feynman motives and deletion-contraction relations*, arXiv:0907.3225v1 [math-ph].

HUMBOLDT UNIVERSITY, 10099 BERLIN, GERMANY (D.K., M.S.) AND RADBOUD UNIVERSITY NIJMEGEN, 6525 AJ NIJMEGEN, THE NETHERLANDS