

ANGLES, SCALES AND PARAMETRIC RENORMALIZATION

FRANCIS BROWN* AND DIRK KREIMER**

1. INTRODUCTION: THE DECOMPOSITION OF $\text{Spec}_{\text{Feyn}}(H)$

In this paper we analyse periods appearing in the renormalization of divergent Feynman graphs Γ . Throughout, for any graph Γ , we let $\Gamma^{[1]}$ be the set of internal edges of Γ , and $\Gamma^{[0]}$ be the set of vertices of Γ .

For each vertex $v \in \Gamma^{[0]}$, we assign a four-momentum $q(v)$; to each edge $e \in \Gamma^{[1]}$, we assign a mass m_e .

Our study holds for quantum field theories quite generally, but we specialise in the following to the case of a scalar field theory with four-valent vertices in four dimensions of space-time, for concreteness.

Amplitudes in quantum field theory can be written as a function of a chosen energy variable $L = \ln(S/S_0)$. Here, S sets the scale of the process under consideration. We take S to be a suitable linear combination of scalar products $q(v) \cdot q(w)$ of external momenta and squared masses m_e^2 such that $0 < S \in \mathbb{R}$. Dimensionless scattering angles Θ like $q(v) \cdot q(w)/S$ and m_e^2/S are defined with respect to the choice of S . Throughout, we will assume that S is chosen such that it only vanishes when all external momenta vanish. In these variables, amplitudes can be calculated as a perturbation expansion in terms of Feynman graphs Γ as $\sum_{\Gamma} \Phi^R(\Gamma)$. Here, the renormalized Feynman rules Φ^R are expressed in terms of such angle and scale variables, and the graphs Γ form a Hopf algebra H . For any choice of angle and scale variables, Φ^R is in the group $\text{Spec}(H)_{\mathbb{C}}$, and the restriction of this group to maps which originate from evaluation of graphs by Feynman rules defines a sub-group $G = \text{Spec}_{\text{Feyn}}(H) \subset \text{Spec}(H)_{\mathbb{C}}$.

Such a chosen decomposition of the variables reflects itself then in a chosen decomposition of the group G into two subgroups G_{1-s} and G_{fin} . Elements $\Phi \in G_{1-s}$ are of the form

$$(1) \quad \Phi(\Gamma) = \sum_{j=1}^{\text{cor}(\Gamma)} p_j L^j,$$

where the coefficients p_j are periods in the sense of algebraic geometry and are independent of the angles $\{\Theta\}$. The integer $\text{cor}(\Gamma)$ is defined below, it basically counts how often divergent sub-graphs are nested in a graph.

We allow for renormalization conditions which are defined by kinematic constraints on Green-functions: we demand that such Green functions, regarded as functions of S and $\{\Theta\}$, vanish (up to a specified order) at a reference point (in $S, \{\Theta\}$ -space) given by $S_0, \{\Theta_0\}$. We implement these constraints graph by graph.

*Supported by CNRS and ERC grant 257638. **Alexander von Humboldt Chair in Mathematical Physics, supported by the Alexander von Humboldt Foundation and the BMBF.

Hence renormalized Green functions as well as renormalized Feynman rules become functions of S, S_0, Θ, Θ_0 . Here, Θ, Θ_0 stand for the whole set of angles in the Feynman rules.

Elements $\Phi_{\text{fin}} \in G_{\text{fin}}$ are of the form

$$(2) \quad \Phi_{\text{fin}}(\Gamma) = c_0^\Gamma(\Theta),$$

with $c_0^\Gamma(\Theta)$ an L -independent function of the angles.

We hence obtain the decomposition of G as a map $\Phi^R \rightarrow (\Phi_{\text{fin}}, \Phi_{1-s}^R)$, which proceeds then by a twisted conjugation:

$$(3) \quad G \ni \Phi^R(S, S_0, \Theta, \Theta_0) = \Phi_{\text{fin}}^{-1}(\Theta_0) \star \Phi_{1-s}^R(S, S_0) \star \Phi_{\text{fin}}(\Theta),$$

with $\Phi_{\text{fin}}(\Theta_0), \Phi_{\text{fin}}(\Theta) \in G_{\text{fin}}$ and $\Phi_{1-s}^R(S, S_0) \in G_{1-s}$. The group law \star and inversion $^{-1}$ are defined through the Hopf algebra underlying G .

In particular, an overall divergent Feynman graph with divergent sub-graphs will evaluate to

$$(4) \quad \Phi^R(\Gamma) = \sum_{j=0}^{\text{cor}(\Gamma)} c_j^\Gamma(\Theta, \Theta_0) L^j,$$

while for an overall convergent graph

$$(5) \quad \Phi^R(\Gamma) = \sum_{j=0}^{\text{cor}(\Gamma)-1} c_j^\Gamma(\Theta, \Theta_0) L^j,$$

and the product structure manifests itself in particular in the facts that in Eq.(??), $c_{\text{cor}(\Gamma)}^\Gamma$ is in fact an angle-independent coefficient, while Eq.(??) is non-leading in its L expansion, (it stops at $c_{\text{cor}(\Gamma)}^\Gamma - 1$) with angle dependent coefficients throughout.

We proceed by first reviewing forests in Feynman graphs, which are the appropriate tool to label divergent sectors. They are most efficiently organized using the Hopf algebra structure of rooted trees. Renormalized Feynman rules for a given Feynman graph Γ evaluate the graph using a sum over all forests in the graph, evaluating divergent sub-graphs to renormalized values in each forest using suitable renormalization conditions. In a final step the remaining overall divergence is subtracted using renormalization conditions on the whole graph.

Each forest corresponds to a sector in the initial Feynman integral which has a localized singularity along the corresponding sub-graphs. Signs in the sum over all forests are arranged such that in the total expression, the Feynman integral allows for integration in all sectors. The underlying combinatorics corresponds then to a Hopf algebra.

We next introduce Feynman rules in parametric space. We put emphasis on giving them as integrands for projective spaces. We discuss in detail subtractions for quadratic sub-divergences in accordance with BPHZ renormalization conditions for the propagator (the 1PI self-energy is supposed to vanish to first order at $q^2 = m^2$, with m^2 being the physical mass).

The resulting unrenormalized amplitudes, after suitable partial integrations, have strictly logarithmic singularities, overall as well as in proper sub-graphs.

Inclusion-exclusion is then the mechanism which eliminates these logarithmic singularities by suitable subtractions along any singular sector, making use of the combinatorial properties of the underlying Hopf algebra structure.

Logarithmic singularities are angle-independent. The same combinatorial inclusion-exclusion properties alluded to above allow us then to compensate singularities by using a different evaluation of graphs in which angle dependences have been eliminated, by choosing enough masses m_e and momenta $q(v)$ to be zero. In particular, we choose an element $\bar{\Phi} \in G_{1-s}$ which maps a Feynman graph to the desired L -polynomial

$$(6) \quad \bar{\Phi}(\Gamma) = \sum_{j=1}^{\text{cor}(\Gamma)} p_j L^j.$$

We define graphs $\bar{\Gamma}$ such that

$$(7) \quad \bar{\Phi}(\Gamma) = \Phi(\bar{\Gamma}).$$

Such graphs (dubbed 1-scale graphs) form a Hopf algebra H_{1-s} again. It separates the periods completely from the angle-dependence of Feynman amplitudes on the level of graphs.

We are then ready to decompose

$$(8) \quad \Phi^R(\Gamma) = \sum_{j=0}^{\text{cor}(\Gamma)} c_j(\Theta) L^j,$$

as announced above. This corresponds to defining a matrix $M_{ij}(\Gamma)$, where i, j run over all forests of a graph Γ . While this matrix defined below plays no prominent role in the conceptual developments discussed here, it will figure prominently in the transition of our approach to concrete algorithms.

For the Hopf algebra of 1-scale graphs, we give renormalization proofs and a derivation of the renormalization group from scratch, using mere combinatorial properties of graph polynomials and Hopf algebras and not allowing ourselves any further analytic input.

The renormalized amplitudes of single-scale graphs are given by explicit integrals involving graph polynomials, and are motivic, i.e., periods of mixed Hodge structures of certain diagrams of hyper-surfaces. In particular, this leads to a rigorous definition of the weight of renormalized amplitudes, and opens the possibility of a qualitative study of renormalized amplitudes from the point of view of algebraic geometry.

Summarizing, we provide an approach which delivers a renormalized Feynman graph as a well-defined sum of projective integrals in parametric space in a form suitable to an algorithmic determination of its periodic content, on combining our results with the studies started in [?, ?].

2. FORESTS

It is useful to collect some notation first.

2.1. Definitions. For a 1PI superficially divergent graph Γ , we define a forest f to be a collection of 1PI proper superficially divergent sub-graphs $\Gamma_i \subset \Gamma$, $i \in \mathcal{I}_\Gamma^f$ for some index set \mathcal{I}_Γ^f , such that either they are disjoint: $\Gamma_i \cap \Gamma_j = \emptyset$, or contained in each other: $\Gamma_i \subset \Gamma_j$ or $\Gamma_j \subset \Gamma_i$. In particular, a forest f is a product of 1PI graphs: $f = \prod_i \gamma_i$. By Γ/f we denote the graph obtained by contracting the graphs γ_i to points in Γ .

For a 1PI superficially divergent graph Γ , we define a maximal forest to be a forest and furthermore, we demand that $p_\Gamma^f := \Gamma / [\cup_{i \in \mathcal{I}_\Gamma^f} \Gamma_i]$ has no divergent sub-graph. We hence call the index set \mathcal{I}_Γ^f maximal for Γ .

For $f \ni \Gamma_i \subset \Gamma$, each index set \mathcal{I}_Γ^f defines an index set \mathcal{I}_i^f of all forests strictly contained in Γ_i , i.e. such that $\Gamma_j \subset \Gamma_i \forall j \in \mathcal{I}_i^f$.

We call a forest complete, if \mathcal{I}_Γ^f is maximal for Γ and \mathcal{I}_i^f maximal for each proper 1PI superficially divergent sub-graph Γ_i of Γ .

Each finite graph Γ has a finite number $|C(\Gamma)|$ of complete forests. Here, the set of all such complete forests is denoted by $C(\Gamma)$. Examples are below.

Such complete forests are in one-to-one correspondence with decorated rooted trees where the set of decorations p_v (at vertices v) is given by 1PI superficially divergent graphs free of sub-divergences,

$$(9) \quad p_v := \Gamma_i / \cup_{j \in \mathcal{I}_i^f} \Gamma_j.$$

From now on, we write in obvious abuse of notation $T \in C(\Gamma)$ for such a decorated rooted tree.

Note that the power set $P_E(T)$ of edges $E(T)$ of such a tree T gives all possible cuts c at the tree T : any $c \in P_E(T)$ defines, for a connected tree T , a union of connected components $T - c$ obtained by removing the edges c , with $R^c(T)$ the unique component containing the root of T , and $P^c(T)$ the union of the remaining components. We have $\cup_{T \in C(\Gamma)} 2^{E(T)}$ as the set of all cuts available altogether, and denote by (c, T) an element of this set.

$P^c(T)$ corresponds to a forest of Γ , with each of its connected components corresponding to a graph γ_i in $f = \prod_i \gamma_i$ ¹.

After having determined the set $C(\Gamma)$, all (non-empty) forests of Γ are in bijection with (non-empty) sets f_c of some cuts (c, T) . We describe them as follows.

If we let $|T|$ be the number of vertices of a tree T , a tree T allows for $2^{|T|}$ cuts including the empty one. For a graph Γ , this gives us $\sum_{T \in C(\Gamma)} 2^{|T|}$ cuts c . By construction, a forest $f = \prod_i \gamma_i$ of a graph Γ assigns to a graph the product $(\Gamma/f) \prod_i \gamma_i$. We have $|C(\Gamma/f)| \prod_i |C(\gamma_i)|$ cuts c_i corresponding to the same forest, and let f_c be the set of cuts (c, T) which correspond to the same forest f .

We often notate a cut (c, T) using T with marked edges, and notate the union f_c then as a sum of such trees. We have $\sum_{T \in C(\Gamma)} 2^{|T|} = \sum_f |f_c|$ by construction².

We can hence label the forests of a graph Γ by subsets of edges on some of the trees $T \in C(\Gamma)$:


$$(10) \quad \sum_f = \sum_{T \in C(\Gamma)} \sum_{c \in f_c} = \sum_{T \in C(\Gamma)} \sum_{c \in P_E(T)}.$$

Furthermore, we identify the empty forest (of Γ) with Γ and write \sum_f^\emptyset when we include it in the sum. If we allow forests also to contain Γ itself, we double the sum of forests and write $\sum_{[f]}$ for the corresponding sum.


¹ $c \rightarrow P^c(T)$ furnishes a surjective map F from $\cup_{T \in C(\Gamma)} 2^{E(T)}$ to the forests f of Γ . The set of pre-images $f_c = F^{-1}(f)$ gives a partition of $\cup_{T \in C(\Gamma)} 2^{E(T)}$ which is a bijection with the forests of Γ .

²The cardinality $|f_c|$ of f_c gives the number of sectors in f and Γ/f .


Example 1. Consider the graph



It has subgraphs



and



We have $\gamma_{34} \subset \gamma_l$ and $\gamma_{34} \subset \gamma_r$. Its forests are

(12) $f_0 = \emptyset, |f_{0c}| = 2,$

(13) $f_1 = \gamma_{34}, |f_{1c}| = 2,$

(14) $f_2 = \gamma_l, |f_{2c}| = 1,$

(15) $f_3 = \gamma_r, |f_{3c}| = 1,$


(16) $f_4 = \gamma_{34}, \gamma_l, |f_{4c}| = 1,$

(17) $f_5 = \gamma_{34}, \gamma_r, |f_{5c}| = 1.$


The forest f_1 is neither maximal nor complete. The forests f_2 and f_3 are both maximal, but incomplete. The forests f_4, f_5 are both complete. Hence, $C(\Gamma) = \{f_4, f_5\}$ is a two-element set.

If we add the graph Γ itself to the forests, we double the set, for each f_i , we now have f_i and $f_i \cup \Gamma$.

The decorated trees T_4, T_5 are complete forests. They are given as:



and

(19) 

We can find the decorations by shrinking all graphs in the subforests of a given forest: we assign to the two maximal complete forests two rooted trees, the root corresponding to the vertex at the outermost box ³.

2.2. Hopf structures. We summarize the relevant Hopf algebra structures [?] as follows.

2.2.1. For trees. For the free commutative algebra of decorated rooted trees H_{Dec} (typically, decorations are provided by either the graphs p_v or their set of edge labels) we have a co-product Δ_T defined by

$$(20) \quad \Delta_T \circ B_+^p(\cdot) = B_+^p(\cdot) \otimes 1 + (\text{id} \otimes B_+^p)\Delta_T,$$

and an antipode given by

$$(21) \quad S(T) = -T - \sum_{c \in P_E(T)} (-1)^{|c|} P^c(T) R^c(T),$$

where $R^c(T)$ contains the root with decoration p and $P^c(T)$ are the other trees in $T - c$. B_+^p are Hochschild 1-cocycles, see [?] for details.

We let $\text{shad} : H_{\text{Dec}} \rightarrow H_\emptyset$ be the map which forgets decorations.

2.2.2. For graphs. For graphs we have a Hopf algebra of graphs H_Γ with co-product

$$(22) \quad \Delta_G(\Gamma) = \Gamma \otimes 1 + 1 \otimes \Gamma + \sum_{\gamma \subset \Gamma} \gamma \otimes \Gamma/\gamma,$$

where γ is a disjoint union $\gamma = \cup_i \gamma_i$ of 1PI graphs which are superficially divergent.

The antipode is given by

$$(23) \quad S(\Gamma) = -\Gamma - \sum_f (-1)^{|f|} \gamma_f \otimes \Gamma/\gamma_f.$$

³Also, we can describe those trees as $T_4 := (((3, 4), 1, 2), 5, 6)$ and $T_5 := (((3, 4), 5, 6), 1, 2)$, where we indicate the tree structure by bracket configurations and decorations by the edge labels of the corresponding primitive graphs. If we notate forests in trees by square brackets [...] corresponding to cuts, then the correspondences are as follows: $f_0 \leftrightarrow (((3, 4), 1, 2), 5, 6) + (((3, 4), 5, 6), 1, 2)$, $f_1 \leftrightarrow ([[3, 4], 1, 2], 5, 6) + ([[3, 4], 5, 6], 1, 2)$, $f_2 \leftrightarrow ([[3, 4], 1, 2], 5, 6)$, $f_3 \leftrightarrow ([[3, 4], 5, 6], 1, 2)$, $f_4 \leftrightarrow ([[3, 4], 1, 2], 5, 6)$, $f_5 \leftrightarrow ([[3, 4], 5, 6], 1, 2)$. The forests corresponding to $f_i \cup \Gamma$ are then notated by replacing the outermost (...) pair of brackets by [...]

We have a Hopf algebra homomorphism $\rho : H_\Gamma \rightarrow H_{\text{Dec}}$ given by $\rho(\Gamma) = \sum_{T \in C(\Gamma)} T$ and with

$$(24) \quad [\rho \otimes \rho] \Delta_G = \Delta_T \rho.$$

For any Hopf algebra $H \in (H_{\text{Dec}}, H_\Gamma)$, we let P be the projection into the augmentation ideal. We set $\sigma := S \star P \equiv m_H(S \otimes P) \Delta$, which vanishes on scalars $\mathbb{Q}\mathbb{I}$. For the Hopf algebra of graphs, one has $\sigma(\Gamma) = \sum_f^\emptyset (-1)^{|f|} f(\Gamma/f)$. We need a well-known lemma:

Lemma 2. *Let id_{Aug} be the identity map $\text{Aug} \rightarrow \text{Aug}$ in the augmentation ideal. We have*

$$(25) \quad \text{id}_{\text{Aug}} = \sum_{j=1}^{\infty} \sigma^{*j} =: \sum_{j=1}^{\infty} \sigma_j.$$

Note that the sums terminate when applied to any element of finite degree in the Hopf algebra. Hopf algebras H allow for a co-radical filtration

$$(26) \quad \mathbb{Q}\mathbb{I} = H^{(0)} \subset H^{(1)} \dots \subset H^{(n)} \subset \dots \subset H.$$

The maps σ_j vanish on elements in the Hopf algebra which are in $H^{(k)}$, $k < j$, and the coradical filtration is defined by the kernels of σ_j : elements in $H^{(k)}$ vanish when acted upon by $\sigma_j, \forall j > k$.

Now any map σ_j above corresponds to a finite sum over forests $\bar{\sigma}_j$. As the empty forest corresponds to the identity map of a graph Γ , we can write for forests

$$(27) \quad \emptyset = \sum_{j=1}^{\infty} \bar{\sigma}_j.$$

The following gives an example for the maps $\bar{\sigma}_j$, acting on the graph Γ of Example ??.

Example 3.

$$(28) \quad \begin{aligned} \bar{\sigma}_1 : & \quad \text{Diagram 1} - \text{Diagram 2} - \text{Diagram 3} \\ & - \text{Diagram 4} + \text{Diagram 5} + \text{Diagram 6}, \end{aligned}$$

$$(29) \quad \begin{aligned} \bar{\sigma}_2 : & \quad \text{Diagram 7} + \text{Diagram 8} + \text{Diagram 9} \\ & - 2 \cdot \text{Diagram 10} - 2 \cdot \text{Diagram 11}, \end{aligned}$$

$$(30) \quad \bar{\sigma}_3 : \quad \text{Diagram 12} + \text{Diagram 13}.$$

Note the multiplicity two generated in two terms in $\bar{\sigma}_2 = \bar{\sigma}_1 \star \bar{\sigma}_1$ in line (??), coming from the fact that the subgraphs γ_2, γ_3 and the cograph Γ/γ_1 are acted upon by $\bar{\sigma}_1$ with the same results.

2.3. (Co-)ideals. For any decorated rooted trees T_1, T_2 we say that T_1 is equivalent to T_2 , $T_1 \sim T_2$, if the trees agree as undecorated trees, $\text{shad}(T_1) = \text{shad}(T_2)$, and at each vertex v , the decorations are by primitive graphs $p_1(v), p_2(v)$ which differ only by the choice of external momenta $q(v)$ at vertices of p_i and masses m_e at internal edges of p_i . We say that two graphs are equivalent, $\Gamma_1 \sim \Gamma_2$, if $C(\Gamma_1)$ and $C(\Gamma_2)$ are two sets of trees which are pairwise equivalent, and similar if the Γ_i are sums of graphs. In particular, if $\Gamma = \sum_i \gamma_i$ say, $C(\Gamma) = \cup_i C(\Gamma_i)$.

2.4. Notation for forests. Finally, we denote a graph with forests generated by a particular $T \in C(\Gamma)$ as follows. For a graph with edges labelled $1, \dots, m$ say, for each $T \in C(\Gamma)$ we can label the vertices $v \in T$ by those edges of Γ which correspond to p_v . Note that this loses the information how and by which edges a decoration p_v was connected to p_w , where w is the vertex above (closer to the root than) the vertex v . This ambiguity will soon give us the freedom to define the desired 1-scale structures.

3. DERIVATION OF RENORMALIZED FEYNMAN RULES IN PARAMETRIC SPACE

We turn to the derivation of Feynman rules. In our Hopf algebra H_Γ , we have graphs Γ with labelled edges $e \in \Gamma^{[1]}$. To a graph Γ , we will assign forms Φ_Γ which depend on the edge labels A_e , the squared masses m_e^2 , and the momenta $q(v)$, $v \in \Gamma^{[0]}$. Physicists may wish to consider these external momenta $q(v)$ as external edges, with a splitting as in say $q(v) = q_1 + q_2$ corresponding to two external edges at v , if so desired (for example to achieve homogeneity in the valence of vertices).

We assume that for a product of graphs $\Gamma_1 \Gamma_2$, labels are not repeated. The forms Φ_Γ have the structure $\Phi_\Gamma = f_\Gamma(\{A_e\})\Omega_\Gamma$, with $f_\Gamma(\{A_e\})$ a function of all the edge variables and Ω_Γ a standard form, see below. With unrepeated labels, $\Phi_{\Gamma_1 \Gamma_2} = f_{\Gamma_1} f_{\Gamma_2} \Omega_{\Gamma_1 \cup \Gamma_2}$.

Renormalized Feynman rules make use of the Hopf algebra H_Γ to construct a linear combination of forms Φ_Γ^R such that it can be integrated against positive real projective $\mathbb{P}^{|\Gamma^{[1]}|-1}$ -space. We write $\Phi^R(\Gamma) \in G = \text{Spec}_{\text{Feyn}}(H)$ for the resulting integral.

3.1. Schwinger parametrization and the exponential integral. We first define the two graph polynomials ψ, φ . Both are configuration polynomials [?]. We define them here though using spanning trees and spanning forests. We have (for a connected graph Γ)

$$(31) \quad \psi_\Gamma := \sum_T \prod_{e \notin T} A_e,$$

for spanning trees T and edges e of Γ . Furthermore, we let $q(v)$ be the external momentum entering a vertex $v \in \Gamma$ (it can be zero), and for a subset of vertices $X \subset \Gamma$, we let $Q(X) = \sum_{v \in X} q(v)$. Then,

$$(32) \quad \varphi_\Gamma := \sum_{T_1 \cup T_2} Q(T_1) \cdot Q(T_2) \prod_{e \notin T_1 \cup T_2} A_e,$$

where $T_1 \cup T_2$ is a spanning two-forest. Note: $Q(T_1) = -Q(T_2)$, $Q(T_1)^2 = Q(T_2)^2 = -Q(T_1) \cdot Q(T_2)$. Note that if Γ has only two distinct vertices v_1, v_2 say at which external momenta $q, -q$ enter (we call such a graph a two-point graph) we can write $\varphi_\Gamma = -q^2 \psi_{\Gamma^\bullet}$, where Γ^\bullet is the graph obtained from Γ by identifying the two

vertices v_1, v_2 . We extend these definitions to products of graphs as follows. For $\gamma = \prod_i \gamma_i$,

$$(33) \quad \psi_\gamma = \prod_i \psi_{\gamma_i}, \quad \varphi_\gamma = \sum_i \left(\varphi_{\gamma_i} \prod_{j \neq i} \psi_{\gamma_j} \right).$$

For 1-scale graphs γ , φ_γ becomes the circular join introduced below, see Eq.(??).

Define $Q_{vw} := q(v) \cdot q(w)$, let $S := \sum_{v,w \in \Gamma(0)} c_{vw} Q_{vw}$ a real ($c_{vw} \in \mathbb{R}$) linear combination of scalar products Q_{vw} which vanishes only when all external momenta $q(v)$ vanish. We say that S is in general kinematic position. Let $\Theta_{vw} := Q_{vw}/S$ and $\Theta_e := m_e^2/S$.

$$(34) \quad \varphi_\Gamma(\Theta) := \frac{\varphi_\Gamma}{S}, \quad \phi_\Gamma(S, \Theta) := S\phi_\Gamma(\Theta), \quad \phi_\Gamma(\Theta) := \varphi_\Gamma(\Theta) + \psi_\Gamma \left(\sum_e A_e \Theta_e \right).$$

We usually write $\phi_\Gamma \equiv \phi_\Gamma(S, \Theta)$ in the decomposed form (and in slight abuse of notation) as $\phi_\Gamma = S\phi_\Gamma(\Theta)$. Extension to products is defined as before.

For a two-point graph Γ , we define

$$(35) \quad \varphi_\Gamma = -q^2 \psi_{\Gamma^\bullet}, \quad \phi_\Gamma(\Theta) := -\psi_{\Gamma^\bullet} + \psi_\Gamma \left(\sum_e A_e \frac{m_e^2}{q^2} \right).$$

For such a two-point graph γ , we also define

$$(36) \quad \bar{\psi}_\gamma := \left(-\psi_{\Gamma^\bullet} + \psi_\Gamma \left(\sum_e A_e \frac{m_e^2}{m^2} \right) \right),$$

with m^2 fixing the scale at which a two-point graph is subtracted.

We have for any $\gamma \subset \Gamma$, with $\gamma = \cup_i \gamma_i$, $\psi_\gamma = \prod_i \psi_{\gamma_i}$,

Proposition 4.

$$(37) \quad \psi_\Gamma = \psi_{\Gamma/\gamma} \psi_\gamma + R_\gamma^\Gamma, \quad |R_\gamma^\Gamma|_\gamma = |\psi(\gamma)|_\gamma + 1,$$

$$(38) \quad \phi_\Gamma(\Theta) = \phi_{\Gamma/\gamma}(\Theta) \psi_\gamma + \bar{R}_\gamma^\Gamma(\Theta), \quad |\bar{R}_\gamma^\Gamma(\Theta)|_\gamma \geq |\psi(\gamma)|_\gamma + 1,$$

and $|\phi_\Gamma| = |\psi_\Gamma| + 1$, and $|U|_V$ is the degree of U in the edge variables of V , and $|U| = |U|_U$.

Note that $\phi_{\Gamma/\gamma}(\Theta)$ can be zero, for example when masses are zero and $Q(T_i) = 0$ for all two-forests of Γ/γ .

Proof. From the definitions via spanning trees and two-forests. \square

We now let \square_Γ be the hypercube $\mathbb{R}_+^{|\Gamma^{(1)}|}$, and consider the integrand obtained from a Schwinger parametrization of a Feynman graph Γ ,

$$(39) \quad \Phi_\Gamma(S, \Theta) := \frac{dA_1 \cdots dA_{|\Gamma^{(1)}|} e^{+\frac{S\phi_\Gamma(\Theta)}{\psi_\Gamma}}}{\psi_\Gamma^2}.$$

This unrenormalized integrand cannot be integrated yet in the edge variables A_e against \square_Γ . Its renormalized counterpart has the form (say for logarithmic divergences, the general case is below and has the same structure)

$$(40) \quad \Phi_\Gamma^R(S, S_0, \Theta, \Theta_0) = \sum_{[f]}^{\emptyset} (-1)^{|f|} \Phi_f(S_0, \Theta_0) \Phi_{\Gamma/f}(S, \Theta).$$

$$(41) \quad = \sum \Phi_{\Gamma'}^{-1}(S_0, \Theta_0) \Phi_{\Gamma''}(S, \Theta),$$

where we used Sweedler's notation $\Delta_G(\Gamma) = \sum \Gamma' \otimes \Gamma''$ in the second line. This is the traditional forest formula⁴.

In the following, we will renormalize this integrand using kinetic renormalization schemes. For that, we let $2s_\Gamma \equiv 2sd(\Gamma)$ be the superficial degree of divergence of Γ (in the example of a massive scalar field theory with quartic interactions):

$$(42) \quad 2s_\Gamma = 4|\Gamma| - 2|\Gamma^{[1]}|.$$

Then, all vertex graphs Γ have $s_\Gamma = 0$ together with $|\Gamma^{[1]}| = 2|\Gamma|$, while for all propagator graphs, $s_\Gamma = 1$ with $|\Gamma^{[1]}| = 2|\Gamma| - 1$.

Let us introduce new variables $A_e \rightarrow a_e, A_e = ta_e$, and $dA_1 \cdots dA_{|\Gamma^{[1]}|} \rightarrow dt \wedge \Omega_\Gamma$, with Ω_Γ the usual $(|\Gamma^{[1]}| - 1)$ -form $A_1 dA_2 \wedge \cdots \wedge dA_{|\Gamma^{[1]}|} - A_2 dA_1 \cdots \pm \cdots$, see Eq.(??). We find

$$(43) \quad \Phi_\Gamma := \frac{dt}{t} \wedge \frac{\Omega_\Gamma e^{t \frac{S\phi_\Gamma(\Theta)}{\psi_\Gamma}}}{t^{s_\Gamma} \psi_\Gamma^2}.$$

We want to study the overall t -integration as a function of the superficial degree of divergence s_Γ first. Concretely, we are interested to define and find the limit in the t -integration

$$(44) \quad \lim_{c \rightarrow 0} \int_c^\infty \Phi_\Gamma,$$

where $c \in \mathbb{R}_+$. We use renormalization conditions on $\Phi_\Gamma \equiv \Phi_\Gamma(S, \Theta)$.

Kinetic renormalization conditions imply that we choose values S_0, Θ_0 for the scale and for the angles, such that the renormalized amplitudes of a graph Γ , together with their first s_Γ derivatives in an expansion around that point, vanish. For $s_\Gamma = 0$, we can simply subtract at a chosen S_0, Θ_0 :

$$(45) \quad \Phi_\Gamma(S, \Theta) \rightarrow \Phi_\Gamma(S, \Theta) - \Phi_\Gamma(S_0, \Theta_0)$$

which takes care of the overall divergence in the graph Γ .

For $s_\Gamma = 1$, we are dealing with a quadratically divergent propagator function. We will subtract at $q^2 = m^2$. Note that there are no angles Θ_{vw} for a two-point function, the Θ_e remain though. Kinetic renormalization conditions are determined by the requirement that the renormalized amplitude vanishes at $q^2 = m^2$, together with its first derivative ∂_{q^2} , so that the pole in the propagator has a on-shell unit residue⁵.

⁴Note that we use $\psi_\emptyset = 1, \phi_\emptyset(\Theta) = 0$.

⁵For a massless propagator, vanishing of Φ_Γ^R at $q^2 = 0$ and of Φ_Γ^R/q^2 at $q^2 = \mu^2$ are also convenient renormalization conditions.

For overall convergent Green functions, the limit $c \rightarrow 0$ can be taken without having to impose overall constraints, the introduction of constraints (which correspond to over-subtractions) to impose some favoured kinematics would be possible though in this set-up.

3.2. $s_\Gamma = 0$. Let us start with the case $s_\Gamma = 0$. The limit is

$$(46) \quad \lim_{c \rightarrow 0} \int_c^\infty [\Phi_\Gamma(S, \Theta) - \Phi_\Gamma(S_0, \Theta_0)] = \frac{\Omega_\Gamma \ln \frac{S \phi_\Gamma(\Theta)}{S_0 \phi_\Gamma(\Theta_0)}}{\psi_\Gamma^2},$$

using that for small $c > 0$,

$$(47) \quad \int_c^\infty \frac{e^{-tX} dt}{t} = -\ln c + \ln X + \gamma_E + \mathcal{O}(c).$$

Here, γ_E is the Euler–Mascheroni constant. Note that we can decompose the logarithm as

$$\ln \frac{S \phi_\Gamma(\Theta)}{S_0 \phi_\Gamma(\Theta_0)} = \ln(S/S_0) + \ln(\phi_\Gamma(\Theta)/\phi_\Gamma(\Theta_0)),$$

(we assume $S/S_0 > 0$). We assume also that the angles Θ, Θ_0 are chosen such that we are off Landau singularities. Approaching such singularities means studying the corresponding variation of the logarithm above.

Let us now look at logarithmic sub-divergences. A typical term in the forest formula provides an integrand of the form

$$(48) \quad \frac{e^{+\frac{S \phi_{\Gamma/f}(\Theta)}{\psi_{\Gamma/f}}} e^{+\frac{S_0 \phi_f(\Theta_0)}{\psi_f}}}{\psi_{\Gamma/f}^2} - \frac{e^{+\frac{S_0 \phi_{\Gamma/f}(\Theta_0)}{\psi_{\Gamma/f}}} e^{+\frac{S \phi_f(\Theta)}{\psi_f}}}{\psi_f^2}.$$

Combining each of the two products of exponentials into a single exponential and using the exponential integral as above delivers

$$(49) \quad M_f^\Gamma := \frac{\ln \frac{S \phi_{\Gamma/f}(\Theta) \psi_f + S_0 \phi_f(\Theta_0) \psi_{\Gamma/f}}{S_0 \phi_{\Gamma/f}(\Theta_0) \psi_f + S \phi_f(\Theta) \psi_{\Gamma/f}}}{\psi_{\Gamma/f}^2 \psi_f^2} \Omega_\Gamma.$$

Summing over all forests including the empty one delivers the renormalized integrand as the homogeneous of degree zero form

$$(50) \quad \Phi_\Gamma^R := \sum_f^\emptyset (-1)^{|f|} M_f^\Gamma.$$

Φ_Γ^R is an integrand which can, this is just a rewriting of the forest formula, be integrated against $\mathbb{P}^{|\Gamma^{[1]}|-1}(\mathbb{R}_+)$. An explicit proof from scratch is given below though, after we decomposed Feynman rules suitably.

Remark 5. *If we write $\varphi_\Gamma = S \sum_{v,w} \Theta_{vw} c_\Gamma^{vw}$ for some monomials*

$$c_\Gamma^{vw} = c_\Gamma^{vw}(A_1, \dots, A_{|\Gamma^{[1]}|}),$$

in edge variables A_e , then

$$(51) \quad -\partial_{\Theta_{v,w}} \frac{\phi_{\Gamma/f}(\Theta)}{\psi_{\Gamma/f}} = \frac{1}{\psi_{\Gamma/f}} c_{\Gamma/f}^{vw} (\partial_{m_1^2, \dots, \partial_{m_{|\Gamma^{[1]}|}^2}).$$

As an extra power of $\psi_{\Gamma/f}$ in the denominator is equivalent to a shift in the dimension, we can re-derive in this way the usual recursion and dimension shift relations between master integrals.

Next we treat other overall degrees of divergence still with logarithmic sub-divergences, and will then treat the general case including quadratic sub-divergences.

3.3. $s_\Gamma = 1$. Let us next look at the case $s_\Gamma = 1$. We have $\Phi_\Gamma = \Phi_\Gamma(q^2, m^2)$ and for $s_\Gamma = 1$ we look at the Taylor expansion $\Phi_\Gamma = \Phi_\Gamma(m^2, m^2) + (q^2 - m^2)\partial_{q^2}\Phi_\Gamma(m^2, m^2) + \mathcal{O}((q^2 - m^2)^2)$. Partially integrating Eq.(??), we find

$$(52) \quad \int_c^\infty \Phi_\Gamma = \int_c^\infty \frac{dt}{t} \wedge \frac{\Omega_\Gamma e^{-t \frac{s_\Gamma \phi_\Gamma^S}{\psi_\Gamma}}}{t \psi_\Gamma^2} = \frac{1}{c} \left[\frac{\Omega_\Gamma e^{-c \frac{s_\Gamma \phi_\Gamma^S}{\psi_\Gamma}}}{\psi_\Gamma^2} \right] - \frac{S \phi_\Gamma^S}{\psi_\Gamma} \int_c^\infty \frac{dt}{t} \wedge \frac{\Omega_\Gamma e^{-t \frac{s_\Gamma \phi_\Gamma^S}{\psi_\Gamma}}}{\psi_\Gamma^2}.$$

We set $S = q^2$, so $\Theta_e = m_e^2/q^2$, and $S_0 = m^2, \Theta_{e,0} = m_e^2/m^2$. If all masses are equal, we simply have $\Theta_0 = 1$.

In Eq.(??) we have on the rhs a boundary term (the term $\sim 1/c$ in $[\dots]$ brackets) and a logarithmically divergent integral $\sim \ln c$. Subtracting at $q^2 = m^2$ gives for the boundary term a contribution:

$$(53) \quad - \left[\frac{\Omega_\Gamma(q^2 - m^2)\psi_{\Gamma^\bullet}}{\psi_\Gamma^3} \right]$$

where Γ^\bullet is the graph obtained from identifying the two external vertices of Γ . Note that this boundary term contains no higher terms in $(q^2 - m^2)$ in the limit $c \rightarrow 0$. So in the limit this is an expression linear in $(q^2 - m^2)$ which hence vanishes in kinematic renormalization.

For the logarithmic divergent integral on the rhs of Eq.(??) we find

$$(54) \quad - \frac{S \phi_\Gamma^S}{\psi_\Gamma} \int_c^\infty \frac{dt}{t} \wedge \frac{\Omega_\Gamma e^{-t \frac{s_\Gamma \phi_\Gamma^S}{\psi_\Gamma}}}{\psi_\Gamma^2} + \frac{S_0 \phi_\Gamma^{S_0}}{\psi_\Gamma} \int_c^\infty \frac{dt}{t} \wedge \frac{\Omega_\Gamma e^{-t \frac{s_0 \phi_\Gamma^{S_0}}{\psi_\Gamma}}}{\psi_\Gamma^2}.$$

Subtraction of the term linear in $(q^2 - m^2)$ delivers a renormalized integrand free of overall divergences which reads as :

$$(55) \quad M_\emptyset^\Gamma = \frac{m^2 \bar{\psi}_{\Gamma^\bullet}}{\psi_\Gamma} \frac{[x_\emptyset^\Gamma \ln(1 + x_\emptyset^\Gamma) + \ln(1 + x_\emptyset^\Gamma) - x_\emptyset^\Gamma] \Omega_\Gamma}{\psi_\Gamma^2},$$

where we write $\phi_\Gamma(q^2, m^2)/\phi_\Gamma(m^2, m^2)$ as $1 + x_\emptyset^\Gamma$, with

$$x_\emptyset^\Gamma = \frac{(q^2 - m^2)\psi_{\Gamma^\bullet}}{m^2 \bar{\psi}_{\Gamma^\bullet}}.$$

We remind the reader of Eq.(??) which gives $\bar{\psi}_{\Gamma^\bullet}$.

3.4. $s_\Gamma = -1$. In that overall convergent case, with $1 + x_\emptyset^\Gamma := \frac{S \phi_\Gamma(\Theta)}{S_0 \phi_\Gamma(\Theta_0)}$ similarly, we have

$$(56) \quad M_\emptyset^\Gamma = \left(\frac{S_0 \phi_\Gamma(\Theta_0)}{\psi_\Gamma} \right)^{-1} \frac{\Omega_\Gamma}{\psi_\Gamma^2 (1 + x_\emptyset^\Gamma)}.$$

Similarly for the cases $s_\Gamma < -1$. One finds for $s = -k$:

$$(57) \quad M_\emptyset^\Gamma = \left(\frac{S_0 \phi_\Gamma(\Theta_0)}{\psi_\Gamma} \right)^{-k} \frac{(k-1)! (-1)^{k-1} \Omega_\Gamma}{\psi_\Gamma^2 (1 + x_\emptyset^\Gamma)^k}.$$

3.5. Quadratic sub-divergences. We saw before that a partial integration improved an overall quadratically divergent integrand to an overall logarithmic one, plus a boundary term which captured the overall quadratic divergence. That boundary term was eliminated by the BPHZ renormalization conditions for the mass term, leaving us to treat a logarithmic divergent integrand.

We want similarly to use partial integrations to treat quadratic sub-divergences.

We will treat quadratic sub-divergences according to their partial ordering, which is discussed in detail below, see §???. To start, assume $\gamma \subset \Gamma$ is a quadratically divergent sub-graph of Γ and γ is itself free of quadratic divergent sub-graphs.

We first consider the Feynman integral in momentum space. Label the edges of γ by $1, \dots, 2|\gamma| - 1$. We use a parametric representation for the internal propagators of γ and integrate out the internal loop momenta of γ . We let e_γ, f_γ denote the two edges (connectors) which connect γ to the other vertices of $\Gamma - \gamma$, with $1/P_{e_\gamma}(q), 1/P_{f_\gamma}(q)$ their propagators.

This gives a factor

$$(58) \quad F_\gamma := \frac{e^{-\frac{\phi_\gamma}{\psi_\gamma}} dA_1 \cdots dA_{2|\gamma|-1}}{P_{e_\gamma}(q)P_{f_\gamma}(q)\psi_\gamma^2},$$

in the Feynman integrand.

Isolating a variable $t_\gamma = A_1$ via $A_i = t_\gamma a_i, i = 2, \dots, |\gamma|^{[1]}$ allows us to study the behaviour of the integrand with respect to the overall divergence of the quadratically divergent sub-graph.

$$(59) \quad F_\gamma := \frac{e^{-t_\gamma \frac{\phi_\gamma}{\psi_\gamma}} da_2 \cdots da_{2|\gamma|-1}}{P_{e_\gamma}(q)P_{f_\gamma}(q)\psi_\gamma^2} \wedge \frac{dt_\gamma}{t_\gamma^2}.$$

In obvious abuse of notation, we still write ϕ_γ, ψ_γ after the change of variables.

Partially integrating $\int_{c_\gamma}^\infty F_\gamma$ in t_γ delivers

$$(60) \quad \int_{c_\gamma}^\infty F_\gamma = \int_{c_\gamma}^\infty \frac{e^{-t_\gamma \frac{\phi_\gamma}{\psi_\gamma}} da_2 \cdots da_{2|\gamma|-1}}{P_{e_\gamma}(q)P_{f_\gamma}(q)\psi_\gamma^2} \wedge \frac{dt_\gamma}{t_\gamma^2}$$

$$(61) \quad = \int_{c_\gamma}^\infty \frac{-1}{P_{e_\gamma}(q)P_{f_\gamma}(q)} \underbrace{\frac{\phi_\gamma e^{-t_\gamma \frac{\phi_\gamma}{\psi_\gamma}} da_2 \cdots da_{2|\gamma|-1}}{\psi_\gamma^3}}_X \wedge \frac{dt_\gamma}{t_\gamma}$$

$$(62) \quad + \frac{1}{P_{e_\gamma}(q)P_{f_\gamma}(q)} \underbrace{\left[\frac{e^{-c_\gamma \frac{\phi_\gamma}{\psi_\gamma}} da_2 \cdots da_{2|\gamma|-1}}{\psi_\gamma^2} \frac{1}{c_\gamma} \right]}_Y.$$

We now impose BPHZ conditions for the sub-graph γ . We start with the boundary term $Y \equiv Y(q^2, m^2)$. We have

$$(63) \quad Y(q^2, m^2) = Y(m^2, m^2) + \frac{\overbrace{[\phi_\gamma(q^2, m^2) - \phi_\gamma(m^2, m^2)]}^{=(q^2-m^2)\psi_\gamma \bullet}}{\psi_\gamma^3} da_2 \cdots da_{2|\gamma|-1} + \mathcal{O}(c_\gamma).$$

In BPHZ conditions we subtract the term $Y(m^2, m^2)$ and the term linear in $(q^2 - m^2)$. Hence, in the limit $c_\gamma \rightarrow 0$, the Y term leaves no contribution.

Let us now turn to the $X \equiv X(q^2, m^2)$ term. We use $\phi_\gamma(m^2, m^2) = m^2 \bar{\psi}_{\gamma\bullet}$. Then⁶, using $(q^2 - m^2) = P_{f_\gamma}(q)$ and $\phi_\gamma(q^2, m^2) = (q^2 - m^2)\psi_{\gamma\bullet} + \phi_\gamma(m^2, m^2)$,

$$(64) \quad \frac{1}{P_{e_\gamma}(q)P_{f_\gamma}(q)}X = \frac{X_1}{P_{e_\gamma}} + \frac{m^2 X_2}{P_{e_\gamma}P_{f_\gamma}},$$

with

$$(65) \quad X_1 := \frac{(q^2 - m^2)\psi_{\gamma\bullet} e^{-\frac{\phi_\gamma}{\psi_\gamma}} dA_1 dA_2 \cdots dA_{2|\gamma|-1}}{\psi_\gamma^3}$$

and

$$(66) \quad X_2 := \frac{\bar{\psi}_{\gamma\bullet} e^{-\frac{\phi_\gamma}{\psi_\gamma}} dA_1 dA_2 \cdots dA_{2|\gamma|-1}}{\psi_\gamma^3}.$$

Note that we returned from t_γ, a_i to A_1, A_2, \dots variables. Summarizing, BPHZ conditions render a renormalized $X_R(q^2, m^2)$ as

$$(67) \quad \begin{aligned} X_R(q^2, m^2) &= P_{f_\gamma}(q)[X_1(q^2, m^2) - X_1(m^2, m^2)] \\ &+ \left[X_2(q^2, m^2) - X_2(m^2, m^2) \left(1 - \frac{P_{f_\gamma}(q)\psi_{\gamma\bullet}}{\psi_\gamma} \right) \right]. \end{aligned}$$

The last term in the second line makes sure that we subtract from $X_2(q^2, m^2)$ its value at $q^2 = m^2$ and its first Taylor coefficient in the expansion in $(q^2 - m^2)$.

We can now continue to parametrize the remaining propagators in $\Gamma - \gamma$ treating quadratic subdivergences in the order as dictated by their partial ordering (see below) and integrate out all loop momenta to find the final renormalized parametric representations below, now for the cases of quadratic and logarithmic subdivergences jointly. Summarizing, we will give them for overall degrees of divergence $s_\Gamma \in \{-1, 0, 1\}$, from overall convergence to quadratic divergence.

3.6. The final result. As before, we let $s_\Gamma = \text{sd}(\Gamma)/2$. Define

$$(68) \quad 1 + x_f^\Gamma := \frac{\frac{S}{S_0} \phi_{\Gamma/f}(\Theta)\psi_f + \phi_f(\Theta_0)\psi_{\Gamma/f}}{\phi_{\Gamma/f}(\Theta_0)\psi_f + \phi_f(\Theta_0)\psi_{\Gamma/f}},$$

and the following series (which could be easily generalized to $l_j(x), j \in \mathbb{Z}$):

$$(69) \quad l_{-1}(x) = \frac{1}{(1+x)} = \sum_{k=0}^{\infty} (-1)^k x^k$$

$$(70) \quad l_0(x) = \ln(1+x) = \sum_{k=1}^{\infty} (-1)^{k-1} \frac{x^k}{k}$$

$$(71) \quad l_1(x) = (q^2 - m^2)l_0(x) + \left(m^2 \partial_{m^2} \frac{1}{x} \right) l_0(x)^{[1]},$$

⁶If we renormalize the subgraph γ at $m_\gamma^2 \neq m_{f_\gamma}^2$, we set $(q^2 - m_\gamma^2) = (q^2 - m_{f_\gamma}^2) + \Delta_m^2$, and compensate by a redefinition of $\bar{\psi}_{\gamma\bullet}$.

where we need to explain the notation in Eq.(??) as follows. First, $l_1(x)$ is needed only for two-point functions, in which case any variable $x = x(q^2, m^2)$, and we denote $x' = \partial_{q^2} x(q^2, m^2)$. Then,

$$(72) \quad m^2 \partial_{m^2} \frac{1}{x_f^\Gamma} = m^2 \frac{\bar{\psi}_{\Gamma/f^\bullet}}{\psi_{\Gamma/f^\bullet}}.$$

Second, for any function $l_i(x)$, we denote by $l_i(x)^{[k]}$ the function $l_i(x) - \sum_{j=0}^i c_{i,j} x^j$, where $c_{i,j}$ is the j -th Taylor coefficient in the expansion of l_i around $x = 0$. Note that $l_0(x) = l_0^{[0]}(x)$ and $l_1(x) = l_1^{[1]}(x)$ by construction.

Let n_2^X be the number of (sub)-graphs $\gamma \subseteq X$ with $\text{sd}(\gamma) = 2$. Define

$$(73) \quad \omega_X := \prod_{\gamma \subseteq X, \text{sd}(\gamma)=2} \frac{\psi_{\gamma^\bullet}}{\psi_\gamma}.$$

For an integer $i \in \mathbb{Z}$, we let $\theta : \mathbb{Z} \rightarrow \{0, 1\}$ be the function $\theta(i) = 1, i > 0$, $\theta(i) = 0, i \leq 0$.

For all n_2^Γ quadratic sub-graphs of Γ , choose a set C_2 of $n_2^\Gamma - s_\Gamma \theta(s_\Gamma)$ distinct edges in Γ such that for each γ at least one connector (the two edges in $\Gamma - \gamma$ which connect to γ) of γ is an element of C_2 . A subscript C_2/I denotes suppression of the edge variables in C_2/I in the corresponding expression.

We define $E_j^{C_2/I}$ to be the j -th elementary symmetric polynomial in the $|C_2/I|$ variables $\frac{m_e^2 \bar{\psi}_{\gamma_e^\bullet}}{\psi_{\gamma_e}}$, $e \in C_2/I$. $E_0^{C_2/I} := 1$. Then, with

$$(74) \quad M(x_f^\Gamma) := \sum_{I \subseteq C_2} (-1)^{|I|} \left\{ \omega_{\Gamma/f} \omega_f \Omega_\Gamma \sum_{j=0}^{|C_2/I|} (-1)^j \left(\frac{\phi_{\Gamma/f}(\Theta_0)}{\psi_{\Gamma/f^\bullet}} \right)^{\theta(|I|+j)\theta(s_\Gamma)} \times \right. \\ \times \left. \left(\frac{\text{den}(1+x_f^\Gamma), \text{ see Eq.(??)}}{\phi_{\Gamma/f}(\Theta_0) \psi_f + \phi_f(\Theta_0) \psi_{\Gamma/f}} \right)^{(s_\Gamma - |I| - j)\theta(-s_\Gamma + |I| + j)} \times \right. \\ \left. \times E_j^{C_2/I} \frac{l_{s_\Gamma - |I| - j}^{[s_\Gamma]}(x_f^\Gamma)}{\underbrace{\psi_{\Gamma/f}^2 \psi_f^2}_{-2|\Gamma|}} \right\}_{C_2/I} \prod_{e \in I} m_e^2 \frac{\bar{\psi}_{\gamma_e^\bullet}}{\psi_{\gamma_e}},$$

we have

Theorem 6.

$$(75) \quad \Phi_\Gamma^R = \sum_f^{\emptyset} (-1)^{|f|} \int_{\mathbb{P}^n(\mathbb{R}_+)} M(x_f^\Gamma),$$

converges when integrated against projective $\mathbb{P}^n(\mathbb{R}_+)$ spaces of dimension $n = |\Gamma^{[1]}| - |C_2| + |I|$ for each $\Omega_{\Gamma_{C_2/I}}$.

Proof. The formula is a straightforward but tedious rewriting of the forest formula in terms of projective integrals, using Eqs.(??,??,??), and factorizing out $\omega_{\Gamma/f}, \omega_f$ as this is convenient for the treatment of 1-scale graphs later on. Again, the direct proof is a consequence of the corresponding result for 1-scale graphs given below, and the factorization of Feynman rules to which we turn in the next section. \square

Proposition 7. *Tadpoles vanish in renormalization.*

Proof. $\frac{S}{S_0} \phi_{\Gamma/f}(\Theta) = \phi_{\Gamma/f}(\Theta_0)$ if Γ/f is a tadpole, so $x_f^\Gamma = 0$ and $l_i(0) = 0$, $i = 0, 1$. \square

Remark 8. *Vanishing tadpoles define an obvious ideal and co-ideal.*

3.7. Auxiliary Feynman rules. To next introduce Feynman rules such that the desired factorization

$$(76) \quad \Phi_\Gamma^R = \Phi_{\text{fin}}(\Theta_0)^{-1} \star \Phi_{1-s}^R \star \Phi_{\text{fin}}(\Theta),$$

can be established, we have to enlarge the set of Feynman rules.

For that, we first introduce auxiliary Feynman graphs. Let p be some decoration of $T \in C(\Gamma)$ at a vertex below the root. Let $E_p^{\bar{p}}$ be the set of edges which connect p to the decoration \bar{p} above p . We can assume $|E_p^{\bar{p}}| \geq 2$ since Γ is one-particle irreducible.

Let $V_p^{\bar{p}} \subseteq p^{[0]}$ be the set of vertices of p which are connected to edges $e \in E_p^{\bar{p}}$. Let $V_p^{\text{ext}} \subset \Gamma^{[0]} \cap p^{[0]}$ be external vertices of Γ ($q(v) \neq 0$) which are also vertices of p . Set $V_p := V_p^{\bar{p}} \cup V_p^{\text{ext}}$. Set $q_p := \sum_{v \in V_p^{\text{ext}}} q(v)$.

Both $E_p^{\bar{p}}, V_p$ have cardinality two or greater.

For V_p , we can hence choose two distinct vertices v_p, w_p . For $E_p^{\bar{p}}$, we can choose a partition into two non-empty sets $E_p^{\bar{p}} = F_p \cup G_p$. Set $q(v_p) = q_p$ and set $q(v) = 0$ for all other vertices in $p^{[0]}$.

Let now $c \in P_E(T)$ be a cut of $T \in C(\Gamma)$, so c is a collection of some edges of T . To such a collection of edges corresponds a set of decorations $D(c)$ at the endpoint (away from the root) of these edges. Each element in $D(c)$ plays the role of a decoration p above.

The corresponding \bar{p} decorate the vertices above p for those edges c .

For each $p \in D(c)$, choose v_p, w_p and partition $E_p^{\bar{p}} = F_p \cup G_p$.

Attach all edges in F_p to v_p and all edges in G_p to w_p . This defines, for each $T \in C(\Gamma)$ and each cut c of T , some new graph Γ_c^T , which clearly depends on choices of v_p, w_p and on choices of partitions of $E_p^{\bar{p}}$. Call these choices the choice of a 1-scale structure.

In particular, consider a complete forest, that is some T with all its edges chosen, $c = P_E(T)$. We chose a 1-scale structure for that complete forest f defining T . It induces a compatible choice for any other forest in the same given tree T (choices can be made independent between different trees in $C(\Gamma)$ though).

Do this for any $T \in C(\Gamma)$.

We now have a set of graphs $\Gamma_f = \sum_{c \in f_c} \Gamma_c^T$ for each forest of Γ . By construction, $|V_p| = 2$ for all $p, \bar{p} \in \Gamma_c^T$. Finally, we set $m_e = 0$ for all edges $e \notin (\Gamma_f/f)^{[1]}$.

Lemma 9. *i) For complete forests, we have the equivalence*

$$\sum_{f \text{ complete}} \Gamma_f \sim \Gamma.$$

ii) For any forest f ,

$$|C(\Gamma_f)| = 1.$$

Note that this implies that under such choices, no new singular sectors are generated. The graphs Γ_c^T are in particular free of overlapping divergences.


Proof. By assumption, all decorations p in any $T \in C(\Gamma)$ are log-divergent. For any choice, consider the pair \bar{p}, p , with $V_p \in p^{[0]}$ as described. Define $t_p^{\bar{p}}$ to be a minimal spanning tree of V_p on internal edges of p , so there is no other spanning tree of V_p in p which contains fewer edges.

Let $x_p \in \bar{p}^{[0]}$ be the vertex into which p is inserted. For any p, \bar{p} decorations of T with \bar{p} above p , and $T \in C(\Gamma)$, there are q, \bar{q} in $T_f = C(\Gamma_f)$ such that the minimal spanning tree $t_q^{\bar{q}} \subseteq t_p^{\bar{p}}$ is a path from v_p to w_p ⁷. Any not necessarily proper subset of cycles $\text{cyc} \subset \bar{p}$ containing x_p has $\text{sd}(\text{cyc}) \leq 0$, as \bar{p} is primitive by construction. As $\text{sd}(\text{cyc} - x_p \cup t_q^{\bar{q}}) < \text{sd}(\text{cyc})$, i) follows.


ii) is obvious. \square

Two examples might be helpful.

Example 10. The graph




say is connected through edges 4, 5, 6 to the graph




Also, at the vertex of edges 9, 10, 13, an external momentum enters Γ . So here, the set V_γ consists of four vertices, three vertices in $V_\gamma^{\Gamma/\gamma}$ and a single vertex in V_γ^{ext} . The graphs $\gamma, \Gamma/\gamma$ form a pair p, \bar{p} . Let x_p be the vertex 4, 5, 6 into which γ is inserted into Γ/γ .

⁷In theories with derivative couplings, where vertices might have negative weight in power-counting, one must (and can) demand that $\text{sd}(t_q^{\bar{q}}) > 0$.

The edges 11, 12, 13, 14 form a minimal spanning tree for the four vertices in $V\gamma$.
In




the connectors 4,5,6 have changed their endpoints. The divergent subgraph



is a 1-scale graph (in particular the external momentum at the vertex of edges 9, 10, 13 is sent to zero), and $\Gamma/\gamma = \Gamma_\gamma/\gamma^2$. V_{γ^2} consists of two vertices, with spanning tree the path t given by edges 12, 14. Any 1PI subgraph of Γ/γ is superficially convergent (if it is a proper subgraph) or log divergent (if it is Γ/γ itself) as Γ/γ is free of sub-divergences. In Γ_γ , such a sub-graph has the vertex x_p replaced by the path t . Its powercounting improves by the weight of the edges in t . So by construction, for any given forest f , we have a graph Γ_f where the subgraphs indexed by the forest are 1-scale graphs.

This ends the first example. Lets consider another one.

Example 11.



We know $|C(\Gamma)| = 2$. It has two maximal complete forests f_4, f_5 . A possible choice of 1-scale structures for them is

$$(82) \quad \Gamma_{f_4} = \begin{array}{c} \text{---} 1 \text{---} \\ \text{---} 2 \text{---} \\ \text{---} 3 \text{---} \\ \text{---} 4 \text{---} \\ \bullet \\ \text{---} 5 \text{---} \\ \text{---} 6 \text{---} \\ \bullet \end{array}, (m_1 = m_2 = m_3 = m_4 = 0)$$

$$(83) \quad \Gamma_{f_5} = \begin{array}{c} \text{---} 5 \text{---} \\ \text{---} 6 \text{---} \\ \text{---} 7 \text{---} \\ \text{---} 8 \text{---} \\ \bullet \\ \text{---} 9 \text{---} \\ \text{---} 10 \text{---} \\ \bullet \end{array}, (m_5 = m_6 = m_3 = m_4 = 0).$$

We then have $\Gamma_{f_0} = \Gamma$, $\Gamma_{f_1} = \Gamma_{m_3=m_4=0}$, $\Gamma_{f_2} = \Gamma_{f_4}$, $\Gamma_{f_3} = \Gamma_{f_5}$, where we refer to the forests in Eqs.(??-??).

We have now constructed graphs such that all divergent sub-graphs depend only on a single-scale. Still, the set $\{v \in \Gamma_f^{[0]} | q(v) \neq 0\}$ might have cardinality > 2 .

But then, we can partition this set into two non-empty subsets X, Y say and choose a vertex $x \in X, y \in Y$ for each. We then set $q(x) := \sum_{v \in X} q(v)$ and $q(y) := \sum_{v \in Y} q(v)$, and all other $q(v) = 0$. This gives a new graph Γ_f^2 which itself is a 1-scale graph.

Define $\phi_{2\Gamma_f} = \varphi_{\Gamma_f^2}$. Note that we use the massless φ on 1-scale graphs. Define furthermore

$$(84) \quad x_2^{\Gamma_f} := \frac{\phi_{\Gamma_f}(\Theta) - \varphi_{2\Gamma_f}}{\varphi_{2\Gamma_f}}.$$

Note that $x_2^{\Gamma_f}$ is independent of S, S_0, Θ_0 , by construction: $x_2^{\Gamma_f} = x_2^{\Gamma_f}(\Theta)$. We have

Lemma 12.

$$(85) \quad \Phi_{\text{fin}}(\Gamma) := \sum_f^{\emptyset} (-1)^{|f|} \int_{\mathbb{P}^n(\mathbb{R}_+)} M(x_2^{\Gamma_f})$$

converges. Here, the notation is defined similarly to Eqs.(??,??).

Proof. By construction, in each sector, singularities drop out, reflected by the scale independence of $x_2^{\Gamma_f}$, and we use the same forest sum as in the proof of convergence for Φ_{1-s}^R . Compared to renormalization by counterterms, where the remainder terms \bar{R}_f^Γ are replaced by $\phi_f(\Theta_0)\psi_{\Gamma/f}$, we now have different remainder terms $\bar{R}_f^{\Gamma_f}$ for them. This is due to the fact that $\phi_{\Gamma/f} = \phi_{\Gamma_f/f}$. Hence, along a forest f , $\phi_\Gamma = \phi_{\Gamma/f}\psi_f + \bar{R}_f^\Gamma$ and $\phi_{\Gamma_f} = \phi_{\Gamma_f/f}\psi_f + \bar{R}_f^{\Gamma_f}$. \square

4. DOUBLE ITERATION

Let us now define

$$(86) \quad x_{g,f}^\Gamma = \frac{\frac{S}{S_0}\phi_{\Gamma_f/g}(\Theta)\psi_g + \phi_g(\Theta_0)\psi_{\Gamma_f/g}}{\phi_{\Gamma_f/g}(\Theta_0)\psi_g + \phi_g(\Theta_0)\psi_{\Gamma_f/g}}$$

and we work with $M(x_{f,g}^\Gamma)$. We then have

$$(87) \quad \Phi^R(\Gamma) = \int_{\mathbb{P}^n(\mathbb{R}_+)} \sum_g^\emptyset (-1)^{|g|} M(x_{\emptyset,g}^\Gamma),$$

$$(88) \quad \Phi_{\text{fin}}(\Gamma) = \int_{\mathbb{P}^n(\mathbb{R}_+)} \sum_f^\emptyset (-1)^{|f|} M(x_{f,\emptyset}^\Gamma).$$

Finally, we define

$$(89) \quad \Phi_f^R(\Gamma) = \int_{\mathbb{P}^n(\mathbb{R}_+)} \sum_g^\emptyset (-1)^{|g|} M(x_{f,g}^\Gamma).$$

Then, first, (due to $\emptyset = \sum_j \bar{\sigma}_j$)

$$(90) \quad \Phi^R(\Gamma) = \sum_{j=1}^{\infty} \Phi_{\bar{\sigma}_j}^R(\Gamma),$$

which is a terminating sum on the rhs due to the finite co-radical filtration $\text{cor}(\Gamma)$ of every graph Γ .

Furthermore, we have our decomposition:

Theorem 13.

$$\Phi_{\bar{\sigma}_j}^R(\Gamma - \Gamma_2) = \sum_{k=0}^{j-1} c_k(\Theta, \Theta_0) L^k,$$

$$\Phi_{\bar{\sigma}_j}^R(\Gamma_2) = \sum_{k=1}^j d_k L^k.$$

In particular, $\Phi_{1-s}^R(\Gamma) := \sum_{j=1}^{\infty} \Phi_{\bar{\sigma}_j}^R(\Gamma_2)$.

This can be written in Sweedler's notation as (see Eq.(??))

$$(91) \quad \Phi_{1-s}^R(\Gamma) = \sum \Phi^{-1}(\Gamma'_2)(m_e = 0)(S_0) \Phi(\Gamma''_2)(S)(m_e = 0),$$

$$(92) \quad \equiv \Phi_2^{-1}(\Gamma') \Phi_2(\Gamma''),$$

where $\Phi_2(\Gamma) = \Phi(\Gamma_2)$ and 1-scale graphs Γ_2 are assumed massless. We use unrenormalized Feynman rules in the affine representation and 1-scale subgraphs which are evaluated to counterterms $\Phi^{-1}(\Gamma'_2)(m_e = 0)(S_0)$. The sum terminates at $\text{cor}(\Gamma)$.

By construction,

$$(93) \quad \Phi_{\text{fin}}(\Gamma)(\Theta) = \sum \Phi_2(\Gamma')^{-1} \Phi(\Gamma'').$$

Then, $\Phi_{1-s}(\Gamma) = \Phi^{-1}(S_0) \star \Phi(S)$, where it is understood that on the rhs of the equation sub- and co-graphs Γ'', Γ' of Γ are projected to 1-scale graphs Γ''_2, Γ'_2 . We finally have our decomposition

$$(94) \quad \Phi^R \equiv S_R^\Phi \star \Phi = \underbrace{(\Phi_2^{-1} \star \Phi(\Theta_0))^{-1}}_{\Phi_{\text{fin}}^{-1}(\Theta_0)} \star \Phi_{1-s}^R \star \underbrace{(\Phi_2^{-1} \star \Phi(\Theta))}_{\phi_{\text{fin}}(\Theta)}.$$

5. EXAMPLES

Next, we work out examples to acquaint the reader with our approach. These examples consider the decomposition into angles and scales. It is not our aim to do the final integrations, but to exhibit the structure of properly renormalized Feynman integrands, suitably decomposed as announced. In the following sections, we exclusively explore properties of G_{1-s} .

5.1. **Overall log.** We start with overall logarithmic divergent graphs.

5.1.1. *Primitive log.* Consider

$$(95) \quad \Gamma = \text{Diagram 1},$$

and

$$(96) \quad \Gamma^2 = \text{Diagram 2}.$$

The graph Γ is the one we are interested in, the 1-scale graph Γ^2 is supposed to have only two vertices at which $q(v) \neq 0$, and all internal masses in it are set to zero.

Trivially, $|C(\Gamma)| = 1$, with $C(\Gamma) = \{T\}$, and $T = \bullet_\Gamma$ primitive in the Hopf algebra of trees, and Γ is primitive in the Hopf algebra of graphs, $\Delta(\Gamma) = \Gamma \otimes \mathbb{I} + \mathbb{I} \otimes \Gamma$.

For the kinematics, we let $S = p_1^2 + p_2^2 + p_3^2 + 2p_1 \cdot p_2 + 2p_2 \cdot p_3 + 2p_3 \cdot p_1$ (which defines the variable angles $\Theta^{ij} = p_i \cdot p_j / S$, $\Theta^e = m_e^2 / S$) and subtract symmetrically say at S_0 , $\Theta_0^{ij} = \frac{1}{3}(4\delta_{ij} - 1)$ and $\Theta_0^e = m_e^2 / S_0$, which specifies the fixed angles Θ_0 .

The renormalized form is

$$(97) \quad \Phi_\Gamma^R = \frac{\ln \frac{\frac{S}{S_0} \phi_\Gamma(\Theta)}{\phi_\Gamma(\Theta_0)}}{\psi_\Gamma^2} \Omega_\Gamma.$$

To find the desired decomposition, we use

$$(98) \quad \Delta^2(\Gamma) = \Gamma \otimes \mathbb{I} \otimes \mathbb{I} + \mathbb{I} \otimes \Gamma \otimes \mathbb{I} + \mathbb{I} \otimes \mathbb{I} \otimes \Gamma.$$

We then have

$$(99) \quad \Phi_\Gamma^R = \Phi_{\text{fin}}^{-1}(\Theta_0)(\Gamma) + \Phi_{1-s}^R(S/S_0)(\Gamma) + \Phi_{\text{fin}}(\Theta)(\Gamma).$$

We have

$$(100) \quad \Phi_{\text{fin}}^{-1}(\Theta_0)(\Gamma) = -\frac{\ln \frac{\phi_\Gamma(\Theta_0)}{\psi_{\Gamma^2} \bullet}}{\psi_\Gamma^2} \Omega_\Gamma,$$


$$(101) \quad \Phi_{1-s}^R(S/S_0)(\Gamma) = \frac{\ln \frac{S}{S_0}}{\psi_{\Gamma^2}^2} \Omega_\Gamma,$$

which integrates to the renormalized value $\Phi_{1-s}^R(S/S_0)(\Gamma) = 6\zeta(3) \ln \frac{S}{S_0}$. Finally,


$$(102) \quad \Phi_{\text{fin}}(\Theta)(\Gamma) = \frac{\ln \frac{\phi_\Gamma(\Theta)}{\psi_\Gamma^2 \bullet}}{\psi_\Gamma^2} \Omega_\Gamma.$$

These integrands indeed all converge, which is synonymous for us to say that they can be integrated against $\mathbb{P}^{|\Gamma^{[1]}|-1}(\mathbb{R}_+)$.


5.1.2. *Log with log sub.* Consider the graph



It has a subgraph




and a co-graph



It suffices to consider a small Hopf algebra (co-)generated by $\mathbb{I}, \Gamma, \gamma, \Gamma/\gamma$, with the only non-trivial co-product

$$(106) \quad \Delta(\Gamma) = \Gamma \otimes \mathbb{I} + \mathbb{I} \otimes \Gamma + \gamma \otimes \Gamma/\gamma.$$

Obviously, $|C(\Gamma)| = 1$. We have $C(\Gamma) = \{T\}$, with



with T just a tree on two vertices⁸.

Forests are $f_0 = \emptyset$ and $f_1 = \gamma$.

⁸In light of [?], note that we can write

$$(108) \quad T = B_+^{\Gamma/\gamma}(\bullet_\gamma),$$

The renormalized integrand is then

$$(110) \quad \Phi_{\Gamma}^R(S/S_0, \Theta, \Theta_0) = \left\{ \frac{\ln \frac{S}{S_0} \phi_{\Gamma}(\Theta)}{\psi_{\Gamma}^2} - \frac{\ln \frac{S}{S_0} \phi_{\Gamma/\gamma}(\Theta) \psi_{\gamma} + \phi_{\gamma}(\Theta_0) \psi_{\Gamma/\gamma}}{\psi_{\Gamma/\gamma}^2 \psi_{\gamma}^2} \right\} \Omega_{\Gamma}.$$

One immediately checks that this is integrable in the edge variables against $\mathbb{P}^{13}(\mathbb{R}_+)$, using nothing more than our cherished remainder properties of the graph polynomials Eqs.(??,??).

For the decomposition as a twisted conjugation, we stick with our conventions that a superscript 2 indicated that a graph is 1-scale with respect to its external momenta flow ($q(v) \neq 0$ only for two vertices, all internal masses zero), while a subscript f means that the graphs in the forest f are made 1-scale as subgraphs, also with zero internal masses in them. Then, $\Gamma_{\emptyset} = \Gamma$ (the forest which is made 1-scale is empty), and we choose

$$(111) \quad \Gamma^2 \equiv \Gamma_{\emptyset}^2 = \text{Diagram},$$

$$(112) \quad \Gamma_{\gamma} = \text{Diagram},$$

and

$$(113) \quad \Gamma_{\gamma}^2 = \text{Diagram}.$$

and $B_+^{\Gamma/\gamma}$ a Hochschild closed map for this Hopf algebra:

$$(109) \quad \rho(\Gamma) = T, \Delta_T(T) = T \otimes \mathbb{I} + \mathbb{I} \otimes T + \bullet_{\gamma} \otimes \bullet_{\Gamma/\gamma} = \Delta_T B_+^{\Gamma/\gamma}(\bullet_{\gamma}),$$

with $\Delta_T B_+^{\Gamma/\gamma} = B_+^{\Gamma/\gamma} \otimes \mathbb{I} + (\text{id} \otimes B_+^{\Gamma/\gamma}) \Delta_T$.

This gives also a new sub-graph γ^2 of Γ_γ ,

$$(114) \quad \gamma^2 = \text{Diagram of a circle with two black dots at the top and bottom. The circle is divided into four quadrants by a horizontal line (edges 11, 13) and a vertical line (edges 14, 12). The top-left quadrant is labeled 7, the top-right 10, the bottom-left 8, and the bottom-right 9.$$

and a new co-graph of Γ_\emptyset^2 ,

$$(115) \quad \Gamma_\gamma^2/\gamma^2 = \Gamma_\emptyset^2/\gamma = \text{Diagram of a circle with two black dots at the top and left. The circle is divided into four quadrants by a horizontal line (edges 6, 3) and a vertical line (edges 5, 4). The top-left quadrant is labeled 1, the top-right 3, the bottom-left 2, and the bottom-right 4.$$

We have by definition

$$(116) \quad \phi_{\Gamma_\emptyset} = \phi_\Gamma = S\phi_\Gamma(\Theta),$$

$$(117) \quad \phi_{\Gamma_\emptyset^2} = S\psi_{\Gamma_\emptyset^2\bullet},$$

and

$$(118) \quad \phi(\Gamma_\gamma) = S\phi_{\Gamma/\gamma}(\Theta)\psi_\gamma + S\psi_{\gamma\bullet}\psi_{\Gamma/\gamma}.$$

Also,

$$(119) \quad \phi(\Gamma_\gamma^2) = S\psi_{\Gamma/\gamma\bullet}\psi_\gamma + S\psi_{\gamma\bullet}\psi_{\Gamma/\gamma}.$$

It follows that

$$(120) \quad 1 + x_2^{\Gamma_\emptyset} = \frac{\phi_{\Gamma_\emptyset}(\Theta)}{\psi_{\Gamma_\emptyset^2\bullet}}$$

and

$$(121) \quad 1 + x_2^{\Gamma_\gamma} = \frac{\phi_{\Gamma_\gamma}(\Theta)}{\psi_{\Gamma_\gamma^2/\gamma^2\bullet}\psi_\gamma + \psi_{\gamma^2\bullet}\psi_{\Gamma_\gamma^2/\gamma^2}}.$$

We also need

$$(122) \quad \Gamma_\gamma^2/\gamma^2\bullet = \text{Diagram of a circle with two black dots at the top and left. The circle is divided into four quadrants by a horizontal line (edges 6, 3) and a vertical line (edges 5, 4). The top-left quadrant is labeled 1, the top-right 3, the bottom-left 2, and the bottom-right 4. There is a loop at the top vertex.$$

and

$$(123) \quad G = \text{Diagram of a circle with two black dots at the top and left. The circle is divided into four quadrants by a horizontal line (edges 6, 3) and a vertical line (edges 5, 4). The top-left quadrant is labeled 1, the top-right 3, the bottom-left 2, and the bottom-right 4. There are loops at both the top and left vertices.$$

We have $\Gamma_\gamma^2/\gamma^2 = \Gamma_\emptyset^2/\gamma$ and $\Gamma/\gamma = \Gamma_\gamma/\gamma^2$. Also, $\psi_\Gamma = \psi_{\Gamma_\emptyset^2}$, $\psi_{\Gamma_\gamma} = \psi_{\Gamma_\gamma^2}$, and $\psi_\gamma = \psi_{\gamma^2}$.

To evaluate $\Phi_{\text{fin}}^{-1}(\Theta_0) \star \Phi_{1-s}^R(S/S_0) \star \Phi_{\text{fin}}(\Theta)$, we need

$$(124) \quad \Delta^2(\Gamma) = \Gamma \otimes \mathbb{I} \otimes \mathbb{I} + \mathbb{I} \otimes \Gamma \otimes \mathbb{I} + \mathbb{I} \otimes \mathbb{I} \otimes \Gamma + \mathbb{I} \otimes \gamma \otimes \Gamma / \gamma + \gamma \otimes \mathbb{I} \otimes \Gamma / \gamma + \gamma \otimes \Gamma / \gamma \otimes \mathbb{I}.$$

We find

$$(125) \quad \begin{aligned} \Phi_{\Gamma}^R &= \Phi_{\text{fin}}(\Theta)(\Gamma) + \Phi_{\text{fin}}^{-1}(\Theta_0)(\Gamma) + \Phi_{1-s}^R(\gamma)\Phi_{\text{fin}}(\Theta)(\Gamma/\gamma) \\ &+ \Phi_{1-s}^R(\Gamma) + \Phi_{\text{fin}}^{-1}(\Theta_0)(\gamma)\Phi_{\text{fin}}(\Theta)(\Gamma/\gamma) + \Phi_{\text{fin}}^{-1}(\Theta_0)(\gamma)\Phi_{1-s}^R(\Gamma/\gamma). \end{aligned}$$

Note that all the terms on the rhs can be expressed using forest sums as in Eq.(??):

$$(126) \quad \Phi_{\Gamma}^R = \underbrace{+\Phi_{\bar{\sigma}_1}^R(\Gamma - \Gamma^2)}_{\mathcal{O}(1)} + \underbrace{\Phi_{\bar{\sigma}_2}^R(\Gamma - \Gamma^2)}_{\mathcal{O}(L)}$$

$$(127) \quad \underbrace{+\Phi_{\bar{\sigma}_1}^R(\Gamma^2)}_{\mathcal{O}(L)} + \underbrace{\Phi_{\bar{\sigma}_2}^R(\Gamma^2)}_{=\Phi_{1-s}^R(\Gamma), \mathcal{O}(L^2)},$$

indicating orders in $L = \ln S/S_0$. We have in particular (we invite the reader to confirm convergence of these expressions again directly from the remainder properties Eqs.(??,??))

$$(128) \quad \Phi_{\text{fin}}(\Theta)(\Gamma) = \left\{ \frac{\ln \frac{\phi_{\Gamma}(\Theta)}{\psi_{\Gamma_0^2}^{\bullet}}}{\psi_{\Gamma_0^2}^2} - \frac{\ln \frac{\phi_{\Gamma/\gamma}(\Theta)}{\psi_{\Gamma/\gamma^2}^{\bullet}}}{\psi_{\Gamma/\gamma^2}^2} \right\} \Omega_{\Gamma}.$$

Furthermore

$$(129) \quad \Phi_{1-s}^R(S/S_0)(\Gamma) = \left\{ \frac{\ln \frac{S}{S_0}}{\psi_{\Gamma^2}^2} - \frac{\overbrace{\ln \frac{S}{S_0} \frac{\psi_{\Gamma^2/\gamma^2}^{\bullet} \psi_{\gamma^2} + \psi_{\gamma^2}^{\bullet} \psi_{\Gamma^2/\gamma^2}}{\psi_{\Gamma^2/\gamma^2}^{\bullet} \psi_{\gamma^2} + \psi_{\gamma^2}^{\bullet} \psi_{\Gamma^2/\gamma^2}}}_{\Upsilon_{\gamma^2; \Gamma^2}(\frac{S}{S_0})}}{\psi_{\Gamma^2/\gamma^2}^2 \psi_{\gamma^2}^2} \right\} \Omega_{\Gamma}$$

The term $\sim \ln^2(S/S_0)$ in Φ_{Γ}^R , generated by Φ_{1-s} only, is obviously $60\zeta(3)\zeta(5)$ by our considerations, while (??) reveals three sources for a term linear in $L = \ln(S/S_0)$: First, we have


$$(130) \quad \int \frac{\psi_{\Gamma/\gamma^2}^{\bullet}}{\psi_{\Gamma^2/\gamma^2}^2 \psi_{\gamma^2} \Upsilon_{\gamma^2; \Gamma^2}(1)} \Omega_{\Gamma},$$

(patently inaccessible by standard methods) but note that $\Upsilon_{\gamma^2; \Gamma^2}(1) = \psi_G$, so that this expression is reduced to the study of graph hypersurfaces, started in [?, ?].


For the other two terms linear in L , they give $\Phi_{\text{fin}}^{-1}(\Theta_0)(\gamma)\Phi_{1-s}^R(\Gamma/\gamma)$ (which is $20\zeta(5)$ times an unknown function of angles Θ_0 , also easily identified in the second term on the right in Eq.(??)), as well as $\Phi_{1-s}^R(\gamma)\Phi_{\text{fin}}(\Theta)(\Gamma/\gamma)$ (which is $6\zeta(3)$ times an unknown function of angles Θ , see also the first term on the right in Eq.(??)).

The constant term in L is again an unknown function of angles Θ, Θ_0 , which equals $\Phi_{\Gamma}^R(1, \Theta, \Theta_0)$.


5.1.3. *Log with quadratic sub.* This is an interesting example as it will involve an overall convergent contribution $l_{-1}^{[0]}$. Kinematics is as before. Let

(131) $\Gamma =$ 


and


(132) $\bar{\Gamma} =$ 

The subgraph is

(133) $\gamma =$ 

The cographs are

(134) $\Gamma/\gamma =$ 

(135) $\bar{\Gamma}/\gamma =$ 

The Hopf algebra structure is simply

(136)
$$\Delta(\Gamma) = \Gamma \otimes \mathbb{I} + \mathbb{I} \otimes \Gamma + \gamma \otimes \Gamma/\gamma,$$

as co-graphs which are tadpoles vanish.

We have

$$\begin{aligned}
 (137) \quad \Phi_\Gamma^R &= \frac{\psi_\gamma \bullet}{\psi_\gamma} \left\{ \frac{l_0^{[0]}(x_\Gamma^\bar{\Gamma})}{\psi_\Gamma^2} - \frac{l_0^{[0]}(x_\gamma^\bar{\Gamma})}{\psi_{\Gamma/\gamma}^2 \psi_\gamma^2} \right\} \Omega_{\bar{\Gamma}} \\
 &+ m_\gamma^2 \frac{\bar{\psi}_\gamma \bullet}{\psi_\gamma} \left\{ \left(\frac{S_0 \phi_\Gamma(\Theta_0)}{\psi_\Gamma} \right)^{-1} \frac{l_{-1}^{[0]}(x_\Gamma^\Gamma)}{\psi_\Gamma^2} \right. \\
 &\quad \left. - \left(\frac{S_0 \phi_{\Gamma/\gamma}(\Theta_0) \psi_\gamma + S_0 \bar{\psi}_\gamma \bullet \psi_{\Gamma/\gamma}}{\psi_{\Gamma/\gamma}} \right)^{-1} \frac{l_{-1}^{[0]}(x_\gamma^\Gamma)}{\psi_{\Gamma/\gamma}^2 \psi_\gamma^2} \right\} \Omega_\Gamma \\
 &- m_\gamma^2 \frac{\bar{\psi}_\gamma \bullet}{\psi_\gamma} \frac{\psi_\gamma \bullet}{\psi_\gamma} \left\{ \left(\frac{S_0 \phi_{\bar{\Gamma}}(\Theta_0)}{\psi_{\bar{\Gamma}}} \right)^{-1} \frac{l_{-1}^{[0]}(x_\Gamma^\Gamma)}{\psi_{\bar{\Gamma}}^2} \right. \\
 &\quad \left. - \left(\frac{S_0 \phi_{\bar{\Gamma}/\gamma}(\Theta_0) \psi_\gamma + S_0 \bar{\psi}_\gamma \bullet \psi_{\bar{\Gamma}/\gamma}}{\psi_{\bar{\Gamma}/\gamma}} \right)^{-1} \frac{l_{-1}^{[0]}(x_\gamma^\Gamma)}{\psi_{\bar{\Gamma}/\gamma}^2 \psi_\gamma^2} \right\} \Omega_{\bar{\Gamma}}.
 \end{aligned}$$

The first line above and the coefficients of m_γ^2 in the two following lines allow for a separate decomposition in G_{1-s}, G_{fin} as expected: the first line depends on scales as well as angles and decomposes as before, the two following lines only depend on angles, taking all definitions into account. Indeed, this must be so as we have an overall dimensionless quantity, and those two lines come from a superficial convergent contribution as they factorize m_γ^2 , and hence the angle m_γ^2/S , multiplying it all out. Here,

$$(138) \quad l_{-1}^{[0]}(x_\Gamma^\Gamma) = \frac{1}{1 + \frac{\frac{S}{S_0}(\phi_\Gamma(\Theta) - \phi_\Gamma(\Theta_0))}{\phi_\Gamma(\Theta_0)}} - 1,$$

and

$$(139) \quad l_{-1}^{[0]}(x_\gamma^\Gamma) = \frac{1}{1 + \frac{\frac{S}{S_0}(\phi_{\Gamma/\gamma}(\Theta) \psi_\gamma - \phi_{\Gamma/\gamma}(\Theta_0) \psi_\gamma)}{\phi_{\Gamma/\gamma}(\Theta_0) \psi_\gamma + \psi_\gamma \bullet \psi_{\Gamma/\gamma}}} - 1,$$

similarly for $l_{-1}^{[0]}(x_\Gamma^\bar{\Gamma})$ and $l_{-1}^{[0]}(x_\gamma^\bar{\Gamma})$.

5.2. Overall quadratic. Next, we turn to overall quadratic amplitudes.

5.2.1. Primitive overall quadratic. We look at the graph

$$(140) \quad \Gamma = \text{---} \bullet \text{---} \bigcirc \text{---} \bullet \text{---} .$$

$C(\Gamma) = \{\bullet_\Gamma\}$. Kinematics are given by $S = q^2, S_0 = m^2, \Theta_{q^2} = (1 - \frac{m^2}{q^2}), \Theta_{q^2}^0 = (1 - \frac{q^2}{m^2}), \Theta_e = m_e^2/q^2, \Theta_e^0 = m_e^2/m^2$. We find immediately

$$\begin{aligned}
 (141) \quad \Phi_\Gamma^R &= (q^2 - m^2) \frac{A_2 A_3 A_4 \ln(1 + x_\Gamma^\Gamma) \Omega_\Gamma}{\psi_\Gamma \psi_\Gamma^2} \\
 &+ m^2 \frac{A_2 A_3 A_4 + (A_2 \Theta_2^0 + A_3 \Theta_3^0 + A_4 \Theta_4^0) \psi_\Gamma \ln^{[1]}(1 + x_\Gamma^\Gamma) \Omega_\Gamma}{\psi_\Gamma \psi_\Gamma^2}.
 \end{aligned}$$

Here,

$$(142) \quad x_\emptyset^\Gamma = \frac{\frac{q^2}{m^2} \Theta_{q^2} A_2 A_3 A_4}{(A_2 A_3 A_4 + (A_2 \Theta_2^0 + A_3 \Theta_3^0 + A_4 \Theta_4^0) \psi_\Gamma)}.$$

This makes it evident that we fulfil the renormalization conditions.

We set $\Phi_\Gamma^R =: (q^2 - m^2) \Phi_\Gamma^{(q^2 - m^2); R} + m^2 \Phi_\Gamma^{m^2; R} + m_2^2 \Phi_\Gamma^{m_2^2; R} + m_3^2 \Phi_\Gamma^{m_3^2; R} + m_4^2 \Phi_\Gamma^{m_4^2; R}$ and can read off the coefficient functions $\Phi_\Gamma^{x; R}$ immediately. Note that they are all dimensionless quantities.

The decomposition into angles and scales happens now for these coefficient functions.

$$(143) \quad \Phi_{1-s}^{q^2 - m^2; R}(\Gamma) = \frac{A_2 A_3 A_4}{\psi_\Gamma} \ln \left(\frac{q^2}{m^2} \right) \frac{\Omega_\Gamma}{\psi_\Gamma^2},$$

$$(144) \quad \Phi_{1-s}^{m^2; R}(\Gamma) = \frac{A_2 A_3 A_4}{\psi_\Gamma} \ln \left(\frac{q^2}{m^2} \right) \frac{\Omega_\Gamma}{\psi_\Gamma^2},$$

$$(145) \quad \Phi_{1-s}^{m_2^2; R}(\Gamma) = \frac{A_2}{1} \ln \left(\frac{q^2}{m^2} \right) \frac{\Omega_\Gamma}{\psi_\Gamma^2},$$

$$(146) \quad \Phi_{1-s}^{m_3^2; R}(\Gamma) = \frac{A_3}{1} \ln \left(\frac{q^2}{m^2} \right) \frac{\Omega_\Gamma}{\psi_\Gamma^2},$$

$$(147) \quad \Phi_{1-s}^{m_4^2; R}(\Gamma) = \frac{A_4}{1} \ln \left(\frac{q^2}{m^2} \right) \frac{\Omega_\Gamma}{\psi_\Gamma^2}.$$

For the angles:

$$(148) \quad \Phi_{\text{fin}}^{q^2 - m^2; R}(\Theta)(\Gamma) = \frac{A_2 A_3 A_4}{\psi_\Gamma} \ln \left(\frac{\varphi_\Gamma + \psi_\Gamma \sum_{j=2}^4 \Theta_j A_j}{\phi_\Gamma} \right) \frac{\Omega_\Gamma}{\psi_\Gamma^2},$$

$$(149) \quad \Phi_{\text{fin}}^{m^2; R}(\Theta)(\Gamma) = \frac{A_2 A_3 A_4}{\psi_\Gamma} \ln \left(\frac{\varphi_\Gamma + \psi_\Gamma \sum_{j=2}^4 \Theta_j A_j}{\phi_\Gamma} \right) \frac{\Omega_\Gamma}{\psi_\Gamma^2},$$

$$(150) \quad \Phi_{\text{fin}}^{m_2^2; R}(\Theta)(\Gamma) = \frac{A_2}{1} \ln \left(\frac{\phi_\Gamma + \psi_\Gamma \sum_{j=2}^4 \Theta_j A_j}{\varphi_\Gamma} \right) \frac{\Omega_\Gamma}{\psi_\Gamma^2},$$


$$(151) \quad \Phi_{\text{fin}}^{m_3^2; R}(\Theta)(\Gamma) = \frac{A_3}{1} \ln \left(\frac{\phi_\Gamma + \psi_\Gamma \sum_{j=2}^4 \Theta_j A_j}{\varphi_\Gamma} \right) \frac{\Omega_\Gamma}{\psi_\Gamma^2},$$

$$(152) \quad \Phi_{\text{fin}}^{m_4^2; R}(\Theta)(\Gamma) = \frac{A_4}{1} \ln \left(\frac{\phi_\Gamma + \psi_\Gamma \sum_{j=2}^4 \Theta_j A_j}{\varphi_\Gamma} \right) \frac{\Omega_\Gamma}{\psi_\Gamma^2}.$$


and

$$\begin{aligned} \Phi_{\text{fin}}^{q^2-m^2;R}(\Theta^0)(\Gamma) &= \frac{A_2 A_3 A_4}{\psi_\Gamma} \ln \left(\frac{\varphi_\Gamma + \psi_\Gamma \sum_{j=2}^4 \Theta_j^0 A_j}{\phi_\Gamma} \right) \frac{\Omega_\Gamma}{\psi_\Gamma^2}, \\ \Phi_{\text{fin}}^{m^2;R}(\Theta^0)(\Gamma) &= \frac{A_2 A_3 A_4}{\psi_\Gamma} \left(\ln \left(\frac{\varphi_\Gamma + \psi_\Gamma \sum_{j=2}^4 \Theta_j^0 A_j}{\phi_\Gamma} \right) \right. \\ &\quad \left. - \frac{\Theta_{q^2}^0 \varphi_\Gamma}{\varphi_\Gamma + \psi_\Gamma \sum_{j=2}^4 \Theta_j^0 A_j} \right) \frac{\Omega_\Gamma}{\psi_\Gamma^2}, \\ \Phi_{\text{fin}}^{m_2^2;R}(\Theta^0)(\Gamma) &= \frac{A_2}{1} \left(\ln \left(\frac{\varphi_\Gamma + \psi_\Gamma \sum_{j=2}^4 \Theta_j^0 A_j}{\phi_\Gamma} \right) - \frac{\Theta_{q^2}^0 \phi_\Gamma}{\varphi_\Gamma + \psi_\Gamma \sum_{j=2}^4 \Theta_j^0 A_j} \right) \frac{\Omega_\Gamma}{\psi_\Gamma^2}, \\ \Phi_{\text{fin}}^{m_3^2;R}(\Theta^0)(\Gamma) &= \frac{A_3}{1} \left(\ln \left(\frac{\varphi_\Gamma + \psi_\Gamma \sum_{j=2}^4 \Theta_j^0 A_j}{\phi_\Gamma} \right) - \frac{\Theta_{q^2}^0 \phi_\Gamma}{\varphi_\Gamma + \psi_\Gamma \sum_{j=2}^4 \Theta_j^0 A_j} \right) \frac{\Omega_\Gamma}{\psi_\Gamma^2}, \\ \Phi_{\text{fin}}^{m_4^2;R}(\Theta^0)(\Gamma) &= \frac{A_4}{1} \left(\ln \left(\frac{\varphi_\Gamma + \psi_\Gamma \sum_{j=2}^4 \Theta_j^0 A_j}{\phi_\Gamma} \right) - \frac{\Theta_{q^2}^0 \phi_\Gamma}{\varphi_\Gamma + \psi_\Gamma \sum_{j=2}^4 \Theta_j^0 A_j} \right) \frac{\Omega_\Gamma}{\psi_\Gamma^2}. \end{aligned}$$


5.2.2. *Overall quadratic with quadratic sub.* The graph Γ which we want to consider is




We also need the graph




We have the sub-graph




and the co-graphs




and



We also need



and



For any X as needed we write

$$(153) \quad \bar{\psi}_X = \frac{1}{m^2} \phi_X(m^2, \Theta^0) = \psi_{X^\bullet} + \left(\sum_{e \in X} \Theta_e^0 A_e \right) \psi_X$$

and

$$(154) \quad \phi_X(q^2, \Theta) = q^2 \left(\psi_{X^\bullet} + \left(\sum_{e \in X} \Theta_e A_e \right) \psi_X \right).$$

We renormalize the subgraph at m_γ^2 and Γ at m^2 . Then, by Thm.??,

$$(155) \quad \Phi_\Gamma^R(q^2, m^2) =$$

$$(156) \quad + \frac{\psi_{\bar{\Gamma}^\bullet} \psi_{\gamma^\bullet}}{\psi_{\bar{\Gamma}} \psi_\gamma} l_1^{[1]}(x_{\bar{\theta}}) \frac{\Omega_{\bar{\Gamma}}}{\psi_{\bar{\Gamma}}^2}$$

$$(157) \quad - \frac{\psi_{\bar{\Gamma}/\gamma} \psi_{\gamma^\bullet}}{\psi_{\bar{\Gamma}/\gamma} \psi_\gamma} l_1^{[1]}(x_{\bar{\gamma}}) \frac{\Omega_{\bar{\Gamma}}}{\psi_{\bar{\Gamma}/\gamma}^2 \psi_\gamma^2}$$

$$(158) \quad + m_\gamma^2 \frac{\bar{\psi}_{\gamma^\bullet}}{\psi_\gamma} l_0^{[1]}(x_\theta) \frac{\Omega_\Gamma}{\psi_\Gamma^2}$$

$$(159) \quad - m_\gamma^2 \frac{\bar{\psi}_{\gamma^\bullet}}{\psi_\gamma} l_0^{[1]}(x_\gamma) \frac{\Omega_\Gamma}{\psi_{\bar{\Gamma}/\gamma}^2 \psi_\gamma^2}$$

$$(160) \quad - m_\gamma^2 \frac{\bar{\psi}_{\gamma^\bullet} \psi_{\gamma^\bullet}}{\psi_\gamma \psi_\gamma} l_0^{[1]}(x_{\bar{\theta}}) \frac{\Omega_{\bar{\Gamma}}}{\psi_{\bar{\Gamma}}^2}$$

$$(161) \quad + m_\gamma^2 \frac{\bar{\psi}_{\gamma^\bullet} \psi_{\gamma^\bullet}}{\psi_\gamma \psi_\gamma} l_0^{[1]}(x_{\bar{\gamma}}) \frac{\Omega_{\bar{\Gamma}}}{\psi_{\bar{\Gamma}/\gamma}^2 \psi_\gamma^2}.$$

Here, $\Omega_{\bar{\Gamma}}$ is the five-form $A_2 dA_3 \wedge \cdots \wedge dA_7 \pm \cdots$, and Ω_{Γ} the six-form $A_1 dA_2 \wedge \cdots \wedge dA_7 \pm \cdots$. Furthermore,

$$(162) \quad l_1^{[1]}(x_{\emptyset}^{\bar{\Gamma}}) = (q^2 - m^2) \ln \left(1 + \frac{(q^2 - m^2)\psi_{\bar{\Gamma}\bullet}}{m^2\bar{\psi}_{\bar{\Gamma}\bullet}} \right)$$

$$(163) \quad + \frac{\bar{\psi}_{\bar{\Gamma}\bullet}}{\psi_{\bar{\Gamma}\bullet}} \left\{ m^2 \ln \left(1 + \frac{(q^2 - m^2)\psi_{\bar{\Gamma}\bullet}}{m^2\bar{\psi}_{\bar{\Gamma}\bullet}} \right) \right.$$

$$(164) \quad \left. - m^2 \frac{(q^2 - m^2)\psi_{\bar{\Gamma}\bullet}}{m^2\bar{\psi}_{\bar{\Gamma}\bullet}} \right\}.$$

and

$$(165) \quad l_1^{[1]}(x_{\gamma}^{\bar{\Gamma}}) = (q^2 - m^2) \ln \left(1 + \frac{(q^2 - m^2)\psi_{\bar{\Gamma}/\gamma\bullet}\psi_{\gamma}}{m^2\bar{\psi}_{\bar{\Gamma}/\gamma\bullet}\psi_{\gamma} + \psi_{\gamma\bullet}\psi_{\Gamma/\gamma}} \right)$$

$$(166) \quad + \frac{\bar{\psi}_{\bar{\Gamma}/\gamma\bullet}}{\psi_{\bar{\Gamma}/\gamma\bullet}} \left\{ m^2 \ln \left(1 + \frac{(q^2 - m^2)\psi_{\bar{\Gamma}/\gamma\bullet}\psi_{\gamma}}{m^2\bar{\psi}_{\bar{\Gamma}/\gamma\bullet}\psi_{\gamma} + \psi_{\gamma\bullet}\psi_{\Gamma/\gamma}} \right) \right.$$

$$(167) \quad \left. - m^2 \frac{(q^2 - m^2)\psi_{\bar{\Gamma}\bullet}\psi_{\gamma}}{m^2\bar{\psi}_{\bar{\Gamma}\bullet}\psi_{\gamma} + \psi_{\gamma\bullet}\psi_{\Gamma/\gamma}} \right\}.$$

$$(168) \quad l_0^{[1]}(x_{\emptyset}^{\Gamma}) = \ln \left(1 + \frac{(q^2 - m^2)\psi_{\Gamma\bullet}}{m^2\bar{\psi}_{\Gamma\bullet}} \right) - \frac{(q^2 - m^2)\psi_{\Gamma\bullet}}{m^2\bar{\psi}_{\Gamma\bullet}},$$

and

$$(169) \quad l_0^{[1]}(x_{\gamma}^{\Gamma}) = \ln \left(1 + \frac{(q^2 - m^2)\psi_{\Gamma/\gamma\bullet}\psi_{\gamma}}{m^2\bar{\psi}_{\Gamma/\gamma\bullet}\psi_{\gamma} + \psi_{\gamma\bullet}\psi_{\Gamma/\gamma}} \right) - \frac{(q^2 - m^2)\psi_{\Gamma/\gamma\bullet}\psi_{\gamma}}{m^2\bar{\psi}_{\Gamma/\gamma\bullet}\psi_{\gamma} + \psi_{\gamma\bullet}\psi_{\Gamma/\gamma}},$$

and similarly for $l_0^{[1]}(x_{\emptyset}^{\bar{\Gamma}})$ and $l_0^{[1]}(x_{\gamma}^{\bar{\Gamma}})$.

One now immediately checks that lines (??)+(??) add to a convergent expression which fulfils the renormalization conditions, and so do (??)+(??) and (??)+(??).

From here on, decomposition into Φ_{1-s} and Φ_{fin} is lengthy but straightforward.

6. GRAPH POLYNOMIALS AND 1-SCALE GRAPHS


6.1. Single-scale graphs.

Definition 14. A *1-scale graph* is a finite disjoint union of connected graphs, each of which has exactly two vertices whose external legs carry non-zero momenta, and with vanishing internal masses m_e for all edges e .

Let G be a connected 1-scale graph with distinguished vertices $v_1 \neq v_2$. Let \bar{G} denote the graph obtained from G by adding a new edge e connecting v_1 and v_2 , and let G^\bullet be the graph obtained from G by identifying v_1 and v_2 . The graphs G , G^\bullet are obtained from \bar{G} by deleting (contracting) the edge e respectively. Thus

$$\psi_{\bar{G}} = \psi_G \alpha_e + \psi_{G^\bullet}.$$

If $G = \cup_{i=1}^n G_i$ has n connected components, then we define G^\bullet to be $\cup_{i=1}^n G_i^\bullet$.



Lemma 15. *Let G be a connected, 1-scale graph with external momentum q . Then the second Symanzik polynomial $\phi(q, G)$ is given by*

$$(170) \quad \phi(q, G) = q^2 \psi_{G^\bullet} .$$

Proof. This follows immediately from the definition of $\phi(q, G)$. \square

We henceforth write ϕ_G for ψ_{G^\bullet} , since all q -dependencies are trivial. Clearly, $\deg \phi_G = \deg \psi_G + 1$. If $G = \cup_{i=1}^n G_i$ with G_i connected, we define

$$(171) \quad \psi_G = \prod_{i=1}^n \psi_{G_i} \quad \text{and} \quad \phi_G = \sum_{i=1}^n \phi_{G_i} \prod_{j \neq i} \psi_{G_j} .$$

Note that this convention is *not* compatible with the contraction-deletion relations, which require that a non-connected graph have vanishing graph polynomial.

Definition 16. Let G be a connected 1-scale graph, and let $I \subset E(G)$ be a set of edges of G . Define $\varepsilon_G(I) \in \{0, 1\}$ to be 1 if I meets both distinguished vertices in G , and 0 otherwise. Equivalently, $\varepsilon_G(I)$ is 0 if the image of I in G^\bullet is isomorphic to I , and equal to 1 otherwise. If $G = \cup_{i=1}^n G_i$, then we define $\varepsilon_G = \prod_{i=1}^n \varepsilon_{G_i}$.

Suppose that G is a connected graph with vertices of degree at most 4, and let $\gamma \subsetneq G$ be a connected subgraph. Define the superficial degree of divergence of γ by

$$sd(\gamma) = 2h_\gamma - E_\gamma .$$

Since γ is a strict subgraph, the average valency of its edges α is < 4 , and hence $E_\gamma < 2V_\gamma$. By Euler's formula $E_\gamma - V_\gamma = h_\gamma - 1$, this implies that $sd(\gamma) \leq 1$. We say that γ is

- *convergent* if $sd(\gamma) < 0$,
- *log-divergent* if $sd(\gamma) = 0$,
- *quadratically divergent* if $sd(\gamma) = 1$.

We say that G has at most log-divergent subgraphs if $2h_I \leq |I|$ for all $I \subset E(G)$.

Remark 17. *Let G be connected in ϕ^4 . Suppose that G has v_i vertices of valency i , where $1 \leq i \leq 4$. A simple computation using Euler's formula shows that*

$$(172) \quad v_3 + 2v_2 = 4 - 2sd(G) .$$

In particular, suppose that G is quadratically divergent. Then $v_3 + 2v_2 = 2$ and (v_2, v_3) is either $(1, 0)$ or $(0, 2)$. In particular, every non-trivial connected quadratic subgraph of a ϕ^4 graph has precisely two 3-valent vertices.

6.2. Circular joins. Let G_1, \dots, G_n denote $n \geq 2$ connected 1-scale graphs.

Definition 18. Label the distinguished vertices of each graph G_i by v_i, w_i . A *circular join* $c(G_1, \dots, G_n)$ of the graphs G_1, \dots, G_n is the graph obtained by gluing the G_i together to form a circle, by identifying the vertices v_i and w_{i+1} , for $i \in \mathbb{Z}/n\mathbb{Z}$. It clearly depends on these choices. Define $c(G) = G^\bullet$.

Lemma 19. *Let $c(G_1, \dots, G_n)$ be any circular join of G_1, \dots, G_n . Then*

$$(173) \quad \psi_{c(G_1, \dots, G_n)} = \phi_{G_1 \cup \dots \cup G_n} .$$

Proof. There is an obvious bijection:

$$T(G) \longleftrightarrow \bigcup_{i=1}^n T(G_1) \times \dots \times T(G_i^\bullet) \times \dots \times T(G_n) ,$$

(disjoint union) where $T(\Gamma)$ denotes the set of spanning trees in a graph Γ . \square

When $n = 2$, the circular join of G_1, G_2 is a 2-vertex join of the graphs $\overline{G}_1, \overline{G}_2$.

Definition 20. Let γ, Γ be single-scale graphs where Γ is connected. We define

$$(174) \quad \Upsilon_{\gamma; \Gamma}(s) = s \psi_\gamma \phi_\Gamma + \phi_\gamma \psi_\Gamma ,$$

where s is an indeterminate which will later correspond to an external momentum.

6.3. Hopf algebras of 1-scale graphs. Let G be a fixed log-divergent graph with at most log-divergent subgraphs, and let H_G denote the finitely-generated \mathbb{Q} -Hopf algebra of graphs generated by G under the coproduct Δ (??). Its underlying vector space is spanned by all logarithmically divergent subgraphs γ of G , their cographs G/γ , and their disjoint unions. That no such cograph G/γ can have quadratic subdivergences is an easy consequence of the definitions.

Now suppose that G is single-scale, and thus carries a pair of distinguished vertices. Suppose that for every divergent subgraph Γ of G there is a choice of two distinguished connectors (Lemma ??) of Γ in G , making Γ single-scale, in such a way that for all divergent subgraphs $\gamma \subsetneq \Gamma$,

$$(175) \quad \varepsilon_\Gamma(\gamma) = 0 .$$

It follows that the cograph Γ/γ inherits the two distinguished vertices of Γ , and therefore is single-scale in a uniquely determined way. The condition (??) ensures that the two distinguished vertices of Γ are never identified in any of its cographs.

In this case, we have a Hopf algebra of 1-scale, log-divergent graphs H_G and a coassociative coproduct obtained by summing over all divergent subgraphs:

$$(176) \quad \begin{aligned} \Delta : H_G &\longrightarrow H_G \otimes_{\mathbb{Q}} H_G \\ \Gamma &\mapsto \sum_{\gamma \subsetneq \Gamma} \gamma \otimes \Gamma/\gamma . \end{aligned}$$

The existence of H_G is proved in §???. Since H_G is graded by the loop number and commutative, it automatically inherits an antipode $S : H_G \rightarrow H_G$.

Variante 21. Suppose that the edges of G are labelled by elements of some set. Then every subgraph and cograph of G , and hence every generator of H_G inherits a labelling. Note that in this case, the choice of two distinguished connectors on each divergent subgraph of G may depend on the labelling. Thus we can have distinct subgraphs γ_1 and γ_2 of G which are isomorphic after forgetting their labels, but

have different choices of distinguished vertices. In the sequel, the set of labels on the edges of a graph will always be distinct.

6.4. Feynman rules for 1-scale graphs. Let G be any graph (which is not necessarily connected), whose edges are labelled $1, \dots, N$, and let α_i denote the corresponding Schwinger parameters. Write

$$(177) \quad \Omega_G = \sum_{i=1}^N (-1)^i \alpha_i d\alpha_1 \wedge \dots \wedge \widehat{d\alpha_i} \wedge \dots \wedge d\alpha_N .$$

Definition 22. Let γ, Γ denote labelled 1-scale graphs where Γ is connected. If γ, Γ have distinct labels, then define rational forms

$$(178) \quad \omega_{\gamma \otimes \Gamma}(s) = \frac{s \phi_\Gamma}{\psi_\gamma \psi_\Gamma^2 \Upsilon_{\gamma; \Gamma}(s)} \Omega_{\gamma \cup \Gamma}$$

where we recall that ψ_γ is the product of the graph polynomials of its components. It follows from the degree formulae stated in lemmas ?? and ?? that the homogeneous degree of $\omega_{\gamma \otimes \Gamma}(s)$ with respect to the Schwinger parameters is

$$(179) \quad \deg \omega_{\gamma \otimes \Gamma}(s) = (E_\gamma - 2h_\gamma) + (E_\Gamma - 2h_\Gamma) .$$

In particular, if γ, Γ are logarithmically divergent then $\omega_{\gamma \otimes \Gamma}(s)$ is of degree 0.

Recall that the Feynman differential form of any labelled graph G is

$$(180) \quad \omega_G = \frac{\Omega_G}{\psi_G^2} .$$

It has degree $E_G - 2h_G$. Since G is not necessarily connected, ψ_G can be a product of graph polynomials. If 1 denotes the empty graph, we extend (??) and (??) by

$$(181) \quad \omega_{\gamma \otimes 1}(s) = 0 \quad \text{and} \quad \omega_{1 \otimes \Gamma}(s) = \omega_\Gamma .$$

In particular, we set $\omega_1 = 0$. It can also be convenient to set $\omega_{\gamma \otimes \Gamma}(s) = 0$ (respectively $\omega_\Gamma = 0$) if $\gamma \cup \Gamma$ (resp. Γ) have repeated labels.

7. RENORMALIZED PARAMETRIC INTEGRALS AND EXAMPLES

We state the main formula for the renormalized amplitudes of 1-scale graphs with logarithmic subdivergences, with some examples. The proofs are spread over the following sections, and the case of quadratic subdivergences is treated in §??.

7.1. Renormalised integrands. Throughout, let H denote a Hopf algebra H_G of labelled 1-scale graphs, as defined in §??, variant ??.

Definition 23. Let $\Gamma \in H$ be a labelled 1-scale graph, with edges labelled $1, \dots, N$. Denote its image under the preparation map (definition ??) by

$$(182) \quad R(\Gamma) = 1 \otimes \Gamma + \sum_{i=1}^n a_i \gamma_i \otimes \Gamma / \gamma_i \quad \in H \otimes_{\mathbb{Q}} H ,$$

where $a_i = \pm 1$ are signs⁹. Let $q \in \mathbb{R}^4$ denote the overall incoming momentum of Γ , let $\mu \in \mathbb{R}^4$ denote a fixed momentum, and set $s = q^2 / \mu^2$. We define

$$(183) \quad \omega_\Gamma^{\text{ren}}(s) = \omega_\Gamma + \sum_{i=1}^n a_i \omega_{\gamma_i \otimes \Gamma / \gamma_i}(s) ,$$

⁹These signs are consistent with the signs in the forest sums $\sum_f^\emptyset (-1)^{|f|}$.

to be the renormalised parametric Feynman integrand. For any linear combination of connected graphs $\xi = \sum_i c_i \Gamma_i \in H$, we define $\omega_\xi^{\text{ren}}(s) = \sum_i c_i \omega_{\Gamma_i}^{\text{ren}}(s)$.

The domain of integration is given by the standard coordinate simplex:

$$(184) \quad \sigma_\Gamma = \{(\alpha_1 : \dots : \alpha_N) \subset \mathbb{P}(\mathbb{R})^{N-1} : \alpha_i \geq 0\} .$$

Let Γ be a log-divergent single-scale graph with at most log-divergent subgraphs. Then by (??), $\omega_\Gamma^{\text{ren}}(s)$ is of degree 0 and we can consider the projective integral

$$(185) \quad f_\Gamma(s) = \int_{\sigma_\Gamma} \omega_\Gamma^{\text{ren}}(s) .$$

We write $f_\Gamma = f_\Gamma(1)$. The following theorem will be proved in §??.

Theorem 24. *The integral (??) converges when $s > 0$.*

The renormalization group equations amount to the following.

Theorem 25. *Write the reduced coproduct $\Delta' \Gamma = \sum_i \gamma_i \otimes \Gamma/\gamma_i$. We have*

$$(186) \quad s \frac{d}{ds} f_\Gamma(s) = \sum_i f_{\gamma_i} f_{\Gamma/\gamma_i}(s) \text{ for } s > 0 .$$

In the case when $\Gamma \in H$ is a primitive graph, i.e., $\Delta \Gamma = 1 \otimes \Gamma + \Gamma \otimes 1$, then $\omega_\Gamma^{\text{ren}}(s) = \omega_\Gamma$ and does not depend on s . In this case, (??) reduces to the residue:

$$f_\Gamma = \int_{\sigma_\Gamma} \frac{\Omega_\Gamma}{\psi_\Gamma^2} .$$

7.2. Relation with BPHZ. Let $\Gamma \in H$ be a 1-scale graph as above, and let $R(\Gamma)$ be given by (??). Then the renormalized Feynman rules are given by Eq. (??):

$$(187) \quad I(s) : H \longrightarrow \mathbb{C}$$

$$I_\Gamma(s) = \int_{\sigma_\Gamma} \left(\frac{\log s}{\psi_\Gamma^2} + \sum_{i=1}^n a_i \frac{\log \left(\frac{s \psi_{\gamma_i} \phi_{\Gamma/\gamma_i} + \phi_{\gamma_i} \psi_{\Gamma/\gamma_i}}{\psi_{\gamma_i} \phi_{\Gamma/\gamma_i} + \phi_{\gamma_i} \psi_{\Gamma/\gamma_i}} \right)}{\psi_{\gamma_i}^2 \psi_{\Gamma/\gamma_i}^2} \right) \Omega_\Gamma$$

where $s = q^2/\mu^2$, assumed to be positive. It follows immediately that $I_\Gamma(1) = 0$ and that $s I_\Gamma'(s) = f_\Gamma(s)$ by definitions (??) and (??). In particular, the convergence of (??) follows from theorem ???. Again, we extend the definition of $I_\bullet(s)$ by linearity to linear combinations of connected graphs $\xi \in H$.

7.2.1. Example 1: The case of a single subdivergence. Let Γ be a log-divergent 1-scale graph with a single log-divergent subgraph $\gamma \subset \Gamma$. Label the edges of Γ from 1 to N , and fix a 1-scale structure on γ . Fixing such a structure involves choosing any two vertices of γ which meet $\Gamma \setminus \gamma$. By definition, $\Delta(\Gamma) = 1 \otimes \Gamma + \Gamma \otimes 1 + \gamma \otimes \Gamma/\gamma$, and therefore the renormalized Feynman rules are:

$$f_\Gamma(s) = \int_{\sigma_\Gamma} \omega_\Gamma^{\text{ren}}(s) = \int_{\sigma_\Gamma} \left(\frac{1}{\psi_\Gamma^2} - \frac{s \phi_{\Gamma/\gamma}}{\psi_\gamma \psi_{\Gamma/\gamma}^2 \Upsilon_{\gamma;G/\gamma}(s)} \right) \Omega_\Gamma ,$$


where $\Upsilon_{\gamma;G/\gamma}(s) = \phi_\gamma \psi_{G/\gamma} + s \psi_\gamma \phi_{G/\gamma}$. The group equation (??) gives

$$f_\Gamma(s) = f_\Gamma + f_\gamma f_{\Gamma/\gamma} \log s$$

Thus all new information about Γ is encoded in f_Γ . We can rewrite this entirely in terms of graph polynomials using the definition of circular joins:


$$(188) \quad f_\Gamma = \int_{\sigma_\Gamma} \left(\frac{1}{\psi_\Gamma^2} - \frac{\psi_{\Gamma^\bullet/\gamma}}{\psi_{\Gamma/\gamma}^2 \psi_\gamma \psi_{c(\gamma, \Gamma/\gamma)}} \right) \Omega_\Gamma .$$

The smallest non-trivial example of such a graph Γ which is simple (i.e., does not reduce to a smaller graph by series-parallel operations) is the wheel with 3 spokes inserted into itself. The five graphs which occur in (??) are illustrated below:




It is surprising that the quantity (??) is not known for this graph, as it is the simplest possible example of a graph which requires renormalization and does not reduce trivially to smaller graphs by series-parallel operations. However, one easily checks that the parametric integration method of [?], [?] applies to this case.

7.2.2. *Example 2: overlapping subdivergences.* Consider the graph Γ below, see also Eq.(??).



It has the three overlapping log-divergent subgraphs shown below. We give them single-scale structures:



They generate a Hopf algebra of single-scale graphs H_Γ . In general, let γ_{ij} denote a 2-edge banana graph on edges i, j with its unique single-scale structure. The coproduct is $\Delta(\Gamma) = 1 \otimes \Gamma + \Gamma \otimes 1 + \sum_{i \in \{l, 34, r\}} \gamma_i \otimes \Gamma/\gamma_i$, hence

$$\Delta^{(1)}(\Gamma) = \gamma_l \otimes \gamma_{56} + \gamma_r \otimes \gamma_{12} + \gamma_{34} \otimes \gamma_{12} \cdot \gamma_{56}$$

where $\gamma_{12} \cdot \gamma_{56} = \Gamma / \gamma_{34}$ is the 1-vertex join of γ_{12} and γ_{56} whose single scale structure is given by the two outer (non-joined) vertices. It follows that

$$\Delta^{(2)}(\Gamma) = \gamma_{34} \otimes \gamma_{12} \otimes \gamma_{56} + \gamma_{34} \otimes \gamma_{56} \otimes \gamma_{12} .$$

The preparation map applied to Γ therefore gives $R(\Gamma) =$

$$1 \otimes \Gamma - (\gamma_l \otimes \gamma_{56} + \gamma_r \otimes \gamma_{12} + \gamma_{34} \otimes \gamma_{12} \cdot \gamma_{56}) + (\gamma_{34} \gamma_{12} \otimes \gamma_{56} + \gamma_{34} \gamma_{56} \otimes \gamma_{12}) ,$$

where $\gamma_{ij} \gamma_{kl}$ denotes the disjoint union $\gamma_{ij} \cup \gamma_{kl}$. Thus

$$\omega_{\Gamma}^{\text{ren}}(s) = \left[\frac{1}{\psi_{\Gamma}^2} - \frac{s \phi_{\gamma_{56}}}{\psi_{\gamma_l} \psi_{\gamma_{56}}^2 \Upsilon_{\gamma_l; \gamma_{56}}(s)} - \frac{s \phi_{\gamma_{12}}}{\psi_{\gamma_r} \psi_{\gamma_{12}}^2 \Upsilon_{\gamma_r; \gamma_{12}}(s)} - \frac{s \phi_{\gamma_{12} \cdot \gamma_{56}}}{\psi_{\gamma_{34}} \psi_{\gamma_{12} \cdot \gamma_{56}}^2 \Upsilon_{\gamma_{34}; \gamma_{12} \cdot \gamma_{56}}(s)} \right. \\ \left. + \frac{s \phi_{\gamma_{56}}}{\psi_{\gamma_{34}} \psi_{\gamma_{12}} \psi_{\gamma_{56}}^2 \Upsilon_{\gamma_{34} \gamma_{12}; \gamma_{56}}(s)} + \frac{s \phi_{\gamma_{12}}}{\psi_{\gamma_{34}} \psi_{\gamma_{56}} \psi_{\gamma_{12}}^2 \Upsilon_{\gamma_{34} \gamma_{56}; \gamma_{12}}(s)} \right] \Omega_{\Gamma}$$


where $\psi_{\gamma_{ij}} = \alpha_i + \alpha_j$, $\phi_{\gamma_{ij}} = \alpha_i \alpha_j$, and

$$\begin{aligned} \psi_{\gamma_l} &= (\alpha_1 + \alpha_2)(\alpha_3 + \alpha_4) + \alpha_3 \alpha_4 & \phi_{\gamma_l} &= \alpha_1(\alpha_2 \alpha_3 + \alpha_2 \alpha_4 + \alpha_3 \alpha_4) \\ \psi_{\gamma_r} &= (\alpha_5 + \alpha_6)(\alpha_3 + \alpha_4) + \alpha_3 \alpha_4 & \phi_{\gamma_r} &= \alpha_5(\alpha_3 \alpha_4 + \alpha_3 \alpha_6 + \alpha_4 \alpha_6) \\ \psi_{\gamma_{12} \cdot \gamma_{56}} &= (\alpha_1 + \alpha_2)(\alpha_5 + \alpha_6) & \phi_{\gamma_{12} \cdot \gamma_{56}} &= \alpha_1 \alpha_2 \alpha_5 + \alpha_1 \alpha_2 \alpha_6 + \alpha_2 \alpha_5 \alpha_6 + \alpha_1 \alpha_5 \alpha_6 \end{aligned}$$

and all remaining polynomials Υ are deduced from these by (??). In particular,

$$\begin{aligned} \Upsilon_{\gamma_l; \gamma_{56}}(s) &= s \psi_{\gamma_l} \phi_{\gamma_{56}} + \phi_{\gamma_l} \psi_{\gamma_{56}} \\ \Upsilon_{\gamma_{34}; \gamma_{12} \cdot \gamma_{56}}(s) &= s \psi_{\gamma_{34}} \phi_{\gamma_{12} \cdot \gamma_{56}} + \phi_{\gamma_{34}} \psi_{\gamma_{12} \cdot \gamma_{56}} \\ \Upsilon_{\gamma_{34} \gamma_{56}; \gamma_{12}}(s) &= s \psi_{\gamma_{34}} \psi_{\gamma_{56}} \phi_{\gamma_{12}} + \psi_{\gamma_{34}} \phi_{\gamma_{56}} \psi_{\gamma_{12}} + \phi_{\gamma_{34}} \psi_{\gamma_{56}} \psi_{\gamma_{12}} \end{aligned}$$

In the case $s = 1$, the terms $\Upsilon_{\gamma_l; \gamma_{56}}(1)$, $\Upsilon_{\gamma_r; \gamma_{12}}(1)$, $\Upsilon_{\gamma_{34}; \gamma_{12} \cdot \gamma_{56}}(1)$ and $\Upsilon_{\gamma_{34} \gamma_{56}; \gamma_{12}}(1) = \Upsilon_{\gamma_{34} \gamma_{56}; \gamma_{12}}(1)$ which occur in the denominator of $\omega_{\Gamma}^{\text{ren}}(1)$ are the graph polynomials of the following four circular join graphs, from left to right:



The renormalization group equation in this case reduces to

$$s f'_{\Gamma}(s) = f_{\gamma_l} f_{\gamma_{56}}(s) + f_{\gamma_r} f_{\gamma_{12}}(s) + f_{\gamma_{34}} f_{\gamma_{12} \cdot \gamma_{56}}(s)$$

Since the γ_{ij} are all isomorphic and primitive, and since $\gamma_r \cong \gamma_l$ this reduces to

$$s f'_{\Gamma}(s) = 2 f_{\gamma_l} f_{\gamma_{56}} + f_{\gamma_{34}} f_{\gamma_{12} \cdot \gamma_{56}}(s)$$

Applying the group equation to $\gamma_{12} \cdot \gamma_{56}$, we see that $f_{\gamma_{12} \cdot \gamma_{56}}(s) = f_{\gamma_{12} \cdot \gamma_{56}} + 2 f_{\gamma_{12}}^2 \log s$ by a similar calculation. All in all, we obtain

$$f_{\Gamma}(s) = f_{\Gamma} + (2 f_{\gamma_l} + f_{\gamma_{12} \cdot \gamma_{56}}) f_{\gamma_{12}} \log s + f_{\gamma_{12}}^3 \log^2 s .$$

8. RESIDUES

Let G be a graph with edges labelled from 1 to N . Then the Schwinger parameters $\alpha_1, \dots, \alpha_N$ define projective coordinates $(\alpha_1 : \dots : \alpha_N)$ on \mathbb{P}^{N-1} . For any strict subset of edges $I \subset E(G)$, let

$$(189) \quad L_I = \bigcap_{i \in I} \{\alpha_i = 0\} \subset \mathbb{P}^{N-1}$$

be the corresponding coordinate linear subvariety. If ω is a regular k -form on a Zariski-open subset of \mathbb{P}^{N-1} , let $v_I(\omega)$ denote the order of vanishing of ω along L_I .

Lemma 26. *Let $I \subsetneq G$ be a subgraph which is not necessarily connected. Then*

$$(190) \quad \psi_G = \psi_I \psi_{G/I} + R,$$

where R is some polynomial of degree strictly greater than $h_I = \deg \psi_I$ in the Schwinger parameters corresponding to I . It follows that $v_I(\psi_G) = h_I$.

Equation (??) is well-known, and its generalization to products of graphs is given by the following corollary. In the sequel we use the following notation: if $I \subset E(G)$, and γ is a subgraph of G , let $I_\gamma = I \cap \gamma$, and let $I_{G/\gamma} = I_G/I_\gamma$.

Corollary 27. *Let G_1, \dots, G_m be connected 1-scale graphs, let $G = \cup_i G_i$, and let $I_i \subset G_i$ be subgraphs for $i = 1, \dots, m$, where at least one $I_i \neq G_i$. Then*

$$(191) \quad v_I(\phi_G) = \sum_{i=1}^m h_{I_i} + \prod_{i=1}^m \varepsilon_{G_i}(I_i)$$

where $I = \cup_{i=1}^m I_i$. Suppose that for every $i = 1, \dots, m$, either $\varepsilon_{G_i}(I_i) = 0$ or $I_i = G_i$. Then the non-zero quotients G_i/I_i are single-scale graphs, and

$$(192) \quad \phi_G = \psi_I \phi_{G/I} + R$$

where R is of degree strictly greater than $\deg \psi_I$ in the variables of ψ_I . Likewise, for any 1-scale graphs $\gamma = \cup_i \gamma_i, \Gamma$ with γ_i, Γ connected, and $I \subset \gamma \cup \Gamma$, we have

$$(193) \quad v_I(\Upsilon_{\gamma; \Gamma}(s)) = \sum_{i=1}^m h_{I_{\gamma_i}} + \varepsilon_\gamma(I_\gamma) \varepsilon_\Gamma(I_\Gamma).$$

If $\varepsilon_{\gamma_i}(I_{\gamma_i}) = 0$ (resp. $\varepsilon_\Gamma(I_\Gamma) = 0$) whenever $\gamma_i \subsetneq I$ (resp. $\Gamma \subsetneq I$), then

$$(194) \quad \Upsilon_{\gamma; \Gamma}(s) = \psi_I \Upsilon_{\gamma/I_\gamma; \Gamma/I_\Gamma}(s) + R.$$

Proof. Use the interpretation of ϕ_G as the graph polynomial of a circular join $\psi_{c(G_1, \dots, G_m)}$. We know from (??) that for any subgraph I in a graph G , the order of vanishing of ψ_G along $\alpha_I = 0$ is equal to the number of loops in I . Equations (??) and (??) follow from computing the number of loops of I in $G = c(G_1, \dots, G_m)$, and applying (??) to G . The corresponding calculation for $\Upsilon_{\gamma; \Gamma}(s)$ follows from this case, by definition (??). \square

8.1. Orders of poles. We compute the orders of the poles of $\omega_{\gamma \otimes \Gamma}(s)$ along L_I .

Lemma 28. *Let γ, Γ denote single-scale graphs, with Γ connected, and let $I_\gamma \subset \gamma, I_\Gamma \subset \Gamma$ and $I = I_\gamma \cup I_\Gamma$. Then*

$$(195) \quad -v_I(\omega_{\gamma \otimes \Gamma}(s)) = 2h_{I_\gamma} + 2h_{I_\Gamma} - \varepsilon_{\gamma \otimes \Gamma}(I),$$

where $\varepsilon_{\gamma \otimes \Gamma} \in \{0, 1\}$ is defined by

$$(196) \quad \varepsilon_{\gamma \otimes \Gamma} = \varepsilon_\Gamma(1 - \varepsilon_\gamma).$$

Proof. By the remarks preceding (??), $v_I(\phi_\Gamma) = h_{I_\Gamma} + \varepsilon_\Gamma(I)$. Likewise it follows from corollary ?? that $v_I(\Upsilon_{\gamma; \Gamma}(s)) = h_{I_\gamma} + h_{I_\Gamma} + \varepsilon_\gamma \varepsilon_\Gamma(I)$. Summing the contributions of each term in (??) gives the formula. \square

From now on let us assume that G is log-divergent and has at most logarithmically divergent subgraphs. A *flag* in G is a nested sequence of divergent subgraphs

$$(197) \quad F : \quad \gamma_1 \subset \gamma_2 \dots \subset \gamma_n \subset G$$

where all inclusions are strict. Given such a flag F , let us write:

$$(198) \quad \gamma_F = \gamma_1 \cup \gamma_2/\gamma_1 \cup \dots \cup \gamma_n/\gamma_{n-1} \quad \text{and} \quad G/\gamma_F = G/\gamma_n .$$

If I is a subset of edges of G , write

$$(199) \quad I_{\gamma_F} = I_{\gamma_1} \cup I_{\gamma_2/\gamma_1} \cup \dots \cup I_{\gamma_n/\gamma_{n-1}} \quad \text{and} \quad I_{G/\gamma_F} = I_{G/\gamma_n} .$$

We say that I_{γ_F} is divergent if I_{γ_1} , and all $I_{\gamma_{i+1}/\gamma_i}$ are either empty or divergent.

Corollary 29. *Let $I \subsetneq E(G)$, and let F be a flag in G . The form $\omega_{\gamma_F \otimes G/\gamma_F}(s)$ has a pole along L_I of order $\leq |I|$, with equality if and only if $I_{\gamma_F}, I_{G/\gamma_F}$ are divergent, and $I_{G/\gamma_F} \subsetneq G_{\gamma_F}$ is a strict subgraph.*

Proof. From equation (??), we have

$$-v_I(\omega_{\gamma_F \otimes G/\gamma_F}(s)) = 2h_{I_{\gamma_F}} + 2h_{I_{G/\gamma_F}} - \varepsilon_{\gamma_F \otimes G/\gamma_F}(I) .$$

It follows that $-v_I(\omega_{\gamma_F \otimes G/\gamma_F}(s)) \leq |I| - \varepsilon_{\gamma_F \otimes G/\gamma_F}(I) \leq |I|$ with equality if and only if $I_{\gamma_F}, I_{G/\gamma_F}$ are divergent and $\varepsilon_{\gamma_F \otimes G/\gamma_F}(I) = 0$. But if I_{G/γ_F} is divergent and a strict subgraph of G/γ_F , $\varepsilon_{G/\gamma_F}(I_{G/\gamma_F}) = 0$ by assumption (??). By (??), this gives $\varepsilon_{\gamma_F \otimes G/\gamma_F}(I) = 0$. In the case when $I_{G/\gamma_F} = G/\gamma_F$, we have $\varepsilon_{G/\gamma_F}(I_{G/\gamma_F}) = 1$, and since $I \subset E(G)$ is strict, we must have $I_{\gamma_F} \subsetneq \gamma_F$, so $\varepsilon_{\gamma_F}(I_{\gamma_F}) = 0$ by assumption (??). Thus the case when $I_{G/\gamma_F} = G/\gamma_F$ gives rise to a pole of order $\leq |I| - 1$. \square

8.2. Blow-ups and differential forms. Let $\emptyset \neq I \subset E(G)$ be a strict subset of edges of G , whose edges are labelled $1, \dots, N$. Consider the blow-up

$$\pi_I : P_I \longrightarrow \mathbb{P}^{N-1}$$

of \mathbb{P}^{N-1} along L_I . We denote the exceptional divisor by \mathcal{E}_I , which is isomorphic to $\mathbb{P}^{|I|-1} \times \mathbb{P}^{|I^c|-1}$, where I^c is the complement of I in $E(G)$.

Lemma 30. *Let ω be a differential form of the shape $(\prod_{k=1}^m P_k^{r_k}) \Omega_G$ where $r_k \in \mathbb{Z}$, and each P_k is a homogeneous polynomial in the α_i of the form*

$$P_k = A_k(\{\alpha_i\}_{i \in I}) B_k(\{\alpha_j\}_{j \in I^c}) + R_k$$

where A_k, B_k, R_k are homogeneous and the degree of R_k in the variables $\alpha_i, i \in I$ is strictly greater than $\deg A_k$. Then the order of the pole of $\pi_I^* \omega$ along \mathcal{E}_I is

$$(200) \quad 1 - \sum_{k=1}^m r_k \deg(A_k) - |I| = 1 - v_I(\omega) - |I|$$

where $v_I(\omega)$ is the order of vanishing of ω along L_I . The pole is simple if and only if $-v_I(\omega) = |I|$, in which case the residue of $\pi_I^* \omega$ along \mathcal{E}_I is given by

$$(201) \quad \text{Res}_{\mathcal{E}_I} \pi_I^* \omega = \prod_{k=1}^m A_k(\alpha_i, i \in I) \Omega_I \otimes \prod_{k=1}^m B_k(\alpha_j, j \in I^c) \Omega_{I^c}$$

Proof. Let $I = \{i_1, \dots, i_a\}$, and $I^c = \{j_1, \dots, j_b\}$. On the affine chart of P defined by $\alpha_{j_b} = 1$, take local affine coordinates $\alpha_{j_1}, \dots, \alpha_{j_{b-1}}, z = \alpha_{i_1}$, and

$$\beta_{i_2} = \frac{\alpha_{i_2}}{\alpha_{i_1}}, \dots, \beta_{i_a} = \frac{\alpha_{i_a}}{\alpha_{i_1}}.$$

In these coordinates, \mathcal{E}_I is given by $z = 0$. On this chart,

$$\Omega_N = \pm z^{a-1} dz \wedge d\beta_{i_2} \wedge \dots \wedge d\beta_{i_a} \wedge d\alpha_{j_1} \wedge \dots \wedge d\alpha_{j_{b-1}}.$$

Equation (??) follows by substituting the new variables into ω and computing the order of the pole in z . Equation (??) follows on taking the residue at $z = 0$. \square

8.3. Calculation of the residues. Let G, π_I be as above.

Proposition 31. *Let F be a flag of divergent subgraphs in G , and let I be a strict subset of edges in G . The form $\pi_I^* \omega_{\gamma_F \otimes G / \gamma_F}(s)$ has a simple pole along the exceptional divisor \mathcal{E}_I if and only if $I_{\gamma_F}, I_{G/\gamma_F}$ are divergent, and $I_{G/\gamma_F} \subsetneq G/\gamma_F$. The residue is*

$$(202) \quad \text{Res}_{\mathcal{E}_I} \pi_I^* \omega_{\gamma_F \otimes G / \gamma_F}(s) = \omega_{I_{\gamma_F} \cup I_{G/\gamma_F}} \otimes \omega_{\gamma_F / I \otimes G / (\gamma_F \cup I)}(s).$$

Proof. Apply lemma ?? to the definition (??) of $\omega_{\gamma_F \otimes G / \gamma_F}(s)$. By (??) and (??), to lowest order terms in the I parameters:

$$\begin{aligned} \phi_{G/\gamma_F} &= \psi_{I_{G/\gamma_F}} \phi_{G/(\gamma_F \cup I)}, \\ \Upsilon_{\gamma_F; G/\gamma_F}(s) &= \psi_{I_{\gamma_F}} \psi_{I_{G/\gamma_F}} \Upsilon_{\gamma_F/I; G/(\gamma_F \cup I)}(s). \end{aligned}$$

By lemma ?? and separating out terms we deduce that $\text{Res}_{\mathcal{E}_I} \pi_I^* \omega_{\gamma_F \otimes G / \gamma_F}(s)$ is

$$\frac{\psi_{I_{G/\gamma_F}}}{\psi_{I_{G/\gamma_F}}^2 \psi_{I_{\gamma_F}}^2 \psi_{I_{G/\gamma_F}}} \Omega_I \otimes \frac{\phi_{G/(\gamma_F \cup I)}}{\psi_{\gamma_F/I} \psi_{G/(\gamma_F \cup I)}^2 \Upsilon_{\gamma_F/I; G/(\gamma_F \cup I)}(s)} \Omega_{G/I}$$

which is exactly $\omega_{I_{\gamma_F} \cup I_{G/\gamma_F}} \otimes \omega_{\gamma_F / I \otimes G / (\gamma_F \cup I)}(s)$. \square

For any meromorphic algebraic form ω on \mathbb{P}^{N-1} , define the total residue to be

$$(203) \quad \text{Res} \omega = \bigoplus_I \text{Res}_{\mathcal{E}_I} \pi_I^* \omega,$$

where the sum is over all strict subsets $I \subsetneq \{1, \dots, N\}$.

Proposition 32. *Let G be a connected log-divergent graph with at most log-divergent subgraphs. For any flag F of divergent subgraphs of G , the total residue is*

$$\text{Res} \omega_{\gamma_F \otimes G / \gamma_F}(s) = (\omega^1 \otimes \omega^{23}) \circ \mu_{13}(\Delta \otimes \Delta)(\gamma_F \otimes G / \gamma_F),$$

where $(\omega^1 \otimes \omega^{23})(x \otimes y \otimes z) = \omega_x \otimes \omega_{y \otimes z}(s)$.

Proof. By the definition of the total residue, $\text{Res} \omega_{\gamma_F \otimes G / \gamma_F}(s)$ is equal to

$$\sum_I \text{Res}_{\mathcal{E}_I} \pi_I^* \omega_{\gamma_F \otimes G / \gamma_F}(s) = \sum_{I_{\gamma_F}, I_{G/\gamma_F} \text{ div.}} \omega_{I_{\gamma_F} \cup I_{G/\gamma_F}} \otimes \omega_{\gamma_F / I_{\gamma_F} \otimes G / (\gamma_F \cup I)}(s),$$

by proposition ??, since the right-hand side vanishes when $I_{G/\gamma_F} = G/\gamma_F$. This can be rewritten

$$\sum_{a \subset \gamma_F \text{ div.}} \sum_{b \subset G/\gamma_F \text{ div.}} (\omega^1 \otimes \omega^{23}) \circ \mu_{13}(a \otimes \gamma_F / a \otimes b \otimes G/\gamma_F / b)$$

which is $(\omega^1 \otimes \omega^{23}) \circ \mu_{13}(\Delta(\gamma_F) \otimes \Delta(G/\gamma_F))$. \square

9. CANCELLATION OF SIMPLE POLES

The proof of convergence of the renormalized Feynman integral follows from a few abstract Hopf-algebra theoretic properties of the residues.

9.1. The set-up. Let H be a commutative, graded Hopf algebra over \mathbb{Q} , and let \mathfrak{D} be a vector space over k , a field of characteristic 0. We think of \mathfrak{D} as some space of differential forms. Suppose we are given a linear map $\omega : H \rightarrow \mathfrak{D}$ such that $\omega(1) = 0$, and a map (the ‘total residue’)

$$\text{Res} : \mathfrak{D} \rightarrow \mathfrak{D} \otimes_k \mathfrak{D}$$

which is related to the coproduct $\Delta : H \rightarrow H \otimes_{\mathbb{Q}} H$ by:

$$(204) \quad \text{Res} \omega = (\omega \otimes \omega) \circ \Delta .$$

In particular, this means that the map Res is coassociative. Now suppose that there exists another linear map

$$\omega^{(2)} : H \otimes_{\mathbb{Q}} H \longrightarrow \mathfrak{D}$$

such that, for all $\xi \in H$,

$$(205) \quad \begin{aligned} \text{(1)} : \quad & \omega^{(2)}(\xi \otimes 1) = 0 \\ \text{(2)} : \quad & \omega^{(2)}(1 \otimes \xi) = \omega(\xi) \\ \text{(3)} : \quad & \text{Res} \omega^{(2)} = (\omega \otimes \omega^{(2)}) \circ \mu_{13}(\Delta \otimes \Delta) . \end{aligned}$$

Definition 33. In this situation, we define the renormalized map

$$(206) \quad \begin{aligned} \omega^{\text{ren}} : H & \longrightarrow \mathfrak{D} \\ \omega^{\text{ren}} & = \omega^{(2)} \circ R \end{aligned}$$

Since we have by definition $R(x) = 1 \otimes x +$ terms of lower order in the coradical filtration, then by assumption **(2)**, we can view $\omega^{\text{ren}} = \omega +$ lower order terms. The following proposition is the main result for the renormalization of logarithmic singularities.

Proposition 34. *With the assumptions (??), $\text{Res} \omega^{\text{ren}} = 0$.*

Proof. Using properties **(1)** – **(3)** above, and theorem ??, we have

$$\begin{aligned} \text{Res} \omega^{\text{ren}} = \text{Res} \omega^{(2)} \circ R & \stackrel{\text{(3)}}{=} (\omega \otimes \omega^{(2)}) \circ \mu_{13}(\Delta \otimes \Delta) \circ R \\ & \stackrel{\text{(1)}}{=} (\omega \otimes \omega^{(2)}) \circ \mu_{13}(\Delta \otimes (\Delta - id \otimes 1)) \circ R \\ & = (\omega \otimes \omega^{(2)}) \circ (1 \otimes R) . \end{aligned}$$

The equality on the third line follows from equation (??). The final expression vanishes by the property that $\omega(1) = 0$, and this follows from **(1)** and **(2)**. \square

Remark 35. *Observe that **(3)** implies (??). For $x \in H$, we have*

$$\text{Res} \omega(x) \stackrel{\text{(2)}}{=} \text{Res} \omega^{(2)}(1 \otimes x) \stackrel{\text{(3)}}{=} (\omega \otimes \omega^{(2)}) \circ \mu_{13}(\Delta \otimes \Delta)(1 \otimes x)$$

If we write $\Delta x = \sum_{(x)} x^{(1)} \otimes x^{(2)}$ using Sweedler’s notation (??), this reduces to

$$\sum_{(x)} (\omega \otimes \omega^{(2)}) \circ (x^{(1)} \otimes 1 \otimes x^{(2)}) \stackrel{\text{(2)}}{=} \sum_{(x)} \omega(x^{(1)}) \otimes \omega(x^{(2)}) = (\omega \otimes \omega) \circ \Delta(x) .$$

10. BLOW-UPS AND MIXED HODGE STRUCTURES

10.1. Hypersurfaces. Fix a 1-scale graph G with N edges. For any subgraph or quotient graph γ of G , the graph hypersurface $X_\gamma = V(\psi_\gamma) \subset \mathbb{P}^{N-1}$ is defined to be the zero locus of the graph polynomial ψ_γ (using convention (??)). If F is a flag of divergent subgraphs in G , define a family of hypersurfaces

$$X_{\gamma_F \otimes G / \gamma_F}^s = V(\Upsilon_{\gamma_F \otimes G / \gamma_F}(s)) \subset \mathbb{P}^{N-1},$$

over \mathbb{P}^1 , with coordinate s . If G has at most logarithmically divergent subgraphs, define $X_{G,s}^{tot} \subset \mathbb{P}^{N-1}$ to be the union of the graph hypersurfaces X_{γ_F} , X_{G/γ_F} , and $X_{\gamma_F \otimes G / \gamma_F}^s$, as F ranges over the set of flags of divergent subgraphs of G . By definition (??), $\omega_G^{\text{ren}}(s) \in \Omega^{N-1}(\mathbb{P}^{N-1} \setminus X_{G,s}^{tot})$. Recall from (??) that σ_G is the standard coordinate simplex, and write $D_I = L_I \cap \sigma_G$ for all $I \subset E(G)$. It is known by [?], Proposition 3.1 and Lemma 7.1, that

$$(207) \quad X_G \cap D_I \neq \emptyset \iff D_I \subset X_G \iff h_I > 0.$$

There is an obvious generalization for the hypersurfaces $X_{\gamma_F \otimes G / \gamma_F}^s$.

Lemma 36. *Let G, F be as above, and let $s > 0$. The following are equivalent:*

- (1) $X_{\gamma_F \otimes G / \gamma_F}^s \cap D_I \neq \emptyset$.
- (2) $D_I \subset X_{\gamma_F \otimes G / \gamma_F}^s$.
- (3) $h_{I_{\gamma_F}} + h_{I_{G/\gamma_F}} > 0$ or $\varepsilon_{\gamma_F}(I) = \varepsilon_{G/\gamma_F}(I) = 1$.
- (4) *The subgraph of $c(\gamma_F, G/\gamma_F)$ defined by the edges I contains a loop.*

In particular, $X_{\gamma_F} \cap D_I \subset X_{\gamma_F \otimes G / \gamma_F}^s \cap D_I$ and $X_{G/\gamma_F} \cap D_I \subset X_{\gamma_F \otimes G / \gamma_F}^s \cap D_I$.

Proof. Since $\Upsilon_{\gamma_F; G/\gamma_F}(s)$ has positive coefficients, $X_{\gamma_F \otimes G / \gamma_F}^s$ meets D_I if and only if $L_I \subset X_{\gamma_F \otimes G / \gamma_F}^s$. By (??), this occurs if and only if $h_{I_{\gamma_F}} + h_{I_{G/\gamma_F}} + \varepsilon_{\gamma_F}(I)\varepsilon_{G/\gamma_F}(I)$ is positive, which is in turn equivalent to (4). So (1)-(4) are equivalent. The last part is obvious by (3) and (??). \square

Thus the intersections of $X_{G,s}^{tot}$ with σ_G are contained in the union of the intersections of each hypersurface $X_{\gamma \otimes G / \gamma}^s$ with σ_G . Hereafter, let $s > 0$.

10.2. The blowup. Define the following set of subsets of edges of G :

$$G^{div} = \{I \subsetneq E(G) : I \text{ minimal such that for some flag } F \text{ of divergent subgraphs, } I \subset c(\gamma_F, G/\gamma_F) \text{ contains a loop}\}$$

The set $\{L_I : I \in G^{div}\}$ is therefore the set of maximal linear coordinate spaces whose real points are contained in $X_{G,s}^{tot} \cap \sigma_G$. Define \mathcal{F}_G to be the set of all intersections $L_{I_1 \cup \dots \cup I_k}$, with $I_j \in G^{div}$, of such coordinate spaces. Following standard practice ([?], §7), one blows up the elements in \mathcal{F}_G in strictly increasing order of codimension. One knows that the space P_G obtained in this way is well-defined. If $\pi : P_G \rightarrow \mathbb{P}^{N-1}$ denotes the blow-up, let Y_G^{tot} be the strict transform of X_G^{tot} and let B be the total inverse image of the linear spaces L_I under π . Then B is a union of the strict transforms of the coordinate hyperplanes $\alpha_i = 0$ with exceptional components \mathcal{E}_I for each $I \in G^{div}$, where \mathcal{E}_I is the strict transform of L_I .

Proposition 37. *The divisor B is normal crossing, and the strict transform of σ_G in P_G does not meet Y_G^{tot} . No non-empty intersection of the components of B is contained in Y_G^{tot} .*

Proof. This can be proved in an identical manner to [?], Proposition 7.3 (ii), (iii) (only part (i) of loc. cit. requires the assumption that the graph be primitive). \square

Example 38. Let G be the graph γ_l in §??. It has one divergent subgraph $\gamma = \{3, 4\}$. The circular join of G/γ and γ is the banana graph with four edges, with graph polynomial $\alpha_1\alpha_2\alpha_3 + \alpha_1\alpha_2\alpha_4 + \alpha_1\alpha_3\alpha_4 + \alpha_2\alpha_3\alpha_4$, which meets σ_G along all D_I for any $I \subset \{1, 2, 3, 4\}$, with $|I| \geq 2$. It follows that $G^{div} = \{\{i, j\}, 1 \leq i < j \leq 4\}$, and \mathcal{F}_G consists of all divisors L_I where $|I| \geq 2$.

Remark 39. *In general, one could simply blow-up all the faces of codimension ≥ 2 for good measure. In this case the space P_G admits an action of the symmetric group on N letters which permutes the coordinates, and B has the structure of a permutohedron with $N!$ vertices. In particular, there are canonical coordinates on P_G in the neighbourhood of each vertex indexed by a $\tau \in \Sigma_N$, given by*

$$\alpha_1^\tau = \alpha_{\tau(1)}, \quad \alpha_2^\tau = \frac{\alpha_{\tau(2)}}{\alpha_{\tau(1)}}, \quad \dots, \quad \alpha_N^\tau = \frac{\alpha_{\tau(N)}}{\alpha_{\tau(N-1)}}.$$

These are precisely the local coordinates in the sector decomposition of K . Hepp.

10.3. The renormalized differential form. We now apply the formalism of §??. Let G be a labelled single-scale overall log-divergent graph with at most log-divergent subgraphs and let $H = H_G$ be the Hopf algebra obtained by fixing single-scale structures on its sub and co-graphs (§??). For every non-empty set of edges $I \subset E(G)$, let \mathbb{P}^I denote the projective space whose coordinates are the Schwinger parameters of I , and let $\mathfrak{D}^I \subset \Omega_{\mathbb{Q}(\alpha_i; i \in I)/\mathbb{Q}}^{|I|-1}$ denote the subspace of regular forms of homogeneous degree 0 on a Zariski open subset of \mathbb{P}^I which are defined over \mathbb{Q} . Let

$$(208) \quad \mathfrak{D} = \bigoplus_{\emptyset \neq I \subset E(G)} \mathfrak{D}^I$$

As in definition (??), set

$$(209) \quad \begin{aligned} \omega^{(2)} : H_G \otimes_{\mathbb{Q}} H_G &\rightarrow \mathfrak{D}^{E(G)} \subset \mathfrak{D} \\ \gamma \otimes \Gamma &\mapsto \omega_{\gamma \otimes \Gamma}(s). \end{aligned}$$

and define $\omega : H_G \rightarrow \mathfrak{D}$ by $\omega(G) = \omega^{(2)}(1 \otimes G) = \Omega_G \psi_G^{-2}$. The total residue $\text{Res} : \mathfrak{D} \rightarrow \mathfrak{D} \otimes_{\mathbb{Q}} \mathfrak{D}$ is given by (??), i.e., the sum of the residues along all exceptional divisors \mathcal{E}_I in the blow-ups of $\mathbb{P}^{E(G)}$ along L_I . Let

$$\pi : P_G \rightarrow \mathbb{P}^{E(G)}$$

denote the blow-up constructed in §??.

Theorem 40. *If $s > 0$, the form $\pi^* \omega_G^{\text{ren}}(s)$ has no poles along B .*

Proof. It is enough (by, for example, the proof of proposition 7.3 in [?]) to show that $\pi_I^* \omega_G^{\text{ren}}(s)$ has no poles along \mathcal{E}_I , where $\pi_I : P^I \rightarrow \mathbb{P}^{E(G)}$ is the blow-up of a single coordinate hyperplane L_I . By proposition ??, it has at most simple poles. The residue is zero by proposition ??, since properties (1) and (2) hold by (??), and property (3) holds by proposition ?? \square

Corollary 41. *If $s > 0$, the renormalized Feynman integral converges:*

$$f_G(s) = \int_{\sigma_G} \omega_G^{\text{ren}}(s) < \infty.$$

Proof. Pull back the integral to P_G . Thus $f_G(s)$ is the integral of $\pi^*\omega^{\text{ren}}(s)$ over the strict transform of σ_G , a compact polytope whose boundary is contained in B . By the previous theorem, the poles of $\pi^*\omega^{\text{ren}}(s)$ are contained in $Y_{G,s}^{\text{tot}}$, which does not meet the new domain of integration by proposition ???. Thus $f_G(s)$ is the integral of a continuous function on a compact domain, and is therefore bounded. \square

10.4. The tangent mixed Hodge structure. We can therefore define the *tangent mixed Hodge structure* of G as follows. Writing Y_G^{tot} for $Y_{G,1}^{\text{tot}}$, let

$$(210) \quad \text{Mot}(G) = H^{N-1}(P_G \setminus Y_G^{\text{tot}}, B \setminus B \cap Y_G^{\text{tot}}) .$$

The renormalized form $\omega_G^{\text{ren}}(1)$ defines a cohomology class

$$[\pi^*(\omega_G^{\text{ren}}(1))] \in H_{DR}^{N-1}(P_G \setminus Y_G^{\text{tot}}, B \setminus B \cap Y_G^{\text{tot}}) ,$$

and the strict transform of the simplex Δ_N defines a relative homology class

$$[\pi^{-1}(\Delta_N)] \in W^0 H_{B,N-1}(P_G \setminus Y_G^{\text{tot}}, B \setminus B \cap Y_G^{\text{tot}}) .$$

Corollary 42. *The lowest log term f_G of the renormalized Feynman integral of G defines a period of the tangent mixed Hodge structure $\text{Mot}(G)$.*

It follows from the renormalization group equations that $f_G(s)$ is a polynomial in $\log s$, whose coefficients are (products of) periods of mixed Hodge structures (??).

11. THE RENORMALIZATION GROUP

We prove that the Feynman rules respect the renormalization group structure.

11.1. Renormalization group equations. It is convenient to define a new set of differential forms as follows. Let λ denote any parameter, and set

$$(211) \quad \nu_G(\lambda) = \frac{\phi_G}{\psi_G} \frac{\lambda}{(\psi_G \lambda + \phi_G)^2} \prod_{e \in E(G)} d\alpha_e ,$$

$$(212) \quad \nu_{\gamma \otimes \Gamma}(\lambda) = \frac{\phi_\Gamma}{\psi_\Gamma} \frac{\lambda}{(\psi_\Gamma \lambda + \phi_\Gamma)^2} \prod_{e \in E(\gamma) \cup E(\Gamma)} d\alpha_e ,$$

where G and γ, Γ are labelled single-scale graphs with disjoint labels. Denote the corresponding renormalized form by $\nu_G^{\text{ren}}(\lambda) = \nu(\lambda) \circ R(G)$. The following lemma enables us to lift the domain of integration to one dimension higher.

Lemma 43. *Let G be single-scale as above. For all $\lambda > 0$ we have :*

$$(213) \quad \int_{\sigma_G} \omega_G^{\text{ren}}(1) = \int_{[0,\infty]^{E(G)}} \nu_G^{\text{ren}}(\lambda) .$$

Proof. We first show that the right-hand side is convergent. For this, it is a simple matter to check that $\nu_{1 \otimes G}(\lambda) = \nu_G(\lambda)$, and $\nu_{G \otimes 1}(\lambda) = 0$. By a similar calculation to proposition ??? (with the small difference that we work in affine rather than projective space) we find that the total residue satisfies

$$\text{Res } \nu(\lambda) = (\nu(\lambda) \otimes \nu(\lambda)) \circ (\Delta \otimes \Delta) ,$$

and therefore by the general set-up of ???, the integral is convergent. The proof of the lemma uses the fact that for any $A, B > 0$, we have

$$(214) \quad \frac{1}{AB} = \int_0^\infty \frac{d\lambda}{(A\lambda + B)^2} .$$

Now let $\sigma_G^\varepsilon = \{(\alpha_1 : \dots : \alpha_{|E(G)|}) \in \mathbb{P}^{|E(G)|-1} : \alpha_i \geq |\varepsilon|\}$, and write the left hand side of (??) as a limit

$$\int_{\sigma_G} \omega_G^{\text{ren}}(1) = \lim_{\varepsilon \rightarrow 0^+} \int_{\sigma_G^\varepsilon} \omega_G^{\text{ren}}(1).$$

Writing the preparation map $R(G) = 1 \otimes G + \sum_\gamma a_\gamma \gamma \otimes G/\gamma$, we obtain

$$\int_{\sigma_G^\varepsilon} \omega_G^{\text{ren}}(1) = \int_{\sigma_G^\varepsilon} \frac{\Omega_G}{\psi_G^2} + \sum_\gamma a_\gamma \int_{\sigma_G^\varepsilon} \frac{\phi_{G/\gamma} \Omega_G}{\psi_\gamma \psi_{G/\gamma}^2 \phi_{\gamma \cup G/\gamma}}.$$

Since σ_G^ε does not meet the coordinate hyperplanes, the graph polynomials are strictly positive on σ_G^ε and we can apply (??) with $(A, B) = (\psi_G, \phi_G)$ to the first factor and with $(A, B) = (\psi_{\gamma \cup G/\gamma}, \phi_{\gamma \cup G/\gamma})$ to the factors in the sum. This gives

$$\int_{\sigma_G^\varepsilon} \int_0^\infty \frac{\phi_G}{\psi_G} \frac{\Omega_G d\lambda}{(\psi_G \lambda + \phi_G)^2} + \sum_\gamma a_\gamma \int_{\sigma_G^\varepsilon} \int_0^\infty \frac{\phi_{G/\gamma}}{\psi_{G/\gamma}} \frac{\Omega_G d\lambda}{(\psi_{\gamma \cup G/\gamma} \lambda + \phi_{\gamma \cup G/\gamma})^2}.$$

We can take the limit as $\varepsilon \rightarrow 0$:

$$\int_{\sigma_G} \int_0^\infty \left(\frac{\phi_G}{\psi_G} \frac{1}{(\psi_G \lambda + \phi_G)^2} + \sum_\gamma a_\gamma \frac{\phi_{G/\gamma}}{\psi_{G/\gamma}} \frac{1}{(\psi_{\gamma \cup G/\gamma} \lambda + \phi_{\gamma \cup G/\gamma})^2} \right) \Omega_G d\lambda$$

since the right-hand side of the integrand (viewed as an integral on $\mathbb{P}^{|E(G)|-1} \times \mathbb{A}^1$) is convergent, by the usual arguments. The integrand defines a projective integral on $\mathbb{P}^{|E(G)|}$ with projective coordinates $(\alpha_1 : \dots : \alpha_{E(G)} : \lambda)$ if we replace $\Omega_G d\lambda$ with $\Omega_G d\lambda + \lambda \prod_i d\alpha_i$. Restricting to any hyperplane $\lambda = \text{constant}$ gives an affine integral which is precisely the right-hand side of (??). \square

Proposition 44. *Let G_1, G_2 denote single-scale graphs with at most logarithmic subdivergences, and denote their images under the preparation map R by*

$$R(G_i) = \sum_{\gamma_i} a_{\gamma_i} \gamma_i \otimes G_i/\gamma_i,$$

where $a_{\gamma_i} = \pm 1$, the sum is over all flags of (possibly empty) divergent subgraphs $\gamma_i \subset G_i$ (??) and $i = 1, 2$. If G_1, G_2 have distinct labels and $s > 0$ then we have

$$(215) \quad \int_{\sigma_{G_1}} \omega_{G_1}^{\text{ren}}(1) \times \int_{\sigma_{G_2}} \omega_{G_2}^{\text{ren}}(s) =$$

$$(216) \quad \int_{\sigma_{G_1 \cup G_2}} \sum_{\gamma_i \subset G_i} a_{\gamma_1} a_{\gamma_2} \frac{\phi_{G_1/\gamma_1} \phi_{G_2/\gamma_2}}{\psi_{G_1/\gamma_1} \psi_{G_2/\gamma_2}} \frac{s \Omega_{G_1 \cup G_2}}{(\Upsilon_{\gamma_1 \cup \gamma_2; G_2/\gamma_2}(s))^2},$$

where all integrals are convergent.

Proof. The two integrals in the first line of (??) are convergent by theorem ???. To check the convergence of (??), write

$$(217) \quad \omega_{\gamma_1 \otimes \gamma_2 \otimes \gamma_3 \otimes \gamma_4}^{(4)}(s) = \frac{\phi_{\gamma_2} \phi_{\gamma_4}}{\psi_{\gamma_2} \psi_{\gamma_4}} \frac{s \Omega_{\gamma_1 \cup \dots \cup \gamma_4}}{\Upsilon_{\gamma_1 \cup \gamma_2 \cup \gamma_3; \gamma_4}^2(s)}$$

and verify, by a similar computation to lemma ?? that $\omega^{(4)}$ has at most simple poles along exceptional divisors \mathcal{E}_I and that its total residue is

$$\text{Res } \omega^{(4)}(s) = (\omega \otimes \omega^{(4)}) \circ \mu_{1357} \circ (\Delta \otimes \Delta \otimes \Delta \otimes \Delta).$$

where $\mu_{1357}(\gamma_1 \otimes \dots \otimes \gamma_8) = \gamma_1 \gamma_3 \gamma_5 \gamma_7 \otimes \gamma_2 \otimes \gamma_4 \otimes \gamma_6 \otimes \gamma_8$. The integrand in (??) is given by $\omega^{(4)}(s) \circ (R \otimes R)(G_1 \otimes G_2)$. We have

$$\begin{aligned} \text{Res } \omega^{(4)} \circ (R \otimes R) &= (\omega \otimes \omega^{(4)}) \circ \mu_{1357} \circ (\Delta \otimes \Delta \otimes \Delta \otimes \Delta) \circ (R \otimes R) \\ &= (\omega \otimes \omega^{(4)}) \circ \mu_{14}((\mu_{13} \circ (\Delta \otimes \Delta) \circ R) \otimes (\mu_{13} \circ (\Delta \otimes \Delta) \circ R)) \\ &= (\omega \otimes \omega^{(4)}) \circ \mu_{14}(1 \otimes R \otimes 1 \otimes R) \\ &= \omega(1) \otimes \omega^{(4)}(R \otimes R) = 0 \end{aligned}$$

The third equality follows from the fact that $\omega^{(4)}(\gamma_1 \otimes \gamma_2 \otimes \gamma_3 \otimes 1) = 0$ and theorem ??, and the vanishing of the last line follows from $\omega(1) = 0$. Using the positivity of graph polynomials (since $s > 0$), we prove as in corollary ?? that (??) converges. For the proof of the main identity, write

$$\int_{\sigma_{G_2}} \omega_{G_2}^{\text{ren}}(s) = \lim_{\varepsilon \rightarrow 0} \left(\int_{\sigma_{G_2}^\varepsilon} \omega_{G_2} + \sum_{\emptyset \neq \gamma_2 \subset G_2} a_{\gamma_2} \int_{\sigma_{G_2}^\varepsilon} \omega_{\gamma_2 \otimes G_2 / \gamma_2}(s) \right),$$

where $\sigma_{G_2}^\varepsilon = \{(\alpha_1 : \dots : \alpha_{|E(G_2)|}) \in \mathbb{P}^{|E(G_2)|-1} : \alpha_i \geq |\varepsilon|\}$. Multiply this expression through by

$$\int_{\sigma_{G_1}} \omega_{G_1}^{\text{ren}}(1) = \int_{[0, \infty]^{E(G_1)}} \nu_{G_1}^{\text{ren}}(\lambda),$$

which holds by lemma ?? for any $\lambda > 0$. Thus, for any choice of λ_{γ_2} 's, we have

$$(??) = \lim_{\varepsilon \rightarrow 0} \left(\int_{X_\varepsilon} \nu_{G_1}^{\text{ren}}(\lambda_{G_2}) \wedge \omega_{G_2} + \sum_{\gamma_2 \subset G_2} a_{\gamma_2} \int_{X_\varepsilon} \nu_{G_1}^{\text{ren}}(\lambda_{\gamma_2}) \wedge \omega_{\gamma_2 \otimes G_2 / \gamma_2}(s) \right),$$

where the domain of integration $X_\varepsilon = [0, \infty]^{E(G_1)} \times \sigma_{G_2}^\varepsilon$. Now substitute

$$\lambda_{G_2} = \frac{s \phi_{G_2}}{\psi_{G_2}} \quad \text{and} \quad \lambda_{\gamma_2} = \frac{\Upsilon_{\gamma_2; G_2 / \gamma_2}(s)}{\psi_{\gamma_2 \cup G_2 / \gamma_2}}.$$

Directly from the definitions, one verifies that

$$\nu_{\gamma_1 \otimes G_1 / \gamma_1}(\lambda_{\gamma_2}) \wedge \omega_{\gamma_2 \otimes G_2 / \gamma_2}(s) = \frac{\phi_{G_1 / \gamma_1} \phi_{G_2 / \gamma_2} s \prod_{e \in E(G_1)} d\alpha_e \wedge \Omega_{G_2}}{\psi_{G_1 / \gamma_1} \psi_{G_2 / \gamma_2} (\Upsilon_{\gamma_1 \cup \gamma_2; G_2 / \gamma_2}(s))^2}.$$

By a similar calculation involving $\nu_{G_1}(\lambda)$ and so on, we conclude that

$$(??) = \lim_{\varepsilon \rightarrow 0} \left(\int_{X_\varepsilon} \sum_{\gamma_i \subset G_i} a_{\gamma_1} a_{\gamma_2} \frac{\phi_{G_1 / \gamma_1} \phi_{G_2 / \gamma_2} s \prod_{e \in E(G_1)} d\alpha_e \wedge \Omega_{G_2}}{\psi_{G_1 / \gamma_1} \psi_{G_2 / \gamma_2} (\Upsilon_{\gamma_1 \cup \gamma_2; G_2 / \gamma_2}(s))^2} \right).$$

Projectivising the integral on the right-hand side gives (??). Therefore the limit $\varepsilon = 0$ is finite and equal to (??). \square

Definition 45. Let γ, Γ be labelled single-scale graphs with disjoint labels. Let

$$(218) \quad \bar{\omega}_{\gamma \otimes \Gamma}(s) = s \frac{\partial}{\partial s} \omega_{\gamma \otimes \Gamma}(s) = \frac{\phi_\gamma \phi_\Gamma}{\psi_\gamma \psi_\Gamma} \frac{s}{(s \psi_\gamma \phi_\Gamma + \phi_\gamma \psi_\Gamma)^2}.$$

In particular, $\bar{\omega}_{1 \otimes \Gamma}(s) = 0$. The following lemma is immediate.

Lemma 46. Let $\gamma_1, \dots, \gamma_n$ be connected 1-scale graphs, and $\gamma = \cup_i \gamma_i$. Then

$$(219) \quad \frac{\phi_\gamma}{\psi_\gamma} = \sum_{i=1}^n \frac{\phi_{\gamma_i}}{\psi_{\gamma_i}}.$$

Proposition 47. *Let Γ be a connected single-scale graph with at most logarithmic subdivergences as above, and write $\Delta'\Gamma = \sum_{\gamma \subset \Gamma} \gamma \otimes \Gamma/\gamma$. Then*

$$(220) \quad \int_{\sigma_{\Gamma}} \bar{\omega}_{\Gamma}^{\text{ren}}(s) = \sum_{\gamma \subset \Gamma} \int_{\sigma_{\gamma}} \omega_{\gamma}^{\text{ren}}(1) \times \int_{\sigma_{\Gamma/\gamma}} \omega_{\Gamma/\gamma}^{\text{ren}}(s) .$$

Proof. If $a_1, \dots, a_n \in H$ are graphs, then lemma ?? implies that

$$(221) \quad \sum_{1 \leq i \leq n} \omega_{a_1 \dots a_{i-1} \otimes a_i \otimes a_{i+1} \dots a_n \otimes b}^{(4)}(s) = \bar{\omega}_{a_1 \dots a_n \otimes b}(s)$$

where empty products are defined to be 1, e.g., $\omega_{1 \otimes a_1 \otimes 1 \otimes b}^{(4)} = \bar{\omega}_{a_1 \otimes b}(s)$. Viewed as an equality of elements in $\text{Hom}(H \otimes_{\mathbb{Q}} H, \mathbb{R})$, proposition ?? states that:

$$\mu \circ (f_{G_1}(1) \otimes f_{G_2}(s)) = \int_{\sigma_{G_1 \cup G_2}} \omega^{(4)}(s) \circ (R \otimes R) \circ (G_1 \otimes G_2) .$$

Apply both sides to $\Delta'(\Gamma) \in H \otimes_{\mathbb{Q}} H$. The left-hand side equates to the right-hand side of (??). By (??) and lemma ??, the integrand on the right-hand side is

$$\omega^{(4)}(s) \circ (R \otimes R) \circ \Delta'(\Gamma) = \bar{\omega}(s) \circ R(\Gamma) = \bar{\omega}_{\Gamma}^{\text{ren}}(s) ,$$

which therefore gives equation (??). \square

11.2. Group equations with angular dependence. The proof of the group equations essentially only uses (??) together with some formal properties of the map R . As a result the proof of the renormalization group equations also goes through in the case when ϕ has non-trivial angular dependencies.

12. QUADRATIC SUBDIVERGENCES

We treat the general case of graphs in massless ϕ^4 with arbitrary subdivergences.

12.1. Single-scale graphs with quadratic subdivergences. Let G be a connected graph in ϕ^4 theory, and let $q \subsetneq G$ be a connected, 1PI subgraph with $sd(q) = 1$. The crucial feature of quadratic subdivergences is that, by remark ??, they have exactly two 3-valent vertices, and two external edges, or *connectors*, which we denote by $\{e_q, f_q\} \in G \setminus q$. Thus q inherits a unique single-scale structure.

We define a *squashing* of G to be the choice, for every q a 1PI connected quadratic subgraph, of an ordered pair of connectors (e_q, f_q) such that $f_{q_1} \neq f_{q_2}$ for all $q_1 \neq q_2$. It is easy to verify (e.g. as a consequence of corollary ?? below) that such a structure always exists. We define the *squashed* graph by:

$$(222) \quad \bar{G} = G / \cup_q f_q$$

Note that \bar{G} has vertices of arbitrary degree. The image \bar{q} in \bar{G} of a quadratic subgraph q in G can have 0 or 1 connectors.

Definition 48. A *single-scale graph with quadratic subdivergences* is a labelled graph G with a choice of squashing. A *tadpole-free subdivergence* of G is a subgraph $\gamma \subsetneq G$ such that γ is 1PI, divergent and

$$(223) \quad \varepsilon_q(\gamma) = 0 \text{ for all } q \supsetneq \gamma ,$$

where q ranges over the set of quadratic ($sd(q) = 1$) connected 1PI subgraphs of G .

The condition (??) ensures that in the cograph G/γ the connectors of q remain unjoined (no tadpoles are spawned). To construct a Hopf algebra from this, we proceed as before. Let G be a 1-scale graph with quadratic subdivergences and suppose that for every divergent tadpole-free subgraph Γ of G there is a choice of two distinguished connectors giving it a 1-scale structure, in a such a way that for all $\gamma \subset \Gamma$ divergent and tadpole-free, we have

$$(224) \quad \varepsilon_\Gamma(\gamma) = 0 .$$

This guarantees that the cographs Γ/γ have a well-defined single-scale structure.

For any such labelled single-scale graph G , let H_G denote the (loop-number graded) \mathbb{Q} -vector space spanned by its tadpole-free divergent subgraphs and their cographs. We therefore have a well-defined map

$$(225) \quad \begin{aligned} \Delta : H_G &\longrightarrow H_G \otimes_{\mathbb{Q}} H_G \\ \Gamma &\mapsto \sum_{\gamma} \gamma \otimes \Gamma/\gamma \end{aligned}$$

where the sum is over all tadpole-free subdivergences (??). It is well known that this map is coassociative and hence defines a Hopf algebra.

12.2. General massless Feynman rules. With G as above, and γ, Γ in H_G , where γ is not necessarily connected, define differential forms as follows:

$$(226) \quad \begin{aligned} \omega_G &= \left(\prod_{q \subseteq G} \frac{\phi_q}{\psi_q} \right) \frac{\Omega_{\overline{G}}}{\psi_{\overline{G}}^2} , \\ \omega_{\gamma \otimes \Gamma}(s) &= \left(\prod_{q \subseteq \gamma} \frac{\phi_q}{\psi_q} \right) \left(\prod_{q \subseteq \Gamma} \frac{\phi_q}{\psi_q} \right) \frac{s \phi_{\overline{\Gamma}}}{\psi_{\overline{\gamma}} \psi_{\overline{\Gamma}}^2 \Upsilon_{\overline{\gamma; \overline{\Gamma}}}(s)} \Omega_{\overline{\gamma \cup \overline{\Gamma}}} , \end{aligned}$$

where the products are over all connected 1PI quadratically-divergent subgraphs q . Note that the forms are in the edge variables of $\overline{\gamma}, \overline{\Gamma}$, and not γ, Γ . Obviously,

$$(227) \quad \omega_{\gamma \otimes 1}(s) = 0 \quad \text{and} \quad \omega_{1 \otimes \Gamma}(s) = \omega_\Gamma .$$

Define renormalized Feynman rules as follows. Given $\Gamma \in H_G$, write as usual

$$\Delta(\Gamma) = 1 \otimes \Gamma + \sum_{i=1}^n a_i \gamma_i \otimes \Gamma/\gamma_i \in H_G \otimes_{\mathbb{Q}} H_G ,$$

and let $\omega^{\text{ren}} = \omega \circ R$, i.e., define


$$\omega_\Gamma^{\text{ren}} = \omega_\Gamma + \sum_{i=1}^n a_i \omega_{\gamma_i \otimes \Gamma/\gamma_i}(s) .$$

The renormalized Feynman integral is then

$$(228) \quad f_\Gamma(s) = \int_{\sigma_{\overline{\Gamma}}} \omega_\Gamma^{\text{ren}}(s) .$$

We show below that (??) converges and satisfies renormalization group equations (theorem ?? and §??). As previously, we write $f_\Gamma = f_\Gamma(1)$ for the lowest log term.

Example 49. Consider the graphs with nested quadratic subdivergences below.



A squashing of this graph is determined by choosing one of the connectors of the subgraph γ . Let it be edge 1. The squashed graphs $\bar{\Gamma}, \bar{\gamma}, \bar{\Gamma}/\bar{\gamma}$ are obtained from the graphs above by contracting edge 1. The renormalized integral is:

$$f_{\Gamma}(s) = \int_{\sigma_{\bar{\Gamma}}} \omega_{\Gamma}^{\text{ren}}(s) = \int_{\sigma_{\bar{\Gamma}}} \left(\frac{\phi_{\bar{\gamma}} \phi_{\bar{\Gamma}}}{\psi_{\bar{\gamma}} \psi_{\bar{\Gamma}}^3} - \frac{\phi_{\bar{\gamma}} \phi_{\bar{\Gamma}/\bar{\gamma}}^2}{\psi_{\bar{\gamma}}^2 \psi_{\bar{\Gamma}/\bar{\gamma}}^3} \Upsilon_{\bar{\gamma}; \bar{\Gamma}/\bar{\gamma}}(s) \right) \Omega_{\Gamma}$$

where $\Upsilon_{\bar{\gamma}; \bar{\Gamma}/\bar{\gamma}}(s) = \psi_{\bar{\gamma}} \phi_{\bar{\Gamma}/\bar{\gamma}} s + \psi_{\bar{\Gamma}/\bar{\gamma}} \phi_{\bar{\gamma}}$, and all other polynomials are:

$$\begin{aligned} \psi_{\bar{\gamma}} &= \alpha_5 \alpha_6 + \alpha_5 \alpha_7 + \alpha_6 \alpha_7, & \phi_{\bar{\gamma}} &= \alpha_5 \alpha_6 \alpha_7, \\ \psi_{\bar{\Gamma}/\bar{\gamma}} &= \alpha_2 \alpha_3 + \alpha_2 \alpha_4 + \alpha_3 \alpha_4, & \phi_{\bar{\Gamma}/\bar{\gamma}} &= \alpha_2 \alpha_3 \alpha_4, \\ \psi_{\bar{\Gamma}} &= \psi_{\bar{\gamma}} \psi_{\bar{\Gamma}/\bar{\gamma}} + (\alpha_3 + \alpha_4) \alpha_5 \alpha_6 \alpha_7, & \phi_{\bar{\Gamma}} &= \psi_{\bar{\gamma}} \phi_{\bar{\Gamma}/\bar{\gamma}} + \alpha_3 \alpha_4 \alpha_5 \alpha_6 \alpha_7. \end{aligned}$$

Note that in general, the graph polynomial of $\bar{\Gamma}$ is obtained from that of Γ by setting the Schwinger parameters of the squashed edges to 0.

12.3. The single-scale Hopf algebra in the general case. The structure of the Hopf algebra is particularly simple: quadratic subdivergences have no non-trivial overlaps with any other subdivergences and can be separated off.

Lemma 50. *Let $q_1 \subsetneq q_2$ be nested quadratic subdivergences. Then $\varepsilon_{q_2}(q_1) = 0$.*

Proof. Let v_1, v_2 denote the two 3-valent vertices of q_2 and denote its connectors by e_1, e_2 . If $\varepsilon_{q_2}(q_1) = 1$, there exists a path from v_1 to v_2 which is contained in q_1 and so q_1 contains v_1, v_2 . Since q_1 has exactly two 3-valent-vertices (which must be v_1, v_2) and all other vertices of q_1 are 4-valent there can be no edge e of q_2 which does not already lie in q_1 , and therefore $q_1 = q_2$. So $q_1 \subsetneq q_2$ implies that $\varepsilon_{q_2}(q_1) = 0$. \square

Quadratic subdivergences are always tadpole-free.

Lemma 51. *Let I be a connected, divergent 1PI subgraph of G . Then the following two statements are equivalent:*

- (1) *I is tadpole-free (??) ,*
- (2) *For all connected 1PI quadratic subgraphs q of G , one of the following holds:*

$$(229) \quad \begin{cases} (i) & I \cap q = \emptyset, \\ (ii) & I \supseteq q, \\ (iii) & I \subsetneq q \text{ and } \varepsilon_q(I) = 0. \end{cases}$$

Proof. Clearly (2) implies (1). In the other direction, let q be a connected quadratic 1PI subgraph of G . Denote its connectors by $\{e_1, e_2\}$. Since I is 1PI, the intersection $I \cap \{e_1, e_2\}$ consists of 0 or 2 elements. Suppose that $I \cap \{e_1, e_2\} = \emptyset$. Since I is connected, either $I \cap q = \emptyset$ or I is contained in q , and (2) holds. Now suppose that

$e_1, e_2 \in I$. In this case, write I as a two-edge join $I = I_q \cup \{e_1, e_2\} \cup A$, where $I_q = I \cap q$ and A is defined to be the complement of $q \cup \{e_1, e_2\}$ in I . Suppose that I_q is strictly contained in q , for otherwise, $I \supseteq q$ and (2) holds. By power counting,

$$(230) \quad sd(A) + sd(I_q) = sd(I) \geq 0 .$$

If I_q is quadratic, then since I is connected and contains the connectors of e_1, e_2 , $\varepsilon_q(I_q) = 1$, which contradicts lemma ???. Therefore I_q is at most logarithmically divergent which implies by (??) that $sd(A) \geq 0$. Therefore the graph $Q = I \cup q$ is quadratically divergent, since $sd(Q) = sd(A) + sd(q) \geq 1$, and therefore $sd(Q) = 1$. But Q strictly contains I and satisfies $\varepsilon_Q(I) = 1$, which contradicts (1). Therefore we must have had $I_q = q$, which completes the proof. \square

Corollary 52. *Let q_1, q_2 be connected 1PI quadratic subdivergences in G . Then they cannot overlap non-trivially: either $q_1 \cap q_2 = \emptyset$, $q_1 \subseteq q_2$, or $q_2 \subseteq q_1$.*

Proof. This follows immediately from lemmas ??? and ???. \square

One can deduce from this the fact that Δ is coassociative.

12.4. Inflating a squashed graph. The Hopf algebra is phrased in terms of the original graph G , but the Feynman rules are expressed in terms of the squashed graph \overline{G} . To compute the poles of the forms (??) in terms of H_G , we therefore require a correspondence between subgraphs of \overline{G} and certain subgraphs of G .

Definition 53. Let G be as above. If $I \subset \overline{G}$, let Q_I denote the set of connected 1PI quadratic subgraphs $q \subset G$ such that $\varepsilon_{\overline{q}}(I) = 1$. Define I^ℓ to be the smallest subgraph of G which contains I and satisfies

$$(231) \quad |I^\ell \cap \{e_q, f_q\}| \in \{0, 2\} \quad \text{for all } q \in Q_I .$$

It is clear that $\overline{I^\ell} = I$, but note that I^ℓ is not necessarily connected, even if I is. We shall call I^ℓ the *inflation* of the graph $I \subset \overline{G}$.

For divergent graphs, the condition (??) is equivalent to being 1PI.

Lemma 54. *Let $J \subset G$ be divergent. Then*

$$J = (\overline{J})^\ell \quad \Leftrightarrow \quad J \text{ is 1PI} .$$

Proof. Suppose that $J \subset G$ is divergent but not 1PI. Then there is an edge $e \in J$ such that $J \setminus e$ has two components, A and B . Since $sd(J) \geq 0$, it follows that at least one component, say A , is quadratic. We can assume that A is 1PI connected. Necessarily $\varepsilon_A(J) = 1$, but J only contains one connector of A , namely the edge e , which violates (??). Therefore if J is of the form I^ℓ for some I , then it is 1PI. Conversely, suppose that J is 1PI. Then J contains \overline{J} and is the minimal such graph satisfying condition (??), since if there is a quadratic subgraph q such that the intersection $J \cap \{e_q, f_q\}$ has exactly one element e , then $J \setminus e$ has one more component than J . \square

Remark 55. *Suppose that $Q \subset G$ is a connected 1PI quadratic subgraph such that $\varepsilon_Q(I^\ell) = 1$ but $I^\ell \cap \{e_Q, f_Q\} = \emptyset$. If I^ℓ is not contained in Q then I^ℓ has at least two components: $I^\ell = I_Q^\ell \cup I_{Q^c}^\ell$ where $I_Q^\ell = I^\ell \cap Q$ and $I_{Q^c}^\ell \subset G \setminus (Q \cup \{e_Q, f_Q\})$. By repeating this for all such subgraphs Q , we obtain a decomposition of I^ℓ into disjoint (but not necessarily connected) pieces $I^\ell = \cup_i I^{\ell, i}$.*

Lemma 56. *For any $I \subset \overline{G}$, and $A \subset G$ a single-scale subgraph, we have $\varepsilon_{\overline{A}}(I) = \varepsilon_{\overline{A}^\ell}(I^\ell)$. Let $Q, I_Q^\ell, I_{Q^c}^\ell$ be as in remark ???. Then for any connected quadratic 1PI subgraph q of G we have*

$$(232) \quad \varepsilon_q(I^\ell) = \varepsilon_q(I_Q^\ell) + \varepsilon_q(I_{Q^c}^\ell) .$$

Proof. For the first part, it is obvious that $\varepsilon_A(I^\ell) = 1$ implies that $\varepsilon_{\overline{A}}(I) = 1$. In the opposite direction, suppose that $\varepsilon_{\overline{A}}(I) = 1$. Then there exists a path γ connecting the two distinguished vertices of \overline{A} . If q is quadratic such that $e_q \in \overline{A}$ and such that $\varepsilon_{\overline{q}}(\gamma \cap \overline{q}) = 1$ then we must have $e_q \in \gamma$. Hence $\gamma \cup \{e_q, f_q\}$ is a path in $I \cup \{f_q\}$ which connects the distinguished vertices of $A \cup \{f_q\}$. Continuing by induction it follows that $\gamma^\ell \subset I^\ell$ connects the distinguished vertices of A^ℓ .

For the second part, we know by corollary ??? that either $q \cap Q = \emptyset$, $Q \subsetneq q$, or $q \subseteq Q$. In the first two cases, clearly $\varepsilon_q(I_Q^\ell) = 0$ and we have $\varepsilon_q(I^\ell) = \varepsilon_q(I_{Q^c}^\ell)$. In the third case we have $\varepsilon_q(I_{Q^c}^\ell) = 0$ and $\varepsilon_q(I^\ell) = \varepsilon_q(I_Q^\ell)$. \square

12.5. Location of the poles. We compute the orders of the poles of the forms (??) along the exceptional divisors \mathcal{E}_I where $I \subset E(\overline{G})$, and relate this to I^ℓ .

Lemma 57. *Let G be as above, and let I be a connected 1PI subset of edges of \overline{G} . Let F be a flag of (tadpole-free) divergent subgraphs of G , and let γ_F be defined by (??). Then $\omega_{\gamma_F \otimes G / \gamma_F}(s)$ has a pole along \mathcal{E}_I of order $1 + p(I)$, where*

$$(233) \quad p(I) = 2h_{I_{\overline{\gamma}_F}} + 2h_{I_{\overline{G}/\overline{\gamma}_F}} - |I| - \varepsilon_{\overline{\gamma}_F \otimes \overline{G} / \overline{\gamma}_F}(I) - \sum_q \varepsilon_{\overline{q}}(I) ,$$

and where the sum is over all connected 1PI quadratic subgraphs q of G .

Proof. It follows from (??) and the definition of ε that

$$(234) \quad v_I \left(\prod_q \frac{\phi_{\overline{q}}}{\psi_{\overline{q}}} \right) = \sum_q \varepsilon_{\overline{q}}(I) .$$

By the calculations in §???, the valuation along L_I is given (cf (??)) by

$$-v_I \left(\frac{s \phi_{\overline{\Gamma}/\overline{\gamma}_F}}{\psi_{\overline{\gamma}_F} \psi_{\overline{\Gamma}/\overline{\gamma}_F}^2 \Upsilon_{\overline{\gamma}_F, \overline{\Gamma}/\overline{\gamma}_F}(s)} \Omega_{\overline{\Gamma}} \right) = 2h_{I_{\overline{\gamma}_F}} + 2h_{I_{\overline{G}/\overline{\gamma}_F}} - \varepsilon_{\overline{\gamma}_F \otimes \overline{\Gamma} / \overline{\gamma}_F}(I) .$$

Using the fact (corollary ??) that there is a bijection between quadratic connected 1PI subgraphs of G and those of $\gamma_F, G/\gamma_F$, the result follows from the two previous expressions and (??). \square

Lemma 58. *Let G, I, F, γ be as in the previous lemma. Maximally decompose I^ℓ into components $I^{\ell,1} \cup \dots \cup I^{\ell,m}$ according to remark ???. Then*

$$(235) \quad p(I) = \sum_{i=1}^m \left(sd(I_{\gamma_F}^{\ell,i}) + sd(I_{G/\gamma_F}^{\ell,i}) - \sum_{I^{\ell,i} \subseteq q} \varepsilon_q(I^{\ell,i}) \right) .$$

In particular, $\omega_{\gamma_F \otimes G / \gamma_F}(s)$ has at most simple poles along exceptional divisors \mathcal{E}_I indexed by I such that $I_{\gamma_F}^\ell, I_{G/\gamma_F}^\ell$ are divergent, tadpole-free, and $I_{G/\gamma_F}^\ell \subsetneq G/\gamma_F$.

Proof. By lemmas ?? and ??, we can write

$$p(I) = 2h_{I_{\overline{\gamma}_F}} + 2h_{I_{\overline{G}/\overline{\gamma}_F}} - |I| - \varepsilon_{\gamma_F \otimes G / \gamma_F}(I^\ell) - \sum_q \varepsilon_q(I^\ell) .$$

By definition of the inflation map, we have

$$|I^{\ell,i}| - |\overline{I^{\ell,i}}| = \sum_{I^{\ell,i} \not\subseteq q} \varepsilon_q(I^{\ell,i}).$$

Since inflation does not change the number of loops, we have by (??):

$$\begin{aligned} p(I) &= \sum_i \left(2h_{I_{\gamma_F}^{\ell,i}} + 2h_{I_{G/\gamma_F}^{\ell,i}} - |I^{\ell,i}| - \sum_{I^{\ell,i} \subseteq q} \varepsilon_q(I^{\ell,i}) \right) - \varepsilon_{\gamma_F \otimes G/\gamma_F}(I^\ell) \\ &= \sum_i \left(sd(I_{\gamma_F}^{\ell,i}) + sd(I_{G/\gamma_F}^{\ell,i}) - \sum_{I^{\ell,i} \subseteq q} \varepsilon_q(I^{\ell,i}) \right) - \varepsilon_{\gamma_F \otimes G/\gamma_F}(I^\ell). \end{aligned}$$

Now observe that every term

$$sd(I_{\gamma_F}^{\ell,i}) + sd(I_{G/\gamma_F}^{\ell,i}) - \sum_{I^{\ell,i} \subseteq q} \varepsilon_q(I^{\ell,i})$$

is ≤ 0 . If $sd(I_{\gamma_F}^{\ell,i}) = sd(I_{G/\gamma_F}^{\ell,i}) = 0$ this is obvious, and one verifies that if $\{sd(I_{\gamma_F}^{\ell,i}), sd(I_{G/\gamma_F}^{\ell,i})\} = \{0, 1\}$ then in each case, $I^{\ell,i}$ is a quadratic 1PI graph, and $\varepsilon_q(I^{\ell,i}) = 1$ for $q = I^{\ell,i}$. One cannot have $sd(I_{\gamma_F}^{\ell,i}) = sd(I_{G/\gamma_F}^{\ell,i}) = 1$. Therefore the pole of $\omega_{\gamma_F \otimes G/\gamma_F}(s)$ along \mathcal{E}_I is at most simple, and this only happens when every $I_{\gamma_F}^{\ell,i}, I_{G/\gamma_F}^{\ell,i}$ are divergent, $\sum_{I^{\ell,i} \subseteq q} \varepsilon_q(I^{\ell,i}) = 0$, and $\varepsilon_{\gamma_F \otimes G/\gamma_F}(I^\ell) = 0$. In other words, I^ℓ is a union of tadpole-free graphs, and by assumption (??), $\varepsilon_{\gamma_F \otimes G/\gamma_F}(I^\ell)$ vanishes in this case provided that $I_{G/\gamma_F} \subsetneq G/\gamma_F$. As in the proof of corollary ??, there is no pole when $I_{G/\gamma_F} = G/\gamma_F$. Therefore we have shown that the poles are simple, and are in one-to-one correspondence with the set of pairs of tadpole-free divergent subgraphs $I_{\gamma_F}^\ell \subset \gamma_F$ and $I_{G/\gamma_F}^\ell \subsetneq G/\gamma_F$. \square

Since by lemma ??, divergent inflated subgraphs of G are the same as 1PI divergent subgraphs of G , the poles of ω_G are indeed indexed by 1PI divergent tadpole-free subgraphs of G , and hence by the terms in the definition of the coproduct.

12.6. Residues and proof of convergence.

Proposition 59. *Let F be a flag of divergent subgraphs in G , and let I be a strict subset of edges in \overline{G} . The form $\omega_{\gamma_F \otimes G/\gamma_F}(s)$ has a simple pole along the exceptional divisor \mathcal{E}_I if and only if $I_{\gamma_F}^\ell, I_{G/\gamma_F}^\ell$ are divergent 1PI and tadpole-free, and $I_{G/\gamma_F}^\ell \subsetneq G/\gamma_F$. Then the residue is*

$$(236) \quad \text{Res}_{\mathcal{E}_I} \omega_{\gamma_F \otimes G/\gamma_F}(s) = \omega_{I_{\gamma_F}^\ell \cup I_{G/\gamma_F}^\ell} \otimes \omega_{\gamma_F/I^\ell \otimes G/(\gamma_F \cup I^\ell)}(s).$$

Proof. Let q be any 1PI connected quadratic subgraph of G . It follows from (??) and (??) that to leading order in the I -variables:

$$\frac{\phi_{\overline{q}}}{\psi_{\overline{q}}} \longrightarrow \begin{cases} \frac{\phi_{\overline{q}}}{\psi_{\overline{q}}} \otimes 1 & \text{if } q \subseteq I^\ell \\ 1 \otimes \frac{\phi_{\overline{q}}}{\psi_{\overline{q}}} & \text{if } q \cap I^\ell = \emptyset \\ 1 \otimes \frac{\phi_{\overline{q}/I}}{\psi_{\overline{q}/I}} & \text{if } I^\ell \subsetneq q \text{ and } \varepsilon_q(I^\ell) = 0 \end{cases}$$

where, following lemma ??, the left hand side of the tensor corresponds to the I variables, and the right-hand side to the I^c variables. It follows that

$$(237) \quad \prod_{q \subseteq G} \frac{\phi_{\overline{q}}}{\psi_{\overline{q}}} \longrightarrow \left(\prod_{q \subseteq I^\ell} \frac{\phi_{\overline{q}}}{\psi_{\overline{q}}} \right) \otimes \left(\prod_{q \subseteq G/I^\ell} \frac{\phi_{\overline{q}}}{\psi_{\overline{q}}} \right)$$

after applying lemma ?? to every connected component of I^ℓ , since every quadratic subdivergence q corresponds to one of the three cases of (??), and there is a bijection between 1PI connected quadratic subdivergences of G and those of I^ℓ and G/I^ℓ . Now apply (??) to the definition (??). The result follows by an identical computation to proposition ??. \square

By the standard properties of the coproduct, and the argument given in proposition ??, for any flag F of divergent subgraphs of Γ , the total residue is

$$(238) \quad \text{Res} \omega_{\gamma_F \otimes \Gamma / \gamma_F}(s) = (\omega^1 \otimes \omega^{23}) \circ \mu_{13}(\Delta \otimes \Delta)(\gamma_F \otimes \Gamma / \gamma_F) ,$$

where $(\omega^1 \otimes \omega^{23})(x \otimes y \otimes z) = \omega_x \otimes \omega_{y \otimes z}(s)$. Note that in this formula, the residue corresponding to a term I is the residue along the divisor indexed by the squashed graph \bar{I} . By (??) and (??), the forms (??) satisfy our hypotheses (??) for renormalization. Since the denominators $\psi_{\bar{q}}$ of the quadratic correction factors are polynomials with positive coefficients, we conclude that:

Theorem 60. *The form $\omega_{\Gamma}^{\text{ren}}(s)$ has no poles along any exceptional divisors \mathcal{E}_I , and therefore the renormalized integral (??) is convergent.*

The proof is similar to the proof of theorem ??.

12.7. Renormalization group equations: quadratic case. The proof of the group equations is essentially identical to §??. We summarize the main steps. For any graph $\gamma \in H_G$ define the quadratic correction factor by

$$(239) \quad Q_\gamma = \prod_{q \subseteq \gamma} \frac{\phi_{\bar{q}}}{\psi_{\bar{q}}}$$

where the product is over all 1PI connected quadratic subgraphs $q \subseteq \gamma$. In addition to the data of the forms (??), define for any $\lambda > 0$,

$$(240) \quad \nu_{\gamma \otimes \Gamma}(s) = Q_\gamma Q_\Gamma \frac{\phi_\Gamma}{\psi_\Gamma} \frac{\lambda}{(\psi_{\gamma \cup \Gamma} \lambda + \phi_{\gamma \cup \Gamma})^2} \prod_{e \in E(\gamma) \cup E(\Gamma)} d\alpha_e ,$$

$$(241) \quad \omega_{\gamma_1 \otimes \gamma_2 \otimes \gamma_3 \otimes \gamma_4}^{(4)}(s) = Q_{\gamma_1} Q_{\gamma_2} Q_{\gamma_3} Q_{\gamma_4} \frac{\phi_{\gamma_2} \phi_{\gamma_4}}{\psi_{\gamma_2} \psi_{\gamma_4}} \frac{\Omega_{\gamma_1 \cup \dots \cup \gamma_4}}{\Upsilon_{\gamma_1 \cup \gamma_2 \cup \gamma_3; \gamma_4}^2}(s) .$$

and extend by linearity to $H \otimes_{\mathbb{Q}} H$ and $H^{\otimes 4}$ respectively. The proof is now identical to the log-divergent case since the correction terms (??) completely factor out of every equation. In brief: the forms $\nu^{\text{ren}}(\lambda)$ are pole-free along the \mathcal{E}_I , and we have

$$\int_{\sigma_{\bar{\mathcal{C}}}} \omega_G^{\text{ren}}(1) = \int_{[0, \infty]^{E(\bar{\mathcal{C}})}} \nu_G^{\text{ren}}(\lambda) .$$

as before. The proof is the same as lemma ??. The analogue of proposition ?? is

$$\int_{\sigma_{\bar{\mathcal{C}}_1}} \omega_{G_1}^{\text{ren}}(1) \times \int_{\sigma_{\bar{\mathcal{C}}_2}} \omega_{G_2}^{\text{ren}}(s) = \int_{\sigma_{\bar{\mathcal{C}}_1 \cup \bar{\mathcal{C}}_2}} \omega_{R(G_1) \otimes R(G_2)}^{(4)}(s)$$

Finally, it is clear from the definition and the property $Q_{\gamma_1 \cup \gamma_2} = Q_{\gamma_1} Q_{\gamma_2}$ that

$$\sum_{1 \leq i \leq n} \omega_{a_1 \dots a_{i-1} \otimes a_i \otimes a_{i+1} \dots a_n}^{(4)}(s) = s \frac{\partial}{\partial s} \omega_{a_1 \dots a_n}(s) ,$$

as in (??). Thus the proof of proposition ?? goes through as before and we obtain

Theorem 61. *The Feynman rules (??) define a cocharacter on H_G . Equivalently, the renormalization group equations hold for graphs with quadratic subdivergences:*

$$\int_{\sigma_{\overline{\Gamma}}} s \frac{\partial}{\partial s} \omega_{\overline{\Gamma}}^{\text{ren}}(s) = \sum_{\gamma \subset \overline{\Gamma}} \int_{\sigma_{\overline{\gamma}}} \omega_{\overline{\gamma}}^{\text{ren}}(1) \times \int_{\sigma_{\overline{\Gamma}/\overline{\gamma}}} \omega_{\overline{\Gamma}/\overline{\gamma}}^{\text{ren}}(s) .$$

for all $\Gamma \in H_G$.

The first part of the theorem follows from the second, by proposition ??.

Remark 62. *These are still the renormalization group equations for massless diagrams. In the presence of masses the result Thm. ?? is a polynomial in squared masses. Repeating the analysis term by term for the coefficient functions verifies the full Callan-Symanzik equations, as expected.*

12.8. The tangent motive in the general case. The definition of the mixed Hodge structure corresponding to G in the general case is similar to the case when G has at most logarithmic subdivergences, if we include extra hypersurfaces X_q for each quadratic 1PI connected subgraph q in the definition of $X_{G,s}^{\text{tot}}$. Then only the differential form $\omega_G^{\text{ren}}(s)$ changes. Thus we can define $\text{Mot}(G)$ via (??), define framings in the same manner, and conclude that f_G is its period, in all cases.

13. REMINDERS ON HOPF ALGEBRAS

13.1. Basic definitions. Consider any commutative, graded Hopf algebra

$$H = \bigoplus_{n \geq 0} H_n$$

over a field k of characteristic zero, where $H_0 = k$. Denote the multiplication by $\mu_2 : H \otimes_k H \rightarrow H$, and more generally let $\mu_n : H^{\otimes n} \rightarrow H$ denote the multiplication of n elements. We frequently write $y_1 \dots y_n$ as a shorthand for $\mu_n(y_1 \otimes \dots \otimes y_n)$ for simplicity. If the coproduct is denoted $\Delta : H \rightarrow H \otimes_k H$, the reduced coproduct is defined by Δ' , where $\Delta' = \Delta - 1 \otimes \text{id} - \text{id} \otimes 1$. It satisfies

$$(242) \quad \Delta'(H_n) \subseteq \bigoplus_{p+q=n, p \geq 1, q \geq 1} H_p \otimes_k H_q .$$

For $n \geq 1$, consider the maps

$$\Delta^{(n)} : H \rightarrow H^{\otimes n+1}$$

obtained by setting $\Delta^{(1)} = \Delta'$ and iterating the reduced coproduct:

$$\Delta^{(n)} = (\text{id} \otimes \Delta^{(n-1)}) \circ \Delta' = (\Delta^{(n-1)} \otimes \text{id}) \circ \Delta' \quad \text{for } n \geq 2 ,$$

The maps $\Delta^{(n)}$ are well-defined by the coassociativity of Δ . For any element $x \in H$, we shall sometimes use the following version of Sweedler's notation and write

$$(243) \quad \Delta^{(n)}(x) = \sum_{(x)} x^{(1)} \otimes \dots \otimes x^{(n+1)} ,$$

where by (??), all elements $x^{(i)}$ have degree ≥ 1 .

Definition 63. The *coradical filtration* is the increasing filtration defined by

$$H^{(i)} = \{x : \Delta^{(i)}x = 0\}$$

where $i \geq 1$, and $H^{(0)} = k$. The set of *primitive* elements in H are the elements in $H^{(1)}$, i.e., which satisfy $\Delta x = 1 \otimes x + x \otimes 1$.

The Hopf algebras we consider in this paper (Hopf algebras of graphs or trees) are graded with respect to the coradical filtration, i.e., Δ is homogeneous with respect to the grading associated to the coradical filtration.

13.2. The preparation map.

Definition 64. The *preparation map* $R : H \rightarrow H \otimes_k H$ is defined by

$$(244) \quad R = 1 \otimes id + \sum_{n \geq 1} (-1)^n (\mu_n \otimes id) \Delta^{(n)} ,$$

where $\mu_1 = id$. It is well-defined since by (??) the sum on the right-hand side terminates when applied to any element in H .

It is convenient to set $R^0 = 1 \otimes id$, and $R^n = (-1)^n (\mu_n \otimes id) \Delta^{(n)}$ for all $n \geq 1$. By Sweedler's notation (??), we can write for $n \geq 1$ and $x \in H$,

$$(245) \quad R^n(x) = (-1)^n \sum_{(x)} x^{(1)} \dots x^{(n)} \otimes x^{(n+1)} .$$

One can also define R recursively in the following way.

Lemma 65. *Using Sweedler's notation (??), the map R satisfies*

$$(246) \quad R(x) = 1 \otimes x - \sum_{(x)} \mu(R(x^{(1)})) \otimes x^{(2)} .$$

Proof. This follows immediately from $R^n = -(\mu \circ R^{n-1} \otimes id) \circ \Delta'$. \square

Since H is commutative and graded, the antipode $S : H \rightarrow H$ is $S = -\mu \circ R$.

13.3. Renormalization property of R . Consider the map defined by:

$$(247) \quad \begin{aligned} \mu_{13} : H^{\otimes 4} &\longrightarrow H^{\otimes 3} \\ x_1 \otimes x_2 \otimes x_3 \otimes x_4 &\mapsto x_1 x_3 \otimes x_2 \otimes x_4 \end{aligned}$$

The following theorem is the main mechanism for the cancellation of poles in renormalization. For want of a suitable reference, we give a complete proof here.

Theorem 66. *The preparation map satisfies the following equation:*

$$(248) \quad \mu_{13} \circ (\Delta \otimes (\Delta - id \otimes 1)) \circ R = 1 \otimes R .$$

Proof. By definition (??), the left-hand side of (??) is given by

$$\sum_{n \geq 0} (-1)^n \mu_{13} (\Delta \mu_n \otimes (\Delta - id \otimes 1)) \circ \Delta^{(n)} ,$$

where $\mu_0 = id$ and $\Delta^{(0)} = 1 \otimes id$. Since Δ is a homomorphism, this can be rewritten:

$$(249) \quad \sum_{n \geq 0} (-1)^n \tilde{\mu}_n (\Delta^{\otimes n} \otimes (\Delta - id \otimes 1)) \circ \Delta^{(n)} ,$$

where $\tilde{\mu}_n : H^{\otimes 2n} \rightarrow H^{\otimes 3}$ is the map defined by

$$\tilde{\mu}_n(y_1 \otimes \dots \otimes y_{2n}) = y_1 y_3 \dots y_{2n-1} \otimes y_2 y_4 \dots y_{2n-2} \otimes y_{2n} .$$

Now let $x \in H$, and write out the following terms

$$(250) \quad \begin{aligned} (\Delta^{\otimes n} \otimes (\Delta - id \otimes 1)) \circ \Delta^{(n)}(x) &= \\ ((\Delta' + id \otimes 1 + 1 \otimes id)^{\otimes n} \otimes (\Delta' + 1 \otimes id)) \circ \Delta^{(n)}(x) \end{aligned}$$

using Sweedler's notation (??). When $n = 0$, (??) gives

$$(251) \quad 1 \otimes 1 \otimes 1 \otimes x + \sum_{(x)} 1 \otimes 1 \otimes x^{(1)} \otimes x^{(2)},$$

and when $n = 1$ (??) has the following six terms:

$$(252) \quad \sum_{(x)} (1 \otimes x^{(1)} \otimes 1 \otimes x^{(2)} + 1 \otimes x^{(1)} \otimes x^{(2)} \otimes x^{(3)} + x^{(1)} \otimes 1 \otimes 1 \otimes x^{(2)} + \\ x^{(1)} \otimes 1 \otimes x^{(2)} \otimes x^{(3)} + x^{(1)} \otimes x^{(2)} \otimes 1 \otimes x^{(3)} + x^{(1)} \otimes x^{(2)} \otimes x^{(3)} \otimes x^{(4)})$$

The general expression can be encoded by an alphabet with three letters a, b, c , which correspond respectively to the maps $1 \otimes id$, $id \otimes 1$, and Δ' . For any word in the letters $\{a, b, c\}$ of length n not ending in b , we associate a term in (??). For this, consider the unique morphism (for the concatenation product) of monoids

$$\phi_1 : \{a, b, c\}^\times \rightarrow \{1, x\}^\times$$

such that $\phi_1(a) = 1x$, $\phi_1(b) = x1$, and $\phi_1(c) = xx$. Now let ϕ_2 be the map which inserts a tensor in between all letters of a word in the letters $\{1, x\}$, and adds superscripts to all letters x in strictly increasing order. Finally, if $w \in \{a, b, c\}^\times$ is a word with at least 2 letters, set $\phi(w) = \phi_2 \circ \phi_1(w)$. We set $\phi(a) = 1 \otimes 1 \otimes 1 \otimes x_1$ and $\phi(c) = 1 \otimes 1 \otimes x^{(1)} \otimes x^{(2)}$ to agree with (??). Hence for $n \geq 2$ we have

$$(253) \quad \phi(a^n) = 1 \otimes x^{(1)} \otimes 1 \otimes x^{(2)} \otimes \dots \otimes 1 \otimes x^{(n)}$$

and the six terms of (??) are $\phi(aa)$, $\phi(ac)$, $\phi(ba)$, $\phi(bc)$, $\phi(ca)$, $\phi(cc)$ in order. With these definitions, the full expansion of (??) in Sweedler notation is just

$$\phi((a + b + c)^n(a + c)).$$

Therefore consider the non-commutative formal power series

$$T = \sum_{n \geq 0} (-1)^n (a + b + c)^n (a + c) \in \mathbb{Q}\langle\langle a, b, c \rangle\rangle.$$

We have shown that (??) is $\tilde{\mu} \circ \phi(T)$. Now it is easy to verify from the definitions that for all (possibly empty) words $w, w' \in \{a, b, c\}^\times$,

$$\tilde{\mu} \circ \phi(w c w') = \tilde{\mu} \circ \phi(w b a w').$$

It follows that $\tilde{\mu} \circ \phi(T) = \tilde{\mu} \circ \phi(\overline{T})$, where

$$\overline{T} = \sum_{n \geq 0} (-1)^n (a + b + ba)^n (a + ba) \in \mathbb{Q}\langle\langle a, b \rangle\rangle.$$

This satisfies $(1 + a + b + ba)\overline{T} = (a + ba)$, which is $(1 + b)(1 + a)\overline{T} = (1 + b)a$. Since $1 + b$ (resp. $1 + a$) is invertible as a non-commutative formal power series in $\mathbb{Q}\langle\langle a, b \rangle\rangle$, this implies that $(1 + a)\overline{T} = a$, which has the unique solution

$$\overline{T} = \sum_{n \geq 1} (-1)^{n+1} a^n.$$

Therefore, by (??), $\tilde{\mu} \circ \phi(\overline{T})$ reduces to

$$\sum_{n \geq 0} (-1)^n \tilde{\mu} \circ \phi(a^{n+1}) = 1 \otimes 1 \otimes x + \sum_{n \geq 1} \sum_{(x)} (-1)^n 1 \otimes x^{(1)} \dots x^{(n)} \otimes x^{(n+1)}$$

which is exactly $1 \otimes R(x)$ by (??). Since H is graded commutative, all the previous formal power series arguments are in fact finite series when evaluated on a specific element x of H , and so this proves that (??) is equal to $1 \otimes R$. \square

13.4. Properties for the renormalization group equations.

Lemma 67. *For any $m, n \geq 0$, the following diagram commutes:*

$$(254) \quad \begin{array}{ccc} H & \xrightarrow{\Delta'} & H^{\otimes 2} \\ \downarrow R^{m+n+1} & & \downarrow R^m \otimes R^n \\ H^{\otimes 2} & \xleftarrow{-\mu_3 \otimes id} & H^{\otimes 4} \end{array}$$

Proof. In Sweedler's notation (??), we have:

$$R^{m+n+1}(x) = \sum_{(x)} (-1)^{m+n+1} x^{(1)} \dots x^{(m+n+1)} \otimes x^{(m+n+2)} .$$

On the other hand, $(R^m \otimes R^n) \circ \Delta'(x)$ can be written

$$\sum_{(x)} (-1)^{m+n} x^{(1)} \dots x^{(m)} \otimes x^{(m+1)} \otimes x^{(m+2)} \dots x^{(m+n+1)} \otimes x^{(m+n+2)} ,$$

which is mapped to $-R^{m+n+1}(x)$ on applying $\mu_3 \otimes id$. \square

We also need the following characterization of additive cocharacters on H . Let K be a field containing k , and let $K[L]$ be the Hopf algebra of K -valued functions on the additive group \mathbb{G}_a , where L is primitive. If $g \in K[L]$, write $g'(L)$ (resp. $g''(L)$), for the usual derivative (resp. second derivative) with respect to L . Since H and $K[L]$ are connected, they have counits (or augmentations) which we denote by $\varepsilon : H \rightarrow H_0 \cong k$ and $\varepsilon : K[L] \rightarrow K$.

Proposition 68. *Suppose that H is graded for its coradical filtration. Then a homomorphism $f : H \rightarrow K[L]$ of augmented algebras is a homomorphism of Hopf algebras if and only if*

$$(255) \quad f_x''(L) = \sum_{(x)} f'_{x^{(1)}}(0) f'_{x^{(2)}}(L) ,$$

where $\Delta'(x) = \sum_{(x)} x^{(1)} \otimes x^{(2)}$, and the dependence on x is written in subscript.

Proof. Let us denote the Lie coalgebras of H and $K[L]$ by

$$\mathfrak{h} = \frac{H_{>0}}{H_{>0}H_{>0}} \quad \text{and} \quad KL \cong \frac{K[L]_{>0}}{K[L]_{>0}K[L]_{>0}}$$

respectively, and let $\bar{f} : \mathfrak{h} \rightarrow KL$ be the map induced by f . One verifies using the fact that H is graded for its coradical filtration, that the map f is a homomorphism of Hopf algebras if and only if the following diagram commutes:

$$\begin{array}{ccc} H & \longrightarrow & \mathfrak{h} \otimes_k H \\ \downarrow f & & \downarrow \bar{f} \otimes f \\ K[L] & \longrightarrow & KL \otimes_K K[L] \end{array}$$

where both horizontal arrows are given by the infinitesimal coaction $(\pi \otimes id) \circ \Delta'$, and $\pi : H_{>0} \rightarrow \mathfrak{h}$ (resp. $\pi : K[L]_{>0} \rightarrow KL$) are the natural quotient maps. The commutativity of the diagram is therefore equivalent to the equation

$$(256) \quad (\pi \otimes id) \circ \Delta' f_x = (\bar{f} \otimes f) \circ (\pi \otimes id) \circ \Delta'(x)$$

for all $x \in H$. Using the fact that $\Delta'(L^n) = \sum_{1 \leq i < n} \binom{n}{i} L^i \otimes L^{n-i}$, we see that

$$(\pi \otimes id) \circ \Delta'(g(L)) = L \otimes (g'(L) - g'(0)) \quad \text{for all } g(L) \in K[L],$$

since the formula is linear and evidently true for $g(L) = L^n$. Now let $x \in H$ satisfy $\varepsilon(x) = 0$, where ε is the counit. Using the Sweedler notation (??), and applying the previous equation to the left-hand side of (??), we obtain

$$(257) \quad L \otimes (f'_x(L) - f'_x(0)) = \sum_{(x)} \bar{f}_{\pi(x^{(1)})}(L) \otimes f_{x^{(2)}}(L).$$

As $\bar{f}_{\pi(x^{(1)})} = \pi(f_{x^{(1)}}(L))$ is the linear term $f'_{x^{(1)}}(0)L$ of $f_{x^{(1)}}(L)$, (??) becomes

$$(258) \quad f'_x(L) - f'_x(0) = \sum_{(x)} f'_{x^{(1)}}(0) f_{x^{(2)}}(L).$$

Since f respects the counit, $f_y(L)$ has no constant term in L for any y of degree ≥ 1 , and so $f_y(0) = 0$. Thus the constant terms on both sides of (??) vanish, and (??) is equivalent to its derivative with respect to L , which is simply (??). \square

In the applications, H is a Hopf algebra of graphs over $k = \mathbb{Q}$, the coefficient field K is \mathbb{C} or \mathbb{R} , and $L = \log s$, where s is the chosen renormalization scale.

REFERENCES

- [1] **C. Bergbauer, D. Kreimer:** *Hopf algebras in renormalization theory: Locality and Dyson-Schwinger equations from Hochschild cohomology*, IRMA Lect.Math.Theor.Phys.10 (2006), 133-164.
- [2] **S. Bloch, H. Esnault, D. Kreimer:** *On motives associated to graph polynomials*, Comm. Math. Phys. 267 (2006), no. 1, 181-225.
- [3] **S. Bloch, D. Kreimer:** *Feynman amplitudes and Landau singularities for 1-loop graphs.*, Comm. Number Th. Phys. 4 (2011), no. 4, 709-753.
- [4] **F. Brown:** *The massless higher-loop two-point function*, Comm. in Math. Physics 287, no. 3, 925-958 (2009).
- [5] **F. Brown:** *On the Periods of some Feynman Integrals*, arXiv:0910.0114 [math.AG] (2009).
- [6] **G. Kirchhoff:** *Ueber die Auflösung der Gleichungen, auf welche man bei der Untersuchung der linearen Vertheilung galvanischer Ströme geführt wird*, Annalen der Physik und Chemie 72 no. 12 (1847), 497-508.

INST.MATH.JUSSIEU, 175 RUE DE CHEVALERET, 75013 PARIS, FRANCE AND HUMBOLDT U.,
UNTER DEN LINDEN 6, 10099 BERLIN, GERMANY